

**PROBABILITY: LIMIT THEOREMS II, SPRING 2020.
HOMEWORK PROBLEMS**

PROF. YURI BAKHTIN

Instructions. You are allowed to work on solutions in groups, but you are required to write up solutions on your own. Please give complete solutions, all claims need to be justified. Late homework will not be accepted. Please let me know if you find any misprints or mistakes.

1. DUE BY WEDNESDAY FEB 26, 11:00AM

1. Let $(X_n)_{n \in \mathbb{N}}$ be an i.i.d. positive sequence, and $S_n = X_1 + \dots + X_n$. Let $N_t = \sup\{n : S_n \leq t\}$. Prove that $S_t = S_{[t]}$ and $(N_t)_{t \in \mathbb{R}_+}$ are stochastic processes.
2. Let $v_1, v_2 : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be C^∞ vector fields on \mathbb{R}^2 . Let N_t be defined as in the previous problem. Let $i(n) = 1$ if n is odd and $i(n) = 2$ if n is even. Let Z_t denote the solution of the ODE

$$\frac{d}{dt}Z_t = v_{i(N_t)}(Z_t), \quad Z_0 = x.$$

Prove that $(Z_t)_{t \geq 0}$ is a stochastic process.

3. Prove that cylinders $C(t_1, \dots, t_n, B)$, $B \in \mathcal{B}(\mathbb{R}^n)$, $t_1, \dots, t_n \geq 0$, $n \in \mathbb{N}$ form an algebra.
4. Prove that elementary cylinders $C(t_1, \dots, t_n, B_1 \times \dots \times B_n)$, $B_1, \dots, B_n \in \mathcal{B}(\mathbb{R})$, $t_1, \dots, t_n \geq 0$, $n \in \mathbb{N}$ form a semi-ring \mathcal{C} (i.e., (i) $\emptyset \in \mathcal{C}$, (ii) if $A, B \in \mathcal{C}$, then $A \cap B \in \mathcal{C}$, (iii) if $A, B \in \mathcal{C}$, then there is n and disjoint $C_1, \dots, C_n \in \mathcal{C}$ such that $A \setminus B = C_1 \cup \dots \cup C_n$). [*A reminder that does not help to solve the problem but just fyi: when extending measures onto the entire σ -algebra one can start with an algebra or just with a semi-ring.*]
5. We can define the cylindrical σ -algebra as the σ -algebra generated by elementary cylinders or by cylinders. Prove that these definitions are equivalent.
6. Let $\mathcal{F}_T = \sigma\{C(t_1, \dots, t_n, B) : t_1, \dots, t_n \in T\}$ for $T \subset \mathbb{T}$. Prove that

$$\mathcal{B}(\mathbb{R}^{\mathbb{T}}) = \bigcup_{\text{countable } T \subset \mathbb{T}} \mathcal{F}_T.$$

7. Prove that $X : \mathbb{T} \times \Omega \rightarrow \mathbb{R}$ is a stochastic process iff X seen as $X : \Omega \rightarrow \mathbb{R}^{\mathbb{T}}$ is $(\Omega, \mathcal{F}) \rightarrow (\mathbb{R}^{\mathbb{T}}, \mathcal{B}(\mathbb{R}^{\mathbb{T}}))$ measurable.
8. Use characteristic functions to prove the existence of a Wiener process (up to continuity of paths).

9. Let $(X_t)_{t \in [0,1]}$ be an (uncountable) family of i.i.d. r.v.'s with nondegenerate distribution. Prove that no modification of this process can be continuous.
10. A multidimensional version of the Kolmogorov–Chentsov theorem. Suppose $d \geq 1$, and there is a stochastic field $X : [0, 1]^d \times \Omega \rightarrow \mathbb{R}$ that satisfies $\mathbb{E}|X(s) - X(t)|^\alpha \leq C|s - t|^{d+\beta}$ for some $\alpha, \beta, C > 0$ and all $t, s \in [0, 1]^d$. Prove that there is a continuous modification of X on $[0, 1]^d$.
11. Show that the Kolmogorov–Chentsov theorem cannot be relaxed: inequality $\mathbb{E}|X_t - X_s| \leq C|t - s|$ is not sufficient for existence of a continuous modification. Hint: consider the following process: let τ be a r.v. with exponential distribution, and define $X_t = \mathbb{1}_{\{\tau \leq t\}}$.
12. Prove that there exists a Poisson process (a process with independent increments that have Poisson distribution with parameter proportional to the length of time increments) such that:
 - (a) its realizations are nondecreasing, taking only whole values a.s.
 - (b) its realizations are continuous on the right a.s.
 - (c) all the jumps of the realizations are equal to 1 a.s.
13. Give an example of a non-Gaussian 2-dimensional random vector with Gaussian marginal distributions.
14. Let $Y \sim \mathcal{N}(a, C)$ be a d -dimensional random vector. Let $Z = AY$ where A is an $n \times d$ matrix. Prove that Z is Gaussian and find its mean and covariance matrix.
15. Prove that an \mathbb{R}^d -valued random vector X is Gaussian iff for every vector $b \in \mathbb{R}^d$, the r.v. $\langle b, X \rangle$ is Gaussian.
16. Prove that $(s, t) \mapsto t \wedge s$ defined for $s, t \geq 0$ is positive semi-definite. Hint:

$$\langle \mathbb{1}_{[0,t]}, \mathbb{1}_{[0,s]} \rangle_{L^2(\mathbb{R}_+)} = t \wedge s.$$

17. Prove that $c : (s, t) \mapsto e^{-|t-s|}$ is positive semi-definite, using an auxiliary Hilbert space H and a $h_t \in H$ such that $\langle h_t, h_s \rangle = c(s, t)$.
18. Prove that if X is a Gaussian vector in \mathbb{R}^d with parameters (a, C) and C is non-degenerate, then the distribution of X is absolutely continuous w.r.t. Lebesgue measure and the density is

$$p_X(x) = \frac{1}{\det(C)^{1/2} (2\pi)^{d/2}} e^{-\frac{1}{2} \langle C^{-1}(x-a), (x-a) \rangle}.$$

19. Use the Chentsov–Kolmogorov theorem to find a condition on the mean $a(t)$ and covariance function $c(s, t)$ that guarantees existence of a continuous Gaussian process with these parameters.
20. Suppose (X_0, X_1, \dots, X_n) is a (not necessarily centered) Gaussian vector. Show that there are constants c_0, c_1, \dots, c_n such that

$$\mathbb{E}(X_0 | X_1, \dots, X_n) = c_0 + c_1 X_1 + \dots + c_n X_n.$$

Your proof should be valid irrespective of whether the covariance matrix of (X_1, \dots, X_n) is degenerate or not.

21. Consider the standard Ornstein–Uhlenbeck process X (Gaussian process with mean 0 and covariance function $c(s, t) = e^{-|t-s|}$).
 - (a) Prove that X has a continuous modification.
 - (b) Find $\mathbf{E}(X_4 | X_1, X_2, X_3)$.
22. Prove that for every centered Gaussian process X with independent increments on $\mathbb{R}_+ = [0, \infty)$ ($X(t) - X(s)$ is required to be independent of $\sigma(X(r), r \in [0, s])$ for $t \geq s \geq 0$), there is a nondecreasing nonrandom function $f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that X has the same f.d.d.'s as Y defined by $Y(t) = W(f(t))$, for a Wiener process W .
23. Let μ be a σ -finite Borel measure on \mathbb{R}^d . Prove existence of Poisson point process and (Gaussian) white noise with leading measure μ .

In both cases, the process X we are interested in is indexed by Borel sets A with $\mu(A) < \infty$, with independent values on disjoint sets, with finite additivity property for disjoint sets and such that

 - (a) $X(B)$ is Poisson with parameter $\mu(B)$ (for Poisson point process).
 - (b) $X(B)$ is centered Gaussian with variance $\mu(B)$ (for white noise).
24. Suppose the process X_t is a Gaussian process, and let H be the Hilbert space generated by $(X_t)_{t \in \mathbb{R}}$, i.e., the space consisting of L^2 -limits of linear combinations of values of X_t . Prove that every element in H is a Gaussian r.v.

2. DUE BY WEDNESDAY MAR 25

I must update the statement in the beginning of this file:

You are allowed to use books or existing electronic/online materials to solve the problems. You are allowed to work on solutions in groups, but you are required to write up solutions on your own. Besides that, you are not allowed to seek external help for specific problems. You are also not allowed to post your solutions online.

You must type/write on the first page of the assignment paper:

I have neither given nor received unauthorized help when working on this assignment.

Please give complete solutions, all claims need to be justified. Late homework will not be accepted.

Please let me know if you find any misprints or mistakes.

1. Suppose ρ is a finite measure on \mathbb{R} and $(Z(B))_{B \in \mathcal{B}(\mathbb{R})}$ is a collection of complex-valued random variables in L^2 such that $\mathbf{E}Z(B) = 0$ for all B and $\mathbf{E}Z(A)\overline{Z(B)} = \rho(A \cap B)$ for all A, B . Prove that if $A = \bigcup_{k=1}^{\infty} B_k$ for pairwise disjoint sets $(B_k)_{k \in \mathbb{N}}$, then the series $\sum_{k=1}^{\infty} Z(B_k)$ converges in L^2 to $Z(A)$.
2. Let A, η be random variables, let a r.v. ϕ be independent of (A, η) and uniform on $[0, 2\pi]$. Prove that $X_t = A \cos(\eta t + \phi)$, $t \in \mathbb{R}$, is a strictly stationary process.
3. Find the covariance function of a stationary process such that its spectral measure is $\rho(dx) = \frac{dx}{1+x^2}$.

4. Prove the law of large numbers $(X_1 + \dots + X_n)/n \xrightarrow{L^2} 0$ for centered wide-sense stationary process X satisfying $\text{cov}(X_0, X_m) \rightarrow 0$ as $m \rightarrow \infty$.
5. Let X be centered wide-sense stationary process with spectral measure ρ . Prove that $\text{cov}(X_0, X_m) \rightarrow 0$ as $m \rightarrow \infty$ is equivalent to $\rho(\{0\}) = 0$. Prove that $\text{cov}(X_0, X_m) \rightarrow 0$ implies $\rho(\{0\}) = 0$. Give an example where $\rho(\{0\}) = 0$ but $\text{cov}(X_0, X_m) \not\rightarrow 0$.
6. Give an example of a wide-sense stationary stochastic process $(X_n)_{n \in \mathbb{N}}$ such that $(X_1 + \dots + X_n)/n$ converges in L^2 to a limit that is not a constant. Explain what is going on in terms of the spectral representation.
7. Let $(X_n)_{n \in \mathbb{Z}}$ be a weakly stationary process. Prove that for any $K \in \mathbb{N}$ and any numbers $a_{-K}, a_{-K+1}, \dots, a_{K-1}, a_K$, the process $(Y_n)_{n \in \mathbb{Z}}$ defined by

$$Y_n = \sum_{k=-K}^K a_k X_{n+k}$$

is weakly stationary. Express the spectral measure of Y in terms of the spectral measure for X .

8. Describe the stationary sequence

$$X_n = \int_0^{2\pi} \cos(n\lambda) Z(d\lambda), \quad n \in \mathbb{N},$$

where $Z(\cdot)$ is the standard white noise on $[0, 2\pi)$ (i.e., a Gaussian orthogonal random measure with structural measure being Lebesgue measure.)

9. Let a stationary process $(X_n)_{n \in \mathbb{Z}}$ satisfy $E|X_0| < \infty$. Prove that with probability 1, $\lim_{n \rightarrow \infty} (X_n/n) = 0$.
10. Find the spectral measure and spectral representation for stationary process $(X_t)_{t \in \mathbb{R}}$ given by

$$X_t = A \cos(t + \phi), \quad t \in \mathbb{R},$$

where A and ϕ are independent, $A \in L^2$ and ϕ is uniform on $[0, 2\pi]$.

11. Consider a map $\theta : \Omega \rightarrow \Omega$. A set A is called (*backward*) *invariant* if $\theta^{-1}A = A$, *forward invariant* if $\theta A = A$. Prove that the collection of backward invariant sets forms a σ -algebra. Give an example of Ω and θ such that the collection of forward invariant sets does not form a σ -algebra.
12. Consider the transformation $\theta : \omega \mapsto \{\omega + \lambda\}$ on $\Omega = [0, 1)$ equipped with Lebesgue measure. Here $\{\dots\}$ denotes the fractional part of a number. This map can be interpreted as rotation of the circle (seen as $[0, 1)$ with endpoints 0 and 1 identified).

A. Prove that this dynamical system is ergodic (i.e., there are no backward invariant Borel sets besides \emptyset and Ω [UPD: or their zero-measure modifications]) if and only if $\lambda \notin \mathbb{Q}$. Hint: for $\lambda \in \mathbb{Q}$ construct a backward invariant subset, for $\lambda \notin \mathbb{Q}$ take the indicator of an invariant set and write down the Fourier series for it (w.r.t. $e^{2\pi i n x}$). What happens to this expansion under θ ?

B. Using the Birkhoff ergodic theorem and Part A, describe the limiting behavior of $(f(\omega) + f(\theta^1\omega) + \dots + f(\theta^{n-1}\omega))/n$ as $n \rightarrow \infty$, where f is any L^1 function on $[0, 1]$.

13. Let (A_k) be a sequence of i.i.d. random matrices with bounded elements. Use the subadditive ergodic theorem to prove that the Lyapunov exponent $\lambda = \lim_{n \rightarrow \infty} \frac{1}{n} \ln \|A_n A_{n-1} \dots A_1\|$ is a.s.-well-defined and nonrandom.
14. Prove that every Gaussian martingale is a process with independent increments.
15. Prove that if W_t is a standard Brownian motion, then $e^{\sigma W_t - \frac{\sigma^2 t}{2}}$ is a martingale for any $\sigma \in \mathbb{R}$.
16. Prove that if W_t is a standard Brownian motion, then $W_t^3 - 3 \int_0^t W_s ds$ is a martingale.
17. Let $X_t = t^\alpha + W_t$, where (W_t) is the standard Wiener process. Find necessary and sufficient conditions on $\alpha \in \mathbb{R}$ to ensure that $\mathbb{P}\{\lim_{t \rightarrow \infty} X_t = +\infty\} = 1$.
18. Prove (without using LIL or anything similarly advanced) that $\lim_{t \rightarrow \infty} W(t)/t = 0$ almost surely.
19. Show that the function

$$P(s, x, t, \Gamma) = P(t - s, x, \Gamma) = \int_{\Gamma} \frac{1}{\sqrt{2\pi(t-s)}} e^{-\frac{(x-y)^2}{2(t-s)}} dy$$

is a Markov transition probability function for the standard Wiener process.

20. Let $(X_t)_{t \geq 0}$ be a stochastic process. We define $\mathcal{F}_{\leq t} = \sigma(X_s, 0 \leq s \leq t)$ and $\mathcal{F}_{\geq t} = \sigma(X_s, s \geq t)$.

Prove that the following two versions of the Markov property are, in fact, equivalent:

1. For all $A \in \mathcal{F}_{\leq t}, B \in \mathcal{F}_{\geq t}, \mathbb{P}(AB|X_t) = \mathbb{P}(A|X_t)\mathbb{P}(B|X_t)$.
 2. For all $B \in \mathcal{F}_{\geq t}, \mathbb{P}(B|\mathcal{F}_{\leq t}) = \mathbb{P}(B|X_t)$.
21. Let W be a Wiener process. The previous problem shows that the process $X_t = W_{-t}, t \in (-\infty, 0]$ is Markov. Find the transition probability function for X_t .
 22. Prove that the process (X_t) given by $X_t = |W_t|$ is Markov. Find its transition probability function.
 23. Let $(N_t)_{t \geq 0}$ be the standard Poisson process. Let $X_t = (-1)^{N_t}$. Prove that (X_t) is a Markov process and find the transition probability function.

3. DUE BY FRIDAY MAY 1, 6PM

You must type/write on the first page of the assignment paper:

I have neither given nor received unauthorized help when working on this assignment.

Please give complete solutions, all claims need to be justified. Late homework will not be accepted.

Please let me know if you find any misprints or mistakes.

1. For a Wiener process W and $b > 0$, compute the density of

$$\tau_b = \inf\{t \geq 0 : W_t = b\}$$

and find $E\tau_b$.

2. We say that a r.v. τ is a stopping time w.r.t. a filtration $(\mathcal{F}_t)_{t \geq 0}$ if for every t , $\{\tau \leq t\} \in \mathcal{F}_t$.

Let $(X_t, \mathcal{F}_t)_{t \geq 0}$ be a continuous process such that $X_0 = 0$. Suppose $a > 0$ and let $\tau = \inf\{t : X(t) > a\}$ and $\nu = \inf\{t : X(t) \geq a\}$. Show that ν is a stopping time w.r.t. $(\mathcal{F}_t)_{t \geq 0}$ and τ is a stopping time w.r.t. $(\mathcal{F}_{t+})_{t \geq 0}$, where $\mathcal{F}_{t+} = \bigcap_{\varepsilon > 0} \mathcal{F}_{t+\varepsilon}$.

3. Show that if $\tau_1 \leq \tau_2 \leq \dots$ are stopping times w.r.t. to a filtration (\mathcal{F}_t) , then $\tau = \lim_{n \rightarrow \infty} \tau_n$ is also a stopping time w.r.t. to (\mathcal{F}_t) .
4. Let $\mathcal{F}_\tau = \{A : A \cap \{\tau \leq t\} \in \mathcal{F}_t\}$ for a filtration (\mathcal{F}_t) and a stopping time τ . Show that \mathcal{F}_τ is a σ -algebra.
5. Give an example of the following: a random variable $\tau \geq 0$ is not a stopping time, \mathcal{F}_τ is not a σ -algebra.
6. Suppose τ is a stopping time w.r.t. (\mathcal{F}_{t+}) . Let us define

$$\tau_n = \frac{[2^n \tau] + 1}{2^n} = \sum_{k \in \mathbb{N}} \frac{k}{2^n} \mathbb{1}_{\{\tau \in [\frac{k-1}{2^n}, \frac{k}{2^n})\}}, \quad n \in \mathbb{N}.$$

Prove that for every $n \in \mathbb{N}$, τ_n is a stopping time w.r.t. $(\mathcal{F}_t)_{t \geq 0}$, $\mathcal{F}_{\tau_n} \supset \mathcal{F}_{\tau+}$, and $\tau_n \downarrow \tau$.

7. Prove: if $(X_t, \mathcal{F}_t)_{t \geq 0}$ is a continuous process, then for any stopping time τ , X_τ is a r.v. measurable w.r.t. \mathcal{F}_τ .
8. Let (X_t, \mathcal{F}_t) be a continuous martingale and let τ be a stopping time w.r.t. \mathcal{F}_t . Prove that the “stopped” process $(X_t^\tau, \mathcal{F}_t)_{t \geq 0}$, where $X_t^\tau = X_{\tau \wedge t}$, is also a martingale.
9. Prove the following theorem (Kolmogorov, 1931) using Taylor expansions of test functions:

Suppose $(P_x)_{x \in \mathbb{R}^d}$ is a (homogeneous) Markov family on \mathbb{R}^d with transition probabilities $P(\cdot, \cdot, \cdot)$. Suppose there are continuous functions $a^{ij}(x)$, $b^i(x)$, $i, j = 1, \dots, d$, such that for every $\varepsilon > 0$, the following relations

hold uniformly in x :

$$P(t, x, B_\varepsilon^c(x)) = o(t), \quad t \rightarrow 0,$$

$$\int_{B_\varepsilon(x)} (y^i - x^i)P(t, x, dy) = b^i(x)t + o(t), \quad t \rightarrow 0,$$

$$\int_{B_\varepsilon(x)} (y^i - x^i)(y^j - x^j)P(t, x, dy) = a^{ij}(x)t + o(t), \quad t \rightarrow 0.$$

where $B_\varepsilon(x)$ is the Euclidean ball of radius ε centered at x .

Then the infinitesimal generator A of the Markov semigroup associated to the Markov family is defined on all functions f such that f itself and all its partial derivatives of first and second order are bounded and uniformly continuous. For such functions

$$Af = \frac{1}{2} \sum_{ij} a^{ij} \partial_{ij} f + \sum_i b^i \partial_i f.$$

10. Suppose $X_t = (X_t^i)_{i=1\dots d}$ satisfies

$$dX_t^i = b^i(X_t)dt + \sum_{j=1}^m \sigma_j^i(X_t)dW_t^j, \quad i = 1, \dots, d.$$

where b^i and σ_j^i are bounded Lipschitz functions and $(W^j)_{j=1}^m$ are jointly independent standard Wiener processes. Taking for granted that X is a Markov process, use the multi-dimensional Itô formula to compute the generator of the Markov semigroup on functions f such that f itself and all its partial derivatives of first and second order are bounded and uniformly continuous. Compare with the result of the previous problem.

11. Consider the Poisson transition probabilities, i.e., fix a number $\lambda > 0$ and for $i \in \mathbb{Z}$ and $t \geq 0$, let $P(i, t, \cdot)$ be the distribution of $i + \pi_{\lambda t}$, where π_s denotes a random variable with Poisson distribution with parameter $s > 0$. In other words,

$$P(i, t, \{j\}) = e^{-\lambda t} \frac{(\lambda t)^{j-i}}{(j-i)!}, \quad i \in \mathbb{Z}, \quad j \in \{i, i+1, \dots\}, \quad t > 0.$$

Find the generator of the Markov semigroup on all bounded test functions $f: \mathbb{Z} \rightarrow \mathbb{R}$.

12. Find the transition probabilities and generator associated to the Ornstein-Uhlenbeck process.
13. Let W^1 and W^2 be two independent Wiener processes w.r.t. a filtration $(\mathcal{F}_t)_{t \geq 0}$, and let X be a bounded process adapted to $(\mathcal{F}_t)_{t \geq 0}$.

For a partition t of time interval $[0, T]$ (i.e., a sequence of times $t = (t_0, t_1, \dots, t_n)$ such that $0 = t_0 < t_1 < t_2 < \dots < t_n = T$), we define

$$Q(t) = \sum_j X_{t_j} (W_{t_{j+1}}^1 - W_{t_j}^1)(W_{t_{j+1}}^2 - W_{t_j}^2).$$

Prove:

$$\lim_{\max(t_{j+1}-t_j) \rightarrow 0} Q(t) = 0 \quad \text{in } L^2.$$

14. Let (\mathcal{F}_t) be a filtration. Suppose that $0 = A_0(t) + A_1(t)W(t)$ for all t , where (A_0, \mathcal{F}_t) and (A_1, \mathcal{F}_t) are processes with C^1 trajectories, and $W(t)$ is a Wiener process w.r.t. (\mathcal{F}_t) . Prove that $A_0 \equiv 0$ and $A_1 \equiv 0$.
15. Suppose $f \in L^2([0, T], \mathcal{B}, \text{Leb})$. For all $h > 0$, we define

$$f_h(t) = \begin{cases} 0, & 0 \leq t < h; \\ \frac{1}{h} \int_{(k-1)h}^{kh} f(s) ds, & kh \leq t < (k+1)h \wedge T. \end{cases}$$

Prove that $f_h \xrightarrow{L^2} f$ and $\|f_h\|_{L^2} \leq \|f\|$. Hint: do not work with generic L^2 functions right away; consider a smaller supply of functions f that behave nicely under this kind of averaging.

16. Compute $\int_0^T W^2(t) dW(t)$ directly as the limit of

$$\sum W^2(t_k)(W(t_{k+1}) - W(t_k))$$

17. The so-called Stratonovich stochastic integral may be defined for a broad class of adapted processes X_t via

$$\int_0^T X_t \circ dW_t \stackrel{\text{def}}{=} \lim_{\max(t_{j+1}-t_j) \rightarrow 0} \sum_j \frac{X_{t_{j+1}} + X_{t_j}}{2} (W_{t_{j+1}} - W_{t_j}) \quad \text{in } L^2.$$

Impose any conditions you need on X and express the difference between the Itô and Stratonovich integrals in terms of quadratic covariation between X and W . Compute $\int_0^T W_t \circ dW_t$. Is the answer a martingale?

18. Let $\mathcal{M}_c^2 = \{\text{square-integrable martingales with continuous paths}\}$. Prove that if $(M_t, \mathcal{F}_t) \in \mathcal{M}_c^2$, then

$$\mathbb{E}[(M_t - M_s)^2 | \mathcal{F}_s] = \mathbb{E}[M_t^2 - M_s^2 | \mathcal{F}_s] = \mathbb{E}[\langle M \rangle_t - \langle M \rangle_s | \mathcal{F}_s], \quad s < t.$$

19. Suppose $(M_t, \mathcal{F}_t) \in \mathcal{M}_c^2$, X is a simple process, and $(X \cdot M)_t = \int_0^t X_s dM_s$. Prove that

$$\mathbb{E}[(X \cdot M)_t - (X \cdot M)_s)^2 | \mathcal{F}_s] = \mathbb{E} \left[\int_s^t X_r^2 d\langle M \rangle_r | \mathcal{F}_s \right], \quad s < t.$$

20. Let $M \in \mathcal{M}_c^2$. Prove that

$$Y \cdot (X \cdot M) = (YX) \cdot M$$

for simple processes X, Y . Find reasonable weaker conditions on X and Y guaranteeing the correctness of this identity in the sense of square integrable martingales.

21. Suppose $(M_t, \mathcal{F}_t) \in \mathcal{M}_c^2$, and X, Y are bounded processes. Prove that

$$\langle X \cdot M, Y \cdot M \rangle_t = \int_0^t X_s Y_s d\langle M \rangle_s.$$

Here, for two processes $M, N \in \mathcal{M}_c^2$ the cross-variation $\langle M, N \rangle_t$ is defined by

$$\langle M, N \rangle_t = \frac{\langle M + N \rangle_t - \langle M - N \rangle_t}{4}.$$

22. Let us define the process X by

$$X_t = e^{\lambda t} X_0 + \varepsilon e^{\lambda t} \int_0^t e^{-\lambda s} dW_s, \quad t \geq 0.$$

Here $\lambda \in \mathbb{R}$, $\varepsilon > 0$, W is a standard Wiener process, and X_0 is a square-integrable r.v., independent of W . Prove that

$$dX_t = \lambda X_t dt + \varepsilon dW_t.$$

23. Prove that if $f : [0, \infty)$ is a deterministic function, bounded on any interval $[0, t]$, then

$$X_t = \int_0^t f(s) dW_s, \quad t \geq 0,$$

is a Gaussian process. Find its mean and covariance function.

24. In the context of Problem 22, find all the values of λ with the following property: there are a and σ^2 such that if $X_0 \sim \mathcal{N}(a, \sigma^2)$, then (X_t) is a stationary process.
25. [Feynman–Kac formula.] Suppose $u_0 : \mathbb{R} \rightarrow [0, \infty)$ and $\phi : [0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$ are smooth bounded functions. Suppose that $u : [0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$ is a smooth function solving the Cauchy problem for the following equation:

$$\begin{aligned} \partial_t u(t, x) &= \frac{1}{2} \partial_{xx} u(t, x) + \phi(t, x) u(t, x), \\ u(0, x) &= u_0(x). \end{aligned}$$

Using the Itô formula and martingales, prove that

$$u(t, x) = \mathbf{E} e^{\int_0^t \phi(t-s, x+W_s) ds} u_0(x + W_t), \quad t > 0, \quad x \in \mathbb{R},$$

where W is a standard Wiener process. (If you need further assumptions on u and/or its derivatives, use PDE theory or just impose any assumptions you need.)