

First-Passage Kinetic Monte Carlo Algorithm for Complex Reaction-Diffusion Systems

Aleksandar Donev¹

Lawrence Postdoctoral Fellow

Lawrence Livermore National Laboratory

In collaboration with:

Vasily V. Bulatov, *LLNL*

CMMD Seminar, July 2009

¹This work performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344 (LLNL-PRES-408395).

- 1 Introduction
- 2 First-Passage Kinetic Monte Carlo
- 3 Results: Radiation Damage in *Fe*

Reaction-Diffusion Particle Models

- Systems of diffusing particles that react with other particles upon collision are a common model in computational materials science: **reaction-diffusion models**.
- Examples include: *diffusion-limited chemical reactions, signal transduction in cells, radiation damage in metals, dopant implantation in semiconductors, epitaxial deposition and growth of thin films, population dynamics, etc.*
- Continuum models are often unable to correctly capture some key property, notably the **strong heterogeneity** in space/time (e.g., *clustering*), and intrinsic **fluctuations** (e.g., *nucleation*)
- **Continuous-Time Markov Chain** models are an attractive but *expensive* alternative:
A collection of Brownian hard spheres that diffuse through a homogeneous continuum and react upon collision with other particles or surfaces.

Example: Chemotaxis in *E. Coli*.




Figure: Bacterial chemotaxis as studied using *Smoldyn* by Karen Lipkow and Steven Andrews [*J. Bacteriol.* 187(1):45-53, (2005)]

Example: Radiation Damage





Figure: Defect creation and clustering in metals during irradiation.

Diffusion Kinetic Monte Carlo




(MNG)

- Some or all unit events are diffusion hops: a set of N hard objects walking randomly on a lattice or in continuum space.
- Upon collision particles *react* (**collision events**).
- Example: Diffusion-controlled annihilation $A + A \rightarrow 0$.
- Great many diffusion hops necessary to bring particles to collisions at low density.

Traditional *synchronous* n -fold event-driven algorithm (BKL). Other types of Poisson events (birth, decay, boundary, etc.) are easy to handle.


Green's Function Diffusion Theory



- Given a region of space Ω , one can determine the probability distributions for when and where (on $\partial\Omega$) a particle will first leave that region (*first-event prediction*).
- Given that a particle has not yet left that region, one can determine the probability of finding the particle at some point inside the region at a given time (*no-event propagation*).

For pairs of particles, reduce to two *independent* **center-of-mass** and **difference walkers**.

First Passage Kinetic Monte Carlo (FPKMC)



(MNG)

- Construct disjoint protective regions (cubes, spheres) at $t = 0$.
- Main events are **(super)hops** to $\partial\Omega$. For each walker (particle or pair) **randomly** draw first passage time from the appropriate PDF.
- Find the earliest time in the queue, propagate the particle/pair to boundary/collision, construct a new protective region, insert back into queue with a new event time, repeat [1, 2]!

Advantages of the Algorithm



- The FPKMC algorithm is **exact** for continuous diffusion problems because it breaks the hard N -body problem into tractable one- and two-body problems.
- It is the first use we know of of **time-dependent Green's functions**.
- The algorithm automatically adjusts to variable timescales: **multiscale**.
- We have a code that implements different types of reactions (annihilation, coalescence, chemical reactions, decay/emission, hard-sphere repulsion).

Disadvantages of the Algorithm

- The method is significantly more **complicated** to implement than BKL KMC and it requires analytical solutions (1-body and 2-body problems).
- Multi-particle reactions cause complications or slowdown (ex., nearly triple collisions).
- One can combine the asynchronous super-hops with local synchronous small hops in a **mixed time-driven/event-driven approach** [3].
- FPKMC can be viewed as a **general-purpose accelerator** that brings particle within interaction range quickly, after which application-specific handling should take over.

FPKMC for Radiation Damage

- Diffusion-reaction model for **radiation damage in metals**: diffusing and reacting *vacancies* and *interstitials* and their clusters



- A Kinetic Monte Carlo (KMC) simulation faithfully follows every **atomistic event**: *cascade insertion*, *diffusion hop*, *annihilation*, *recombination*, *clustering*, *dissociation*, *trapping*, *escape*, etc [4].

Radiation Damage KMC Model

- Very simple additive hard-sphere model for **testing** purposes, based on work by Barbu *et al.*

- **Species:**

- *monomers*, including highly-mobile interstitials (I) and less-mobile vacancies (V), with diffusion coefficient

$$D_1 = D_0 e^{-E_m/kT}$$

- mobile *cluster species*, including dimers (I_2 and V_2) and trimers (I_3 and V_3), with radius

$$R_c \sim R_0 + (R_1 - R_0)c^{1/3}$$

- *immobile* species representing clusters larger than any of the mobile species (I_c and V_c)
- *Frenkel pairs* (IV), inserted randomly with some rate

Model contd.

• Reactions:

- *Coalescence*: $I + I \rightarrow I_2$ or $V + V_3 \rightarrow V_{c=4}$
- *Partial annihilation*: $I_2 + V_{c=4} \rightarrow V_2$
- *Decay or emission*: $V_{c=5} \rightarrow V_{c=4} + V$, or $I_2 \rightarrow I + I$, with rate

$$\Gamma_c = \Gamma_0 D_1 a^{-2} c^{2/3} e^{-E_b(c)/kT},$$

$$E_b(c) = E_f + [E_b(2) - E_f] \frac{c^{2/3} - (c-1)^{2/3}}{2^{2/3} - 1}.$$

Validation




Figure: Comparison of the density profile between FPKMC (symbols) and CEA OKMC code from LAKIMOCA [5] (lines) simulations of a $0.287\mu\text{m}$ -thick film of α -iron subjected to 120 seconds of electron radiation at a temperature $T = 200^\circ\text{C}$.

Computational Extrapolation

- We can develop FPKMC models that match *accelerator* (e.g., JANNUS) experiments (**high dose rate**) then use the same models to simulate material behavior over a *nuclear reactor* lifetime (**low dose rate**): **computational extrapolation**
- *Approximate* scaling of results from the accelerated testing experiments is achieved when the temperature is raised sufficiently so as to keep the ratio of vacancy diffusion coefficient to irradiation rate constant.
- FPKMC allows us to quantify the mismatch with **atomistic fidelity** up to *realistic doses* (10dpa)!
- But the model is **too simple** to capture realistic physics! Work is underway at CEA to develop more detailed and accurate models.

Lab vs. Reactor

Swelling ($\Delta V / V$)





Figure: Swelling due to 10dpa electron irradiation in a pure Fe film

Future Directions

- **Apply** FPKMC to a wider range of problems, notably, **systems biology** (compare to *Smoldyn* library).
- Extend first-passage to *lattice models* (discrete space), notably, defect diffusion through **dense alloys**.
- Approximately handle *continuous interaction potentials* (ala Brownian Dynamics), and most importantly, **long-ranged interactions** (e.g., electrostatic or elastic).
- When very fast species (e.g., interstitials) are present they slow down even event-driven algorithms: multiscale methods to utilize this **separation of time-scales**.
- **Parallelization** of event-driven algorithms remains an important challenge.
- The particle model can be coupled to a more efficient *continuum* model in a **hybrid method** (see next talk).

References/Questions?



T. Oppelstrup, V. V. Bulatov, G. H. Gilmer, M. H. Kalos, and B. Sadigh.
First-Passage Monte Carlo Algorithm: Diffusion without All the Hops.
Phys. Rev. Lett., 97(23):230602, 2006.



T. Oppelstrup, V. V. Bulatov, A. Donev, M. H. Kalos, G. H. Gilmer, and B. Sadigh.
First-Passage Kinetic Monte Carlo Method.
Preprint, [arXiv:0905.3575](https://arxiv.org/abs/0905.3575), 2009.



A. Donev.
Asynchronous event-driven particle algorithms.
SIMULATION: Transactions of The Society for Modeling and Simulation International, 85(4):229–242, 2008.



A. Donev, V. V. Bulatov, T. Oppelstrup, G. H. Gilmer, B. Sadigh, and M. H. Kalos.
A First-Passage Kinetic Monte Carlo Algorithm for Complex Diffusion-Reaction Systems.
Preprint, [arXiv:0905.3576](https://arxiv.org/abs/0905.3576), 2009.



C. Domain, C. S. Becquart, and L. Malerba.
Simulation of radiation damage in Fe alloys: an object kinetic Monte Carlo approach.
Journal of Nuclear Materials, 335(1):121–145, 2004.