

Random Matrix Theory, homework 2, due October 5.

Problem 1. The Circular Unitary Ensemble is a log-correlated random field. Let $(e^{i\theta_k})_{1 \leq k \leq N}$ be the eigenvalues of a Haar-distributed matrix in $U(N)$. The eigenangles have joint probability distribution

$$\mathbb{P}(d\theta) = \frac{1}{N!} \prod_{1 \leq i < j \leq N} |e^{i\theta_i} - e^{i\theta_j}|^2 \frac{d\theta_1}{2\pi} \dots \frac{d\theta_N}{2\pi}.$$

- (i) Prove that $\chi = \sum_{k=1}^N \delta_{\theta_k}$ is a determinantal point process with correlation kernel

$$K(x, y) = K^{(N)}(x, y) = \frac{1}{2\pi} \frac{\sin N \frac{x-y}{2}}{\sin \frac{x-y}{2}}$$

with respect to the Lebesgue measure on $(0, 2\pi)$.

- (ii) Let $\phi : [0, 2\pi) \rightarrow \mathbb{R}$ be bounded measurable. Prove that

$$\mathbb{E} \prod_{k=1}^N (1 + \phi(\theta_k)) = \sum_{n \geq 0} \frac{1}{n!} \int_{(0, 2\pi)^n} \prod_{j=1}^n \phi(x_j) \det_{n \times n} K(x_i, x_j) dx_1 \dots dx_n.$$

You will need to explain why the right hand side converges.

- (iii) Read Section 3 in the book *Trace ideals and applications*.
 (iv) Let $A \subset [0, 2\pi)$ be measurable. On $L^2(A)$, define $K\phi$ the convolution operator with kernel $K\phi$, where ϕ is bounded measurable:

$$(K\phi)(f)(x) = \int K(x, y)\phi(y)f(y)dy.$$

Prove that $K\mathbb{1}_A$ is trace-class with spectrum in $[0, 1]$. Let $X = \chi(A)$. Show that

$$\log \mathbb{E}(e^{i\xi X}) = \log \det(\text{Id} + K\mathbb{1}_A(e^{i\xi} - 1)) = - \sum_{k=1}^{\infty} \frac{(1 - e^{i\xi})^k}{k} \text{Tr}((K\mathbb{1}_A)^k).$$

- (v) The formula $\log \mathbb{E}(e^{i\xi X}) = \sum_{\ell=1}^{\infty} C_{\ell}(X) \frac{(i\xi)^{\ell}}{\ell!}$ defines the cumulants $C_{\ell}(X)$ of the random variable X . Prove that for any $\ell \geq 3$,

$$C_{\ell}(X) = (-1)^{\ell}(\ell - 1)! \text{Tr}(K\mathbb{1}_A - (K\mathbb{1}_A)^{\ell}) + \sum_{j=2}^{\ell-1} \alpha_{j\ell} C_j(X)$$

for some universal constants $\alpha_{j\ell}$.

- (vi) Take $A = [0, x)$ ($x \in (0, 2\pi)$) in this question and the next one. Prove that

$$C_2(X) = \int_0^x du \int_x^{2\pi} dv |K(u, v)|^2 \underset{N \rightarrow \infty}{\sim} \pi^{-2} \log N.$$

- (vii) Prove that $C_{\ell}(X/\sqrt{\log N})$ converges to 0 as $N \rightarrow \infty$ for any $\ell \geq 3$. For this you can first prove the trace inequality

$$0 \leq \text{Tr}(K\mathbb{1}_A - (K\mathbb{1}_A)^{\ell}) \leq (\ell - 1) \text{Tr}(K\mathbb{1}_A - (K\mathbb{1}_A)^2).$$

Show that $(X - \mathbb{E}X)/\sqrt{\log N}$ converges weakly to a Gaussian random variable with variance π^{-2} . Compare this result to the case of N independent uniform points on the circle.

- (viii) Consider $X_k = \chi([0, x_k)) - Nx_k/(2\pi)$ where $x_k = N^{-\alpha_k}$, $0 < \alpha_1 < \dots < \alpha_{\ell} < 1$. Prove a joint central limit theorem for the random variables X_1, \dots, X_{ℓ} as $N \rightarrow \infty$. Compare this result to the case of N independent uniform points on the circle.

Exercise 1. Fluctuations for the Ginibre ensemble. Consider the joint distribution of eigenvalues from the Ginibre ensemble,

$$\mathbb{P}(dz) = \frac{1}{Z_N} \prod_{1 \leq i < j \leq N} |z_i - z_j|^2 \prod_{i=1}^N e^{-N|z_i|^2} dA(z_i)$$

where dA is the Lebesgue measure on \mathbb{C} . Let \mathcal{C} be a smooth Jordan curve, with interior A , finite length $\ell(\mathcal{C})$, strictly included in the unit disk $\{|z| < 1\}$. Let $X_{\mathcal{C}} = \chi(A) - \mathbb{E}(\chi(A))$ where $\chi = \sum_{i=1}^N \delta_{z_i}$. By mimicking the method from Problem 1, prove the weak convergence

$$\frac{X_{\mathcal{C}}}{\ell(\mathcal{C})^{1/2} N^{1/4}} \rightarrow \mathcal{N}(0, c)$$

as $N \rightarrow \infty$, with some c independent of \mathcal{C} . What about joint convergence of $(X_{\mathcal{C}_1}, \dots, X_{\mathcal{C}_n})$ where all Jordan curves $\mathcal{C}_1, \dots, \mathcal{C}_n$ satisfy the above assumptions?

Exercise 2. The semicircle law for band matrices. Let H_N be a symmetric matrix with $H_N(i, j)$ a standard Bernoulli random variable when $|i - j| \leq W/2$ or $||i - j| - N| \leq W/2$, 0 otherwise. All entries are independent, up to the symmetry constraint. Assume $1 \ll W \leq N$.

Prove that the empirical spectral measure of $W^{-1/2}H_N$ converges (in probability, say) to the semicircle distribution $\varrho(s) = (2\pi)^{-1} \sqrt{(4 - s^2)_+}$.

Open problem 1. In Exercise 1, what happens when the Jordan curve is not smooth and has infinite length? In particular, if $\log \text{var}(X_{\mathcal{C}}) \sim \alpha(\mathcal{C}) \log N$, does $\alpha(\mathcal{C})$ only depend on the Hausdorff dimension of \mathcal{C} ? Or the Minkowski dimension?

Open problem 2. In Exercise 2, let u_1, \dots, u_N be the L^2 -normalized eigenvectors of H_N and $\alpha \in (0, 1)$, $D > 0$.

Assume $\alpha < 1/2$. Prove that there exists $\delta > 0$ such that for N greater than some $N_0(\alpha, D)$, with probability at least $1 - N^{-D}$ the following holds: for any $k \in \llbracket 1, N \rrbracket$, $\|u_k\|_{\infty} > N^{-1/2+\delta}$.

Assume $\alpha > 1/2$. Prove that for any $\delta > 0$, for N greater than some $N_0(\alpha, D, \delta)$, with probability at least $1 - N^{-D}$ the following holds: for any $k \in \llbracket 1, N \rrbracket$, $\|u_k\|_{\infty} < N^{-1/2+\delta}$.