

Generalized flows, intrinsic stochasticity, and turbulent transport

Weinan E and Eric Vanden Eijnden

Courant Institute of Mathematical Sciences, New York University, New York, NY 10012

Communicated by Grigory Isaakovich Barenblatt, University of California, Berkeley, CA, April 13, 2000 (received for review February 11, 2000)

The study of passive scalar transport in a turbulent velocity field leads naturally to the notion of generalized flows, which are families of probability distributions on the space of solutions to the associated ordinary differential equations which no longer satisfy the uniqueness theorem for ordinary differential equations. Two most natural regularizations of this problem, namely the regularization via adding small molecular diffusion and the regularization via smoothing out the velocity field, are considered. White-in-time random velocity fields are used as an example to examine the variety of phenomena that take place when the velocity field is not spatially regular. Three different regimes, characterized by their degrees of compressibility, are isolated in the parameter space. In the regime of intermediate compressibility, the two different regularizations give rise to two different scaling behaviors for the structure functions of the passive scalar. Physically, this means that the scaling depends on Prandtl number. In the other two regimes, the two different regularizations give rise to the same generalized flows even though the sense of convergence can be very different. The “one force, one solution” principle is established for the scalar field in the weakly compressible regime, and for the difference of the scalar in the strongly compressible regime, which is the regime of inverse cascade. Existence and uniqueness of an invariant measure are also proved in these regimes when the transport equation is suitably forced. Finally incomplete self similarity in the sense of Barenblatt and Chorin is established.

Recent efforts to understand the fundamental physics of hydrodynamic turbulence have concentrated on the explanation of observed violations of Kolmogorov’s scaling. These violations reflect the occurrence of large fluctuations in the velocity field on the small scales, a phenomenon referred to as intermittency. Some progress in the understanding of intermittency has been achieved recently through the study of simple model problems that include Burgers equation (1, 2) and the passive advection of a scalar by a velocity field of known statistics (3–6). This paper is a summary of the many interesting mathematical issues that arise in the problem of passive scalar advection together with our understanding of these issues. We put some of our results in the perspective of a new phenomenological model proposed recently by Barenblatt and Chorin (7, 8) using the formalism of incomplete self similarity.

Generalized Flows

Consider the transport equation for the scalar field $\theta^\kappa(x, t)$ in \mathbb{R}^d :

$$\frac{\partial \theta^\kappa}{\partial t} + (u(x, t) \cdot \nabla) \theta^\kappa = \kappa \Delta \theta^\kappa. \quad [1]$$

We will be interested in θ^κ in the limit as $\kappa \rightarrow 0$. It is known from classical results that if u is Lipschitz continuous in x , then as $\kappa \rightarrow 0$, θ^κ converges to θ , the solution of

$$\frac{\partial \theta}{\partial t} + (u(x, t) \cdot \nabla) \theta = 0. \quad [2]$$

Furthermore, if we define $\{\varphi_{s,t}(x)\}$ as the flow generated by the velocity field u , satisfying the ordinary differential equations (ODEs)

$$\frac{d\varphi_{s,t}(x)}{dt} = u(\varphi_{s,t}(x), t), \quad \varphi_{s,s}(x) = x \quad [3]$$

for $s < t$, then the solution of the transport equation in 2 for the initial condition $\theta^\kappa(x, 0) = \theta_0(x)$ is given by

$$\theta(x, t) = \theta_0(\varphi_{0,t}^{-1}(x)) = \theta_0(\varphi_{t,0}(x)). \quad [4]$$

This classical scenario breaks down when u fails to be Lipschitz continuous in x , which is precisely the case for fully developed turbulent velocity fields. In this case, Kolmogorov’s theory of turbulent flows suggests that u is Hölder continuous only with an exponent roughly equal to $\frac{1}{3}$ for $d = 3$. In such situations, the solution of the ODEs in 3 may fail to be unique (so long as u is continuous, Peano’s theorem tells us that solutions to 3 do exist), and we then have to consider probability distributions on the set of solutions in order to solve the transport equation in 2. This is the essence of the notion of generalized flows proposed by Brenier (9, 10) (see also refs. 11 and 12).

There are two ways to think about probability distributions on the solutions of the ODEs in 3. We can think of them either as probability measures on the path-space (functions of t) supported by paths that are solutions of 3, or we can think of them as transition probability at time t if the starting position at time s is x . In the classical situation, when u is Lipschitz continuous, this transition probability degenerates to a point mass centered at the unique solution of 3. When Lipschitz condition fails, this transition probability may be nondegenerate and the system in 3 is intrinsically stochastic.

There is a parallel story for the case when u is a white-in-time random process defined on a probability space $(\Omega, \mathcal{F}, \mathcal{P})$. We will denote the elements in Ω by ω and indicate the dependence on realization of the random velocity field by a super- or subscript ω . In connection with the transport equation in 2, it is most natural to consider the stochastic ODEs,

$$d\varphi_{s,t}^\omega(x) = u(\varphi_{s,t}^\omega(x), t) dt, \quad \varphi_{s,s}^\omega(x) = x, \quad [5]$$

in the Stratonovich sense. In this case, it is shown (13) that if the local characteristic of u is spatially twice continuously differentiable, then the system in 5 has a unique solution. Such conditions are not satisfied by typical turbulent velocity fields on the scale of interest. When the regularity condition on u fails, there are at least two natural ways to regularize 3 or 5. The first is to add diffusion:

$$d\varphi_{s,t}^{\omega,\kappa}(x) = u(\varphi_{s,t}^{\omega,\kappa}(x), t) dt + \sqrt{2\kappa} d\beta(t) \quad [6]$$

and consider the limit as $\kappa \rightarrow 0$. We will call this the κ -limit. The second is to smooth out the velocity field. Let ψ_ε be defined

Abbreviation: ODE, ordinary differential equation.

*To whom reprint requests should be addressed at: 207 Fine Hall, PACM, Washington Road, Princeton, NJ 08544. E-mail: weinan@princeton.edu.

as $\psi_\varepsilon(x) = \varepsilon^{-d}\psi(x/\varepsilon)$, where ψ is a standard mollifier: $\psi \geq 0$, $\int_{\mathbb{R}^d} \psi dx = 1$, ψ decays fast at infinity. Let $u^\varepsilon = u \star \psi_\varepsilon$ and consider

$$d\varphi_{s,t}^{\omega,\varepsilon}(x) = u^\varepsilon(\varphi_{s,t}^{\omega,\varepsilon}(x), t) dt \quad [7]$$

in the limit as $\varepsilon \rightarrow 0$. We will call this the ε -limit. Physically, κ plays the role of molecular diffusivity, and ε can be thought of as a crude model of the viscous cutoff scale. The κ -limit corresponds to the situation when the Prandtl number, defined here as the ratio of ε and κ , tends to zero, $Pr \rightarrow 0$, whereas the ε -limit corresponds to the situation when the Prandtl number diverges, $Pr \rightarrow \infty$. The following questions naturally arise:

- (i) How do the flows and the passive scalar behave statistically in the κ - and ε -limits?
- (ii) Does there exist a unique statistical steady state when the transport equation in **1** is suitably forced?
- (iii) What are the statistical and geometrical properties of solutions in the statistical steady state?

Below we address these questions by using a specific model introduced by Kraichnan (14).

Before proceeding further, we relate the regularized flows in **6** and **7** to the solutions of the transport equations. Consider the κ -regularization first. It is convenient to introduce the backward transition probability

$$g_\omega^\kappa(x, t | dy, s) = \mathbf{E}_\beta \delta(y - \varphi_{t,s}^{\omega,\kappa}(x)) dy, \quad s < t, \quad [8]$$

where the expectation is taken with respect to $\beta(t)$, and $\varphi_{t,s}^{\omega,\kappa}(x)$ is the flow inverse to $\varphi_{s,t}^{\omega,\kappa}(x)$ defined in **6** (i.e., $\varphi_{s,t}^{\omega,\kappa}(x)$ is the forward flow, and $\varphi_{t,s}^{\omega,\kappa}(x)$ is the backward flow). The action of g_ω^κ generates a semigroup of transformation

$$S_{t,s}^{\omega,\kappa} \psi(x) = \int_{\mathbb{R}^d} \psi(y) g_\omega^\kappa(x, t | dy, s), \quad [9]$$

for all test functions ψ . $\theta_\omega^\kappa(x, t) = S_{t,s}^{\omega,\kappa} \psi(x)$ solve the transport equation in **1** for the initial condition $\theta_\omega^\kappa(x, s) = \psi(x)$. Similarly, for the flow in **7**, define

$$S_{t,s}^{\omega,\varepsilon} \psi(x) = \psi(\varphi_{t,s}^{\omega,\varepsilon}(x)), \quad s < t. \quad [10]$$

$\theta_\omega^\varepsilon(x, t) = S_{t,s}^{\omega,\varepsilon} \psi(x)$ solves the transport equation

$$\frac{\partial \theta^\varepsilon}{\partial t} + (u^\varepsilon(x, t) \cdot \nabla) \theta^\varepsilon = 0, \quad [11]$$

with initial condition $\theta(x, s) = \psi(x)$. Similar definitions can be given for forward flows but we will restrict attention to the backward ones because we are interested primarily in scalar transport. The results given below generalize trivially to forward flows.

Kraichnan Model

In ref. 14, Kraichnan introduced one of the simplest model of passive scalar by considering the advection by a Gaussian spatially nonsmooth and white-in-time velocity field. The fact that white-in-time velocity fields may exhibit intermittency was first recognized by Majda (15, 16). Definitive work on the Kraichnan model was done afterwards in refs. 3–6.

We will consider a generalization of the Kraichnan model introduced in ref. 17 (see also ref. 18). The velocity u is assumed to be a statistically homogeneous, isotropic and stationary Gaussian field with mean zero and covariance

$$\mathbf{E} u_\alpha(x, t) u_\beta(y, s) = (C_0 \delta_{\alpha\beta} - c_{\alpha\beta}(x - y)) \delta(t - s). \quad [12]$$

We assume that u has a correlation length ℓ_0 , i.e. the covariance in **12** decays fast for $|x - y| > \ell_0$. Consequently, $c_{\alpha\beta}(x) \rightarrow C_0 \delta_{\alpha\beta}$

as $|x|/\ell_0 \rightarrow \infty$. On the other hand, we will be interested mainly in small-scale phenomena, for which $|x| \ll \ell_0$. In this range, we take $c_{\alpha\beta}(x) = d_{\alpha\beta}(x) + O(|x|^2/\ell_0^2)$ with

$$d_{\alpha\beta}(x) = A d_{\alpha\beta}^P(x) + B d_{\alpha\beta}^S(x), \quad [13]$$

and

$$d_{\alpha\beta}^P(x) = D \left(\delta_{\alpha\beta} + \xi \frac{x_\alpha x_\beta}{|x|^2} \right) |x|^\xi, \quad [14]$$

$$d_{\alpha\beta}^S(x) = D \left((d + \xi - 1) \delta_{\alpha\beta} - \xi \frac{x_\alpha x_\beta}{|x|^2} \right) |x|^\xi.$$

D is a parameter with dimension $[\text{length}]^{2-\xi} [\text{time}]^{-1}$. The dimensionless parameters A and B measure the divergence and rotation of the field u . $A = 0$ corresponds to incompressible fields with $\nabla \cdot u = 0$. $B = 0$ corresponds to irrotational fields with $\nabla \times u = 0$. The parameter ξ measures the spatial regularity of u . For $\xi \in (0, 2)$, the local characteristic of u fails to be twice differentiable, and this fact has important consequences for both the transport equation in **2** and the systems of ODEs in **3** or **5**.

Existing physics literature concentrates on the κ -limit for the Kraichnan model. Let $\mathcal{S}^2 = A + (d - 1)B$, $\mathcal{C}^2 = A$, $\mathcal{P} = \mathcal{C}^2/\mathcal{S}^2$. $\mathcal{P} \in [0, 1]$ is a measure of the degree of compressibility of u . The pioneering work of Gawędzki and Vergassola (17) (see also ref. 18) identifies two different regimes for the κ -limit:

- (i) The strongly compressible regime when $\mathcal{P} \geq d/\xi^2$. In this regime, g_ω^κ converges to a flow of maps, i.e., there exists a two-parameter family of maps $\{\varphi_{t,s}^\omega(x)\}$ such that

$$g_\omega^\kappa(x, t | dy, s) \rightarrow \delta(y - \varphi_{t,s}^\omega(x)) dy. \quad [15]$$

Moreover, particles have finite probability to coalesce under the flow of $\{\varphi_{t,s}^\omega(x)\}$. In other words the flow is not invertible.

- (ii) When $\mathcal{P} < d/\xi^2$, g_ω^κ converges to a “generalized stochastic flow”

$$g_\omega^\kappa(x, t | dy, s) \rightarrow g_\omega(x, t | dy, s), \quad [16]$$

and the limit g_ω is a nontrivial probability distribution in y . This means that the image of a particle under the flow defined by the velocity field u is nonunique and has a nontrivial distribution. In other words, particle trajectories branch.

The same classification of the flows was obtained by Le Jan and Raimond (18) using Wiener chaos expansion with no explicit reference to the κ -limit. In contrast, our primary motivation is to study the limit of physical regularizations.

The following result answers question *i* and also points out that there are three different regimes if both the κ - and the ε -limits are considered.

THEOREM 1. *In the strongly compressible regime when $\mathcal{P} \geq d/\xi^2$, there exists a two-parameter family of random maps $\{\varphi_{t,s}^\omega(x)\}$, such that for all smooth test functions ψ and for all (s, t, x) , $s < t$,*

$$\mathbf{E} (S_{t,s}^{\omega,\kappa} \psi(x) - \psi(\varphi_{t,s}^\omega(x)))^2 \rightarrow 0, \quad [17]$$

as $\kappa \rightarrow 0$, and

$$\mathbf{E} (\psi(\varphi_{t,s}^{\omega,\varepsilon}(x)) - \psi(\varphi_{t,s}^\omega(x)))^2 \rightarrow 0 \quad [18]$$

as $\varepsilon \rightarrow 0$. Moreover, the limiting flow $\{\varphi_{t,s}^\omega(x)\}$ coalesces in the sense that for almost all (t, x, y) , $x \neq y$, we can define a time τ such that $-\infty < \tau < t$ a.s. and

$$\varphi_{t,s}^\omega(x) = \varphi_{t,s}^\omega(y) \quad \text{for } s \leq \tau. \quad [19]$$

In the weakly compressible regime when $\mathcal{P} \leq (d + \xi - 2)/2\xi$, there exists a random family of generalized flows $g_\omega(x, t | dy, s)$, such that for all test function ψ ,

$$S_{t,s}^\omega \psi(x) = \int_{\mathbb{R}^d} \psi(y) g_\omega(x, t | dy, s) \quad [20]$$

satisfies

$$\mathbf{E} (S_{t,s}^{\omega,\kappa} \psi(x) - S_{t,s}^\omega \psi(x))^2 \rightarrow 0, \quad [21]$$

as $\kappa \rightarrow 0$ for all (s, t, x) , $s < t$, and

$$\mathbf{E} \left(\int_{\mathbb{R}^d} \eta(x) (\psi(\varphi_{t,s}^{\omega,\varepsilon}(x)) - S_{t,s}^\omega \psi(x)) dx \right)^2 \rightarrow 0, \quad [22]$$

as $\varepsilon \rightarrow 0$ for all (s, t) , $s < t$, and for all test functions η . Moreover, $g_\omega(x, t | dy, s)$ is nondegenerate in the sense that

$$S_{t,s}^\omega \psi^2(x) - (S_{t,s}^\omega \psi(x))^2 > 0 \quad a.s. \quad [23]$$

In the intermediate regime when $(d + \xi - 2)/2\xi < \mathcal{P} < d/\xi^2$, there exists a random family of generalized flows $g_\omega(x, t | dy, s)$, such that for all test functions ψ and for all (s, t, x) , $s < t$,

$$\mathbf{E} (S_{t,s}^{\omega,\kappa} \psi(x) - S_{t,s}^\omega \psi(x))^2 \rightarrow 0 \quad [24]$$

as $\kappa \rightarrow 0$. In the ε -limit, the flows $\varphi_{t,s}^{\omega,\varepsilon}(x)$ converge in the sense of distributions, i.e., there exists a family of probability densities $\{G_n(x_1, \dots, x_n, t | y_1, \dots, y_n, s) dy_1 \cdots dy_n\}$, $n = 1, 2, \dots$, such that

$$\begin{aligned} \mathbf{E} \psi(\varphi_{t,s}^{\omega,\varepsilon}(x_1), \dots, \varphi_{t,s}^{\omega,\varepsilon}(x_n)) &\rightarrow \int_{\mathbb{R}^d \times \dots \times \mathbb{R}^d} \psi(y_1, \dots, y_n) \\ &\times G_n(x_1, \dots, x_n, t | y_1, \dots, y_n, s) dy_1 \cdots dy_n, \end{aligned} \quad [25]$$

as $\varepsilon \rightarrow 0$ for any continuous function ψ with compact support. Furthermore, the ε -limit coalesces in the sense that

$$\begin{aligned} G_2(x_1, x_2, t | y_1, y_2, s) &= \tilde{G}_2(x_1, x_2, t | y_1, y_2, s) \\ &+ A(y_1, x_1, x_2, t, s) \delta(y_1 - y_2), \end{aligned} \quad [26]$$

with $A > 0$ when $t > s$. Here \tilde{G}_2 is the absolutely continuous part of G_2 with respect to the Lebesgue measure. Similar statements hold for the other G_n . In particular, the G_n differ from the moments of the κ -limit g_ω , defined in 24.

Rephrasing the content of this result, we have strong convergence to a family of flow maps in the strongly compressible regime for both the κ -limit and ε -limit. In the weakly compressible regime, we have strong convergence to a family of generalized flows for the κ -limit, but weak convergence to the same limit for the ε -regularization. In fact, using the terminology of Young measures (19), the limiting generalized flow $\{g_\omega(x, t | dy, s)\}$ is nothing but the Young measure for the sequence of flow maps $\{\varphi_{s,t}^{\omega,\varepsilon}(x)\}$. Finally, in contrast to what is observed in the other two regimes, the ε -limit and κ -limit are not the same in the intermediate regime. As we will see below, the structure functions of the passive scalar field scale differently in the two limits.

From Theorem 1, it is natural to define the solution of the transport equation in 2 for the initial condition $\theta_\omega(x, s) = \theta_0(x)$ as

$$\theta_\omega(x, t) = S_{t,s}^\omega \theta_0(x) = \int_{\mathbb{R}^d} \theta_0(y) g_\omega(x, t | dy, s), \quad [27]$$

for the weakly compressible and the intermediate regimes in the κ -limit (nondegenerate cases), and as

$$\theta_\omega(x, t) = \theta_0(\varphi_{t,s}^\omega(x)) \quad [28]$$

for the strongly compressible regime. In the intermediate regime in the ε -limit, it makes sense to look at the limiting moments of $\theta_\omega^\varepsilon(x, t)$, because we have as $\varepsilon \rightarrow 0$

$$\begin{aligned} \mathbf{E}(\theta_\omega^\varepsilon(x_1, t) \cdots \theta_\omega^\varepsilon(x_n, t)) &\rightarrow \int_{\mathbb{R}^d \times \dots \times \mathbb{R}^d} \theta_0(y_1) \cdots \theta_0(y_n) \\ &\times G_n(x_1, \dots, x_n, t | y_1, \dots, y_n, s) dy_1 \cdots dy_n. \end{aligned} \quad [29]$$

It should be noted that when g_ω is nondegenerate, there exists an anomalous dissipation mechanism for the scalar, whereas no such anomalous dissipation is present in the coalescence cases (17). The presence of anomalous dissipation is the primary reason why the transport equation in 2 has a statistical steady state (invariant measure) if it is appropriately forced, as we will show later.

Details of the proof of Theorem 1 are given (W.E and E.V.E., unpublished results). Crucial to the proof is the study of $P(\rho | r, s)$, defined through ε -regularization as

$$\begin{aligned} &\int_0^\infty \eta(r) P(\rho | r, s - t) dr \\ &= \lim_{\varepsilon \rightarrow 0} \mathbf{E} \eta(|\varphi_{t,s}^{\omega,\varepsilon}(y) - \varphi_{t,s}^{\omega,\varepsilon}(z)|), \end{aligned} \quad [30]$$

where η is a test function, and similarly through κ -regularization. Here $\rho = |y - z|$ and $s < t$. $P(\rho | r, s)$ can be thought of as the probability density that two particles have distance r at time $s < t$ if their final distance is ρ at time t . For the Kraichnan model, P satisfies the backward equation

$$-\frac{\partial P}{\partial s} = -\frac{\partial}{\partial r} (b(r)P) + \frac{\partial^2}{\partial r^2} (a(r)P) \quad [31]$$

for the final condition $\lim_{s \rightarrow 0^-} P(\rho | r, s) = \delta(r - \rho)$ and with $a(r)$, $b(r)$ such that

$$\begin{aligned} a(r) &= D(S^2 + \xi C^2) r^\xi + O(r^2/\ell_0^2), \\ b(r) &= D((d - 1 + \xi)S^2 - \xi C^2) r^{\xi-1} + O(r/\ell_0^2). \end{aligned} \quad [32]$$

For $r \gg \ell_0$, $a(r)$ tends to C_0 , $b(r)$ to $C_0(d - 1)/r$, and the equation in 31 reduces to a diffusion equation with constant coefficient. The equation in 31 is singular at $r = 0$. The proof of Theorem 1 is essentially reduced to the study of this singular diffusion equation. This is also the main step for which the white-in-time nature of the velocity field is crucial.

Structure Functions

We now study some consequences of Theorem 1 for the passive scalar θ_ω defined in 27 or 28. We note that the scaling of the second-order structure function is the same for the κ - and the ε -limits in the strongly and the weakly compressible cases[†], but it differs in the intermediate regime as a result of the difference between the limits in 24 and 25. For simplicity of presentation, we assume that θ_0 is isotropic and Gaussian. Denote $(n \in \mathbb{N})$

$$S_{2n}(|x - y|, t) = \mathbf{E}(\theta_\omega(x, t) - \theta_\omega(y, t))^{2n}, \quad [33]$$

$$\text{or } S_{2n}(|x - y|, t) = \lim_{\varepsilon \rightarrow 0} \mathbf{E}(\theta_\omega^\varepsilon(x, t) - \theta_\omega^\varepsilon(y, t))^{2n}, \quad [33']$$

in the intermediate regime in the ε -limit. In the strongly compressible case, we have for both the κ - and the ε -limits

$$S_2(r, t) = O(r^\xi), \quad [34]$$

[†]For the weakly compressible case, because the ε -limit is a weak limit one has to consider the structure functions of the mollified temperature field $\psi_\varepsilon * \theta^\varepsilon$ in order to obtain the limiting scaling as $\varepsilon \rightarrow 0$.

with

$$\zeta = \frac{2 - d - \xi + 2\xi\mathcal{P}}{1 + \xi\mathcal{P}}. \quad [35]$$

In the weakly compressible case, we have for both the κ - and the ε -limits

$$S_2(r, t) = O(r^{2-\xi}). \quad [36]$$

In the intermediate regime, the limits differ, and the κ -limit scales as in 36, whereas the ε -limit scales as in 34. The equations in 34 and 36 can be derived on expressing S_2 in terms of P ; the details are given (W.E and E.V.E., unpublished work).

It is interesting to discuss the higher order structure functions both in the nondegenerate and coalescence cases in 34 and 36, because their scalings highlight very different behaviors of the scalar. We consider first the coalescence cases, which are simpler. In these cases, because of the absence of dissipative anomaly, all higher order structure functions can again be expressed in terms of P , and it can be shown (17) that

$$S_{2n}(r, t) = O(r^\zeta), \quad [37]$$

with ζ given by 35 for all $n \geq 1$. In fact, coalescence implies that the temperature field θ_ω tends to become flat except possibly on a zero-measure set, where it presents shock-like discontinuities. Such a situation with two kinds of spatial structures for θ_ω is usually referred to as bifractal, and, in simple cases, one may identify ζ with the codimension of the set supporting the discontinuities of θ_ω (20–22).

Nondegenerate cases are more complicated. In these cases, one expects that θ_ω presents a spatial behavior much richer than in the coalescence cases, with all kinds of scalings present. This is the multi-fractal situation for which the higher order structure functions behave as

$$S_{2n}(r, t) = O(r^{\zeta_{2n}}), \quad [38]$$

with $\zeta_{2n} < n(2 - \xi)$ for $n > 1$. The actual value of the ζ_n 's cannot be obtained by dimensional analysis, and one has to resort to various sophisticated perturbation techniques (see refs. 3–6). We will consider again the scaling of the structure functions at statistical steady state in *Incomplete Self Similarity*.

One Force, One Solution Principle for Temperature

We now turn to question *iii* and consider the existence of a statistical steady state for the transport equation with appropriate forcing. We restrict attention to the nondegenerate cases, which include the weakly compressible regime and the intermediate regime in the κ -limit. Indeed, in these regimes the nondegeneracy of $g_\omega(x, t | dy, s)$ as a probability distribution in y implies dissipation of energy or, phrased differently, decay in memory in the semigroup $S_{t,s}$ generated by $\{g_\omega\}$. We show that the anomalous dissipation is strong enough that the forced transport equation has a unique invariant measure for both the weakly compressible regime and the intermediate regime in the κ -limit. This result, however, depends on the finiteness of ℓ_0 . In limit as $\ell_0 \rightarrow \infty$ an invariant measure exists only for the weakly compressible regime.

We will consider (compare with 2)

$$\frac{\partial \theta}{\partial t} + (u(x, t) \cdot \nabla) \theta = b(x, t), \quad [39]$$

where b is a white-noise forcing such that

$$\mathbf{E} b(x, t) b(y, s) = B(|x - y|) \delta(t - s). \quad [40]$$

$B(r)$ is assumed to be smooth and rapidly decaying to zero for $r \gg L$; L will be referred to as the forcing scale. The solution of 39 for the initial condition $\theta_\omega(x, s) = \theta_0(x)$ is understood as

$$\theta_\omega(x, t) = S_{t,s}^\omega \theta_0(x) + \int_s^t S_{t,\tau}^\omega b(x, \tau) d\tau. \quad [41]$$

Define the product probability space $(\Omega_u \times \Omega_b, \mathcal{F}_u \times \mathcal{F}_b, \mathcal{P}_u \times \mathcal{P}_b)$, and the shift operator $T_\tau \omega(t) = \omega(t + \tau)$, with $\omega = (\omega_u, \omega_b)$. We have

THEOREM 2 (One force, one solution I). *For $d > 2$, in the weakly compressible regime and in the intermediate regime in the κ -limit for almost all ω , there exists a unique solution of 39 defined on $\mathbb{R}^d \times (-\infty, \infty)$. This solution can be expressed as*

$$\theta_\omega^*(x, t) = \int_{-\infty}^t S_{t,s}^\omega b(x, s) ds. \quad [42]$$

Furthermore, the map $\omega \rightarrow \theta_\omega^*$ satisfies the invariance property

$$\theta_{T_\tau \omega}^*(x, t) = \theta_\omega^*(x, t + \tau). \quad [43]$$

Theorem 2 is the “one force, one solution” principle articulated in ref. 23. Because of the invariance property 43, the map in 42 leads to a natural invariant measure. As a consequence we have

COROLLARY 3. *For $d > 2$, in the weakly compressible regime and in the intermediate regime in the κ -limit, there exists a unique invariant measure on $L_{loc}^2(\mathbb{R}^d \times \Omega)$ for the dynamics defined by 39.*

The connection between map 42 and the invariant measure, together with uniqueness, is explained in ref. 23. The restriction on the dimensionality in *Theorem 2* arises because the velocity field has a finite correlation length ℓ_0 : *Theorem 2* is changed into *Theorem 4* below in the limit as $\ell_0 \rightarrow \infty$, which can be considered after appropriate redefinition of the velocity field.

We sketch the proof of *Theorem 2*. Basically, it amounts to verifying that the dissipation in the system is strong enough in the sense that

$$\mathbf{E} \left(\int_{T_1}^{T_2} S_{t,t+s}^\omega b(x, s) ds \right)^2 \rightarrow 0, \quad [44]$$

as $T_1, T_2 \rightarrow -\infty$ for fixed x and t . The average in 44 is given explicitly by

$$\int_{T_1}^{T_2} \int_0^\infty B(r) P(0 | r, s) dr ds, \quad [45]$$

where P satisfies 31. The convergence of the integral in 44 depends on the rate of decay in $|s|$ of $P(0 | r, s)$. The latter can be estimated by studying the equation in 31 (?), which yields $P(0 | r, s) \sim Cr^\alpha |s|^{-d/2}$ with $\alpha = (d - 1 - \xi(\xi + 1)P)/(1 + \xi P)$ for $|s|$ large and $r \ll \ell_0$. Hence, the integral in s in 45 tends to zero as $T_1, T_2 \rightarrow -\infty$ if $d > 2$. It follows that the invariant measure in 42 exists provided $d > 2$.

We now ask what happens if we let $\ell_0 \rightarrow \infty$ in order to emphasize the effect of the inertial range of the velocity? This question, however, has to be considered carefully because the velocity field with the covariance in 12 diverges as $\ell_0 \rightarrow \infty$. The right way to proceed is to consider an alternative velocity v , taken to be Gaussian and white-in-time but *nonhomogeneous*, with covariance

$$\begin{aligned} \mathbf{E} v_\alpha(x, t) v_\beta(y, s) \\ = (c_{\alpha\beta}(x - a) + c_{\alpha\beta}(a - y) - c_{\alpha\beta}(x - y)) \delta(t - s). \end{aligned} \quad [46]$$

For finite ℓ_0 , one has $v(x, t) = u(x, t) - u(a, t)$, where a is arbitrary but fixed. However, v makes sense in the limit as $\ell_0 \rightarrow \infty$.

Denote by $\vartheta_\omega(x, t)$ the temperature field advected by v , i.e., the solution of transport equation **39** with u replaced by v :

$$\frac{\partial \vartheta}{\partial t} + (v(x, t) \cdot \nabla) \vartheta = b(x, t). \quad [47]$$

Restricting to zero initial condition, it follows from the homogeneity of the forcing that the single-time moments of θ_ω and ϑ_ω coincide for finite ℓ_0 , but in contrast to θ_ω , ϑ_ω makes sense as $\ell_0 \rightarrow \infty$. Thus, ϑ_ω is a natural process to study the limit as $\ell_0 \rightarrow \infty$, and we have

THEOREM 4 (One force, one solution II). *In the limit as $\ell_0 \rightarrow \infty$ in the weakly compressible regime, for almost all ω , there exists a unique solution of **47** defined on $\mathbb{R}^d \times (-\infty, \infty)$. This solution can be expressed as*

$$\vartheta_\omega^*(x, t) = \int_{-\infty}^t S_{t,s}^\omega b(x, s) ds, \quad [48]$$

where $S_{t,s}^\omega$ is the semigroup for the generalized flow associated with the velocity defined in **46** in the limit as $\ell_0 \rightarrow \infty$. Furthermore the map $\omega \rightarrow \vartheta_\omega^*$ satisfies the invariance property

$$\vartheta_{T,\omega}^*(x, t) = \vartheta_\omega^*(x, t + \tau). \quad [49]$$

As a direct result we also have

COROLLARY 5. *In the limit as $\ell_0 \rightarrow \infty$, in the weakly compressible regime there exists a unique invariant measure on $L_{loc}^2(\mathbb{R}^d \times \Omega)$ for the dynamics defined by **47**.*

Notice that, as $\ell_0 \rightarrow \infty$, the anomalous dissipation is not strong enough in the intermediate regime in the κ -limit, for which no statistical steady state with finite energy exists.

The proof of *Theorem 4* proceeds as the one for *Theorem 2*, but the estimate for P in **45** changes as $P(0|r, s) \sim Cr^\alpha |s|^{-(\alpha+1)/(2-\varepsilon)}$ with $\alpha = (d-1 - \xi(\xi+1)P)/(1 + \xi P)$ for $|s|$ large and $\rho \ll \ell_0$. It follows that the integral in s in **45** converges as $T_1, T_2 \rightarrow -\infty$ in the weakly compressible regime only.

One Force, One Solution Principle for the Temperature Difference

Because no anomalous dissipation is present in the coalescence cases, i.e., the strongly compressible regime and the intermediate regime in the ε -limit, no invariant measure for the temperature field exists in these regimes. It makes sense, however, to ask about the existence of an invariant measure for the temperature difference, i.e., to consider

$$\delta\theta_\omega(x, y, t) = \int_T^t S_{t,s}^\omega (b(x, s) - b(y, s)) ds, \quad [50]$$

in the limit as $T \rightarrow -\infty$. When θ_ω^* exists, one has

$$\delta\theta_\omega^*(x, y, t) = \lim_{T \rightarrow -\infty} \delta\theta_\omega(x, y, t) = \theta_\omega^*(x, t) - \theta_\omega^*(y, t), \quad [51]$$

but it is conceivable that $\delta\theta_\omega^*$ exists in the coalescence cases even though θ_ω^* is not defined. The reason is that coalescence of the generalized flow implies that the temperature field flattens with time, which is a dissipation mechanism as far as the temperature difference is concerned. Of course, this effect has to overcome the fluctuations produced by the forcing, and the existence of an invariant measure such as **50** will depend on how fast particles coalesce under the flow, which happens only in the limit as $\ell_0 \rightarrow \infty$ (i.e., for the alternate velocity defined in **46**), as we show now.

For finite ℓ_0 , if we were to consider two particles separated by much more than the correlation length ℓ_0 , the dynamics of their distance under the flow is governed by the equation in **31** for $r \gg \ell_0$, i.e., by a diffusion equation with constant diffusion coefficient on the scale of interest. It follows that no tendency of coalescence is observed before the distance becomes smaller

than ℓ_0 , which, as shown below, does not happen fast enough to overcome the fluctuations produced by the forcing. In other words,

LEMMA 6. *In the coalescence cases, for finite ℓ_0 , there is no invariant measure with finite energy for the temperature difference.*

Consider now the limit as $\ell_0 \rightarrow \infty$, and let $\delta\vartheta_\omega(x, y, t) = \vartheta_\omega(x, t) - \vartheta_\omega(y, t)$, where ϑ_ω solves the equation in **47**. The temperature difference $\delta\vartheta_\omega$ satisfies the transport equation

$$\begin{aligned} \frac{\partial \delta\vartheta}{\partial t} + (v(x, t) \cdot \nabla_x + v(y, t) \cdot \nabla_y) \delta\vartheta \\ = b(x, t) - b(y, t). \end{aligned} \quad [52]$$

We have

THEOREM 7 (One force, one solution III). *In the limit as $\ell_0 \rightarrow \infty$, for almost all ω , in the strongly and the weakly compressible regimes, as well as in the intermediate regime if the flow is nondegenerate, there exists a unique solution of **52** defined on $\mathbb{R}^d \times (-\infty, \infty)$. This solution can be expressed as*

$$\delta\vartheta_\omega^*(x, y, t) = \int_{-\infty}^t S_{t,s}^\omega (b(x, s) - b(y, s)) ds, \quad [53]$$

where $S_{t,s}^\omega$ is the semigroup for the generalized flow associated with the velocity defined in **46** in the limit as $\ell_0 \rightarrow \infty$. Furthermore, the map $\omega \rightarrow \delta\vartheta_\omega^*$ satisfies the invariance property

$$\delta\vartheta_{T,\omega}^*(x, y, t) = \delta\vartheta_\omega^*(x, y, t + \tau). \quad [54]$$

An immediate consequence of this theorem is

COROLLARY 8. *In the limit as $\ell_0 \rightarrow \infty$, in the strongly and the weakly compressible regimes, as well as in the intermediate regime if the flow is nondegenerate, there exists a unique invariant measure on $L_{loc}^2(\mathbb{R}^d \times \Omega)$ for the dynamics defined by **52**.*

The proof of *Theorem 7* proceeds similarly to the proof of *Theorem 2*. In the nondegenerate cases, one studies the convergence of (compare **44**)

$$\mathbb{E} \left(\int_{T_1}^{T_2} S_{t,t+s}^\omega (b(x, s) - b(y, s)) ds \right)^2 \rightarrow 0 \quad [55]$$

as $T_1, T_2 \rightarrow -\infty$ for fixed x and t . The average in **55** can be expressed in terms of P , and it can be shown (unpublished results) that the expression in **55** converges as $T_1, T_2 \rightarrow -\infty$ in the nondegenerate cases. In the strongly compressible regimes, because of the existence of a flow of maps, **55** is replaced by

$$\mathbb{E} \left(\int_{T_1}^{T_2} (b(\varphi_{t,s}^\omega(x), s) - b(\varphi_{t,s}^\omega(y), s)) ds \right)^2. \quad [56]$$

This average can again be expressed in terms of P , and it can be shown that the convergence of the time integral in **56** depends on the rate at which P loses mass at $r = 0+$ (i.e., the rate at which particles coalesce). The analysis of the equation in **31** shows that the process is fast enough that the integral over s in **56** tends to zero as $T_1, T_2 \rightarrow -\infty$ in the strongly compressible regime. In contrast, the equivalent of **56** in the intermediate regime in the ε -limit can be shown to diverge as $T_1, T_2 \rightarrow -\infty$.

It can be shown that the invariant measure has finite correlation functions of all orders, even though these results do not by themselves imply uniqueness of stationary solutions to the n -point Fokker–Planck equation. The task of studying the passive scalar is now changed to the study of the short distance behavior of these correlation functions.

Incomplete Self Similarity

We finally consider the scaling of the structure functions based on the invariant measure $\delta\vartheta^*$ defined in 53. Denote

$$\mathcal{S}_n(|x - y|) = \mathbf{E}|\delta\vartheta_\omega^*(x, y, t)|^n. \quad [57]$$

The dimensional parameters are $B_0 = B(0)$ ([temperature]²[time]⁻¹), D ([length]^{2- ξ} [time]⁻¹), L ([length]). It follows that

$$\mathcal{S}_n(r) = \left(\frac{B_0 r^{2-\xi}}{D}\right)^{n/2} f_n\left(\frac{r}{L}\right), \quad [58]$$

where the f_n are dimensionless functions that cannot be obtained by dimensional arguments. For instance, the scalings in 37 and 38 correspond to different f_n . It is, however, obvious from the equation 58 that, provided the limit exists and is nonzero

$$\lim_{L \rightarrow \infty} \mathcal{S}_n(r) = C_n \left(\frac{B_0 r^{2-\xi}}{D}\right)^{n/2} O(r^{n(2-\xi)/2}), \quad [59]$$

where $C_n = \lim_{r \rightarrow \infty} f_n(r/L)$ are numerical constants. The scaling in 59 is usually referred to as the normal scaling because, consistent with Kolmogorov's picture, it is independent of the forcing or dissipation scales. In contrast, anomalous scaling is a statement that the structure functions diverge in the limit of infinite forcing scale, $L \rightarrow \infty$. In the spirit of Barenblatt and Chorin (7, 8), we may say that normal scaling holds in case of complete self similarity, whereas anomalous scaling is equivalent to incomplete self similarity.

It is interesting to discuss the existence or nonexistence of the limit in 59 for both the coalescence and nondegenerate cases. When the flow coalesces, because of the existence of a flow of maps and the absence of dissipative anomaly, the \mathcal{S}_{2n} of even order $2n \geq 2$ can be computed exactly (17). It gives $\mathcal{S}_{2n}(r) = \infty$ for $n \geq \zeta/(2 - \xi)$, whereas

$$\mathcal{S}_{2n}(r) = O(r^\zeta) \quad \text{for } n < \frac{\zeta}{2 - \xi}, \quad [60]$$

where ζ is given in 35. Thus, for $n < \zeta/(2 - \xi)$,

$$f_{2n}(r) = O\left(\left(\frac{r}{L}\right)^{\zeta - n(2-\xi)}\right). \quad [61]$$

1. E. W., Khanin, K., Mazel, A., Sinai, Ya. G. *Phys. Rev. Lett.* **78**, 1904–1907.
2. E. W. & Vanden Eijnden, E. (1999) *Phys. Rev. Lett.* **83**, 2572–2575.
3. Shraiman, B. & Siggia, E. (1994) *Phys. Rev. E* **49**, 2912–2927.
4. Chertkov, M., Falkovich, G., Kolokolov, I. & Lebedev, V. (1995) *Phys. Rev. E* **51**, 5609–5627.
5. Gawędzki, K. & Kupiainen, A. (1995) *Phys. Rev. Lett.* **75**, 3834–3837.
6. Balkovsky, E. & Lebedev, V. (1998) *Phys. Rev. E* **58**, 5776–5795.
7. Barenblatt, G. I. & Chorin, A. J. (1997) *Comm. Pure Appl. Math.* **50**, 381–398.
8. Barenblatt, G. I., & Chorin, A. J. (1998) *SIAM Rev.* **40**, 265–291.
9. Brenier, Y. (1989) *J. Am. Math. Soc.* **2**, 225–255.
10. Brenier, Y. (1993) *Arch. Rational Mech. Anal.* **122**, 323–351.
11. DiPerna, R. J. & Majda, A. J. (1986) *Comm. Math. Phys.* **108**, 667–689.
12. Shnirelman, A. I. (1994) *Geom. Funct. Anal.* **4**, 586–620.

It follows that f_{2n} and hence \mathcal{S}_{2n} , tend to zero as $L \rightarrow \infty$ for $2 \leq n < \zeta/(2 - \xi)$, whereas they are infinite for all L for $n \geq \zeta/(2 - \xi)$. In fact, in the coalescence case, it can be shown (17) that on scales much larger than the forcing scale L , the structure functions of order $n < \zeta/(2 - \xi)$ behave as

$$\mathcal{S}_{2n}(r) \sim C_{2n} r^{n(2-\xi)} \quad \text{as } r/L \rightarrow \infty. \quad [62]$$

Thus in the coalescence case, it is more natural to consider the limit as $L \rightarrow 0$ of the structure functions, for which the expression in 62 shows the absence of intermittency corrections.

In the non-degenerate case, one has

$$\mathcal{S}_2(r) = O(r^{2-\xi}), \quad [63]$$

whereas perturbation analysis gives for the higher order structure functions (3–6)

$$\mathcal{S}_{2n}(r) = O(r^{\zeta_{2n}}), \quad [64]$$

with $\zeta_{2n} < n(2 - \xi)$ for $2n > 2$. It follows that $f_{2n}(r) = O(1)$, whereas

$$f_{2n}(r) = O\left(\left(\frac{r}{L}\right)^{\zeta_{2n} - n(2-\xi)}\right), \quad 2n > 2. \quad [65]$$

In other words, as $L \rightarrow \infty$, \mathcal{S}_2 has a limit that exhibits normal scaling, whereas the \mathcal{S}_{2n} , $2n > 2$, diverge. This may be closely related to the argument in refs. 7 and 8 that, in appropriate limits, intermittency corrections may disappear, and higher than fourth order structure functions may not exist. We note, however, that Barenblatt and Chorin were discussing the case of infinite Reynolds number (here infinite Peclet number, $\kappa \rightarrow 0$) at finite L , whereas we require $L \rightarrow \infty$.

We thank many people for helpful discussions, including G. Barenblatt, E. Balkovsky, M. Chertkov, A. Chorin, U. Frisch, J. Goodman, A. Majda, and S. R. S. Varadhan. We are particularly grateful to K. Gawędzki for pointing out an error in the first draft of this paper. WE is partially supported by a Presidential Faculty Fellowship from National Science Foundation (NSF). E.V.E. is partially supported by the NSF Grant DMS-9510356.

13. Kunita, H. (1990) *Stochastic Flows and Stochastic Differential Equations* (Cambridge Univ. Press, Cambridge).
14. Kraichnan, R. H. (1968) *Phys. Fluids* **11**, 945–963.
15. Majda, A. J. (1993) *J. Stat. Phys.* **73**, 515–542.
16. Majda, A. J. & Kramer, P. R. (1999) *Phys. Rep.* **314**, 237–574.
17. Gawędzki, K. & Vergassola, M. (2000) *Physica D* **138**, 63–90.
18. Le Jan, Y. & Raimond, O. (1999) math.PR/9909147.
19. Tartar, L. (1983) *Systems of Nonlinear Partial Differential Equations* Vol. III, pp. 263–285, NATO Math. Phys. Sci, Reidel, Dordrecht, the Netherlands).
20. Parisi, G. & Frisch, U. (1985) *Turbulence and Predictability in Geophysical Fluid Dynamics* (North-Holland, Amsterdam), pp. 83–87.
21. Jaffard, S. (1997) *SIAM J. Math. Anal.* **28**, 944–970.
22. Jaffard, S. (1997) *SIAM J. Math. Anal.* **28**, 971–978.
23. E. W., Khanin, K., Mazel, A. & Sinai, Ya. G. (2000), in press.