

Change of Variables

Liming Pang

Another commonly used technique for integration is Change of Variable, also called Integration by Substitution.

Recall the Chain Rule for differentiation: If $y = f(u)$ and $u = g(x)$, then

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = \frac{df}{du}(g(x)) \frac{dg}{dx}(x)$$

The above implies that

$$dy = df(g(x)) = f'(g(x))g'(x) dx = f'(g(x)) dg(x) = f'(u) du$$

so we obtain the following theorem:

Theorem 1. (*Change of Variables*) If $f(u)$ is continuous and $u = g(x)$ is continuously differentiable, then

$$\int f(g(x))g'(x) dx = \int f(u) du$$

Remark 2. There are two possible situations to apply the method of change of variable. The first case is when we can write the given integral in the form of $\int f(g(x))g'(x) dx$ by a good choice of $g(x)$, and then apply the formula. See Example 4,5,6

The second case is to use the theorem in the reversed way: let x be the intermediate variable by letting $x = g(t)$, so $\int f(x) dx = \int f(g(t))g'(t) dt$. See Example 7

Remark 3. We can also write the change of variable in differential notation:

$$\int f(g(x)) dg(x) = \int f(u) du$$

where $u = g(x)$

Example 4. Compute $\int x^2 + 2x + 1 dx$

$$\begin{aligned}\int x^2 + 2x + 1 dx &= \int (x + 1)^2 dx \\ &= \int (x + 1)^2 d(x + 1) \\ &= \frac{1}{3}(x + 1)^3\end{aligned}$$

Example 5. Compute $\int x\sqrt{1 + x^2} dx$

$$\begin{aligned}\int x\sqrt{1 + x^2} dx &= \int \frac{1}{2}\sqrt{1 + x^2} d(1 + x^2) \\ &= \frac{1}{2} \int \sqrt{1 + x^2} d(1 + x^2) \\ &= \frac{1}{2} \times \frac{2}{3}(1 + x^2)^{\frac{3}{2}} + C \\ &= \frac{1}{3}(1 + x^2)^{\frac{3}{2}} + C\end{aligned}$$

Example 6. Compute $\int 2xe^{-x^2} dx$

$$\begin{aligned}\int 2xe^{-x^2} dx &= - \int e^{-x^2} d(-x^2) \\ &= -e^{-x^2} + C\end{aligned}$$

Example 7. Compute $\int \ln x dx$

Let $x = e^t$, so $t = \ln x$.

$$\begin{aligned}\int \ln x dx &= \int \ln e^t de^t = \int tde^t \\ &= te^t - \int e^t dt \\ &= te^t - e^t + C \\ &= x \ln x - x + C\end{aligned}$$

There is also a corresponding formula for definite integrals:

Theorem 8. *If $f(u)$ is continuous and $u = g(x)$ is continuously differentiable, then*

$$\int_a^b f(g(x))g'(x) dx = \int_{g(a)}^{g(b)} f(u) du$$

Proof. Let F be an antiderivative of f , then

$$\int_a^b f(g(x))g'(x) dx = F(g(x)) \Big|_a^b = F(u) \Big|_{g(a)}^{g(b)} = \int_{g(a)}^{g(b)} f(u) du$$

□

Example 9. *Evaluate $\int_1^e \frac{1+\ln x}{x} dx$*

There are different alternatives to apply the rules:

Method I:

$$\begin{aligned} \int_1^e \frac{1 + \ln x}{x} dx &= \int_1^e (1 + \ln x) d \ln x \\ &= \int_{\ln 1}^{\ln e} (1 + u) du \\ &= \int_0^1 (1 + u) du \\ &= u + \frac{1}{2}u^2 \Big|_0^1 \\ &= \frac{3}{2} \end{aligned}$$

Method II:

$$\begin{aligned} \int_1^e \frac{1 + \ln x}{x} dx &= \int_1^e (1 + \ln x) d(1 + \ln x) \\ &= \int_{1+\ln 1}^{1+\ln e} u du \\ &= \int_1^2 u du \\ &= \frac{1}{2}u^2 \Big|_1^2 \\ &= \frac{3}{2} \end{aligned}$$

Method III: Let $x = e^t$

$$\begin{aligned}\int_1^e \frac{1 + \ln x}{x} dx &= \int_0^1 \frac{1 + \ln e^t}{e^t} de^t \\ &= \int_0^1 \frac{1+t}{e^t} (e^t)' dt \\ &= \int_0^1 \frac{1+t}{e^t} e^t dt \\ &= \int_0^1 1+t dt \\ &= \frac{3}{2}\end{aligned}$$

Example 10. Evaluate $\int_0^1 \sqrt{1+x} dx$

$$\begin{aligned}\int_0^1 \sqrt{1+x} dx &= \int_0^1 \sqrt{1+x} d(1+x) \\ &= \int_1^2 \sqrt{u} du \\ &= \frac{4}{3}\sqrt{2} - \frac{2}{3}\end{aligned}$$

Example 11. Compute $\int_0^1 x^3 \sqrt{1+x^2} dx$

Method I:

$$\begin{aligned}\int_0^1 x^3 \sqrt{1+x^2} dx &= \frac{1}{2} \int_0^1 x^2 \sqrt{1+x^2} d(1+x^2) \\ &= \frac{1}{2} \int_0^1 ((1+x^2) - 1) \sqrt{1+x^2} d(1+x^2) \\ &= \frac{1}{2} \int_{1+0^2}^{1+1^2} (u-1) u^{\frac{1}{2}} du \\ &= \frac{1}{2} \int_1^2 u^{\frac{3}{2}} - u^{\frac{1}{2}} du \\ &= \frac{1}{2} \left(\frac{2}{5} u^{\frac{5}{2}} - \frac{2}{3} u^{\frac{3}{2}} \right) \Big|_1^2 = \frac{2}{15} + \frac{2}{15} \sqrt{2}\end{aligned}$$

There is another way of writing the above computation, though the reason behind are the same:

Let $u = 1 + x^2$, then $du = d(1 + x^2) = (1 + x^2)'du = 2xdx$

$$\begin{aligned}
 \int_0^1 x^3 \sqrt{1+x^2} dx &= \int_{1+0^2}^{1+1^2} x^3 \sqrt{u} \frac{1}{2x} du \\
 &= \int_1^2 \frac{x^2}{2} \sqrt{u} du \\
 &= \int_1^2 \frac{u-1}{2} \sqrt{u} du \\
 &= \frac{1}{2} \int_1^2 u^{\frac{3}{2}} - u^{\frac{1}{2}} du \\
 &= \frac{1}{2} \left(\frac{2}{5} u^{\frac{5}{2}} - \frac{2}{3} u^{\frac{3}{2}} \right) \Big|_1^2 \\
 &= \frac{2}{15} + \frac{2}{15} \sqrt{2}
 \end{aligned}$$

Method II: Let $x = \sqrt{t^2 - 1} = (t^2 - 1)^{\frac{1}{2}}$

$$\begin{aligned}
 \int_0^1 x^3 \sqrt{1+x^2} dx &= \int_1^{\sqrt{2}} ((t^2 - 1)^{\frac{1}{2}})^3 (1 + ((t^2 - 1)^{\frac{1}{2}})^2)^{\frac{1}{2}} d(t^2 - 1)^{\frac{1}{2}} \\
 &= \int_1^{\sqrt{2}} (t^2 - 1)^{\frac{3}{2}} (1 + t^2 - 1) ((t^2 - 1)^{\frac{1}{2}})' dt \\
 &= \int_1^{\sqrt{2}} (t^2 - 1)^{\frac{3}{2}} t \left(\frac{1}{2} (t^2 - 1)^{-\frac{1}{2}} 2t \right) dt \\
 &= \int_1^{\sqrt{2}} t^2 (t^2 - 1) dt \\
 &= \int_1^{\sqrt{2}} t^4 - t^2 dt \\
 &= \frac{t^5}{5} - \frac{t^3}{3} \Big|_1^{\sqrt{2}} \\
 &= \frac{2}{15} + \frac{2}{15} \sqrt{2}
 \end{aligned}$$