How many ways can one draw a graph?

János Pach[*]
Courant Institute, NYU
and Hungarian Academy of Sciences
pach@cims.nyu.edu

Géza Tóth[†]
Rényi Institute
Hungarian Academy of Sciences
geza@renyi.hu

Dedicated to Miklós Simonovits on his sixtieth birthday

Abstract

Using results from extremal graph theory, we determine the asymptotic number of *string graphs* with n vertices, i.e., graphs that can be obtained as the intersection graph of a system of continuous arcs in the plane. The number becomes much smaller, for any fixed d, if we restrict our attention to systems of arcs, any two of which cross at most d times. As an application, we estimate the number of different drawings of the complete graph K_n with n vertices under various side conditions.

1 Introduction

Given a simple graph G, is it possible to represent its vertices by simply connected regions in the plane so that two regions overlap if and only if the corresponding two vertices are adjacent? In other words, is G isomorphic to the *intersection graph* of a set of simply connected regions in the plane? This deceptively simple extension of propositional logic and its generalizations are often referred to in the literature as *topological inference problems* [CGP98a], [CGP98b],[CHK99]. They have proved to be relevant in the area of geographic information systems [E93], [EF91] and in graph drawing [DETT99]. In spite of many efforts [K91a], [K98] (and false claims [SP92], [ES93]), until very recently no algorithm was known for their solution. Two years ago, we showed [PT02] that the problem is *decidable.* Shortly after a more elegant proof was found by Schaefer and Stefankovič [SS01a], who went on to prove that the question is in NP [SS01b].

Since each element of a finite system of regions in the plane can be replaced by a simple continuous arc ("string") lying in its interior so that the intersection pattern of these arcs is the same as that of the original regions, it is enough to restrict our attention to *string graphs,* i.e., to intersection graphs of planar curves. As far as we know, these graphs were first studied in 1959 by S. Benzer [B59], who investigated the topology of genetic structures. Somewhat later they were also considered by F. W. Sinden [S66] in Bell Labs, who was interested in electrical networks realizable by printed circuits. Sinden collaborated with R. L. Graham, who popularized the notion among combinatorists at a conference in Keszthely (Hungary), in 1976 [G78]. Soon after G. Ehrlich, S. Even, and R. E. Tarjan [EET76] studied string graphs (see also

[*]Supported by NSF grant CR-00-98246, PSC-CUNY Research Award 62450-0031 and OTKA-T-032452.
[†]Supported by OTKA-T-032452 and OTKA-T-038397.

[K83] and [EPL72] for a special case). The aim of this paper is to estimate the *number* of different string graphs on n vertices.

To formulate our main result precisely, we have to agree on the terminology. Let G be a simple graph with vertex set $V(G)$ and edge set $E(G)$. A *string representation* of G is an assignment of simple continuous arcs to the elements of $V(G)$ such that two arcs cross each other if and only if the corresponding vertices of G are adjacent. Graph G is a *string graph* if it has a string representation. We assume that any two arcs share only finitely many points and that at each common point the arcs *properly cross*, i.e., one arc passes from one side of the other arc to the other side. An intersection point of two arcs is called a *crossing*.

For any $d > 0$, graph G is a *string graph of rank d* if it has a string representation with the property that any two strings have at most d crossings.

A class $\mathcal{P}$ of labeled graphs, which is closed under isomorphism, is said to be a *property*. A property $\mathcal{P}$ is called *hereditary* if every induced subgraph of every member of $\mathcal{P}$ belongs to $\mathcal{P}$. Let $\mathcal{P}^n$ denote the set of all (labeled) graphs on the vertex set $\{1, 2, \ldots, n\}$ that belong to $\mathcal{P}$. In the combinatorics literature, the function $|\mathcal{P}^n| \leq 2^{\binom{n}{2}}$ is often called the *speed* of property $\mathcal{P}$, and there are several well known estimates on its growth rate as n increases.

Let $\mathcal{S}$ and $\mathcal{S}_d$ denote the classes of all string graphs and all string graphs of rank d, respectively. Clearly, these are hereditary properties and we have $\mathcal{S}_1 \subseteq \mathcal{S}_2 \subseteq \cdots \subseteq \mathcal{S}$. Our first goal is to estimate their speeds.

Theorem 1. *For the number $|\mathcal{S}^n|$ of all string graphs on n labeled vertices, we have*

$$2^{\frac{3}{4}\binom{n}{2}} \leq |\mathcal{S}^n| \leq 2^{\left(\frac{3}{4}+o(1)\right)\binom{n}{2}}.$$

Theorem 2. *For any $d > 0$, the number $|\mathcal{S}_d^n|$ of all string graphs of rank d satisfies $|\mathcal{S}_d^n| \leq 2^{o(n^2)}$.*

We do not have any better lower bound on $|\mathcal{S}_d^n|$ than $2^{\Omega(n \log n)}$, which follows from the fact that the vertex set has this many different permutations.

A *drawing* of a graph is a mapping f which assigns to each vertex of G a distinct point in the plane and to each edge uv a continuous arc between $f(u)$ and $f(v)$, not passing through the image of any other vertex. For simplicity, the point assigned to a vertex is also called a *vertex* and an arc assigned to an edge is also called an *edge* of the drawing, and, if this leads to no confusion, it is also denoted by uv. We assume that (a) two edges have only finitely many points in common, and (b) if two edges share an interior point p, then they properly cross at p. Two drawings of G are said to be *essentially equivalent* if the set of crossing pairs of edges is the same in the two drawings. Otherwise, they are *essentially different.*

Let $\Delta(n)$ and $\bar{\Delta}(n)$ denote the number of essentially different drawings and essentially different straight-line drawings, resp., of the complete graph K_n with n vertices. For any $d > 0$, let $\Delta_d(n)$ denote the number of drawings with the property that any two edges have at most d points in common. Clearly, we have

$$\bar{\Delta}(n) \leq \Delta_1(n) \leq \Delta_2(n) \leq \Delta_3(n) \leq \ldots \leq \Delta(n),$$

for every n.

In Sections 2 and 3, we review the extremal graph theoretic tools used in this paper and establish Theorem 1, respectively. In Section 4 we prove Theorem 2, first in the special case $d = 1$, and then we extend the argument to the general case.

In Section 5, we deduce the following estimates.

Theorem 3. *For the number of essentially different drawings of K_n under various restrictions, we have*
(i) $2^{\Omega(n\log n)} \le \bar{\Delta}(n) \le 2^{O(n\log n)}$;
(ii) $2^{\Omega(n^2)} \le \Delta_1(n) \le 2^{O(n^2\log n)}$;
(iii) $2^{\Omega(n^2\log n)} \le \Delta_d(n) \le 2^{o(n^4)}$, *for any fixed* $d \ge 2$*;*
(iv) $2^{\Omega(n^4)} \le \Delta(n) \le 2^{O(n^4)}$.

2 Tools from extremal graph theory

One of the central questions in extremal graph theory [B78] is the following. Given a graph H, what is the maximum number of edges that a graph of n vertices can have if it does not contain H as a (not necessarily induced) subgraph? This quantity is usually denoted by $\mathrm{ex}(n, H)$.

Obviously, the property that a graph is H-free, is hereditary. Let $\mathrm{Forb}(n, H)$ denote the *speed* of this property, i.e., the number of graphs on n labeled vertices that do not contain H as a subgraph. It turns out that the growth rate of these functions crucially depends on the *chromatic number* $\chi(H)$ of H.

Theorem 2.1. (Erdős-Stone [ES46], Erdős-Simonovits [ES66]) *For any graph H, we have*

$$\mathrm{ex}(n, H) = \left(1 - \frac{1}{\chi(H) - 1}\right)\frac{n^2}{2} + o(n^2).$$

Theorem 2.2. (Erdős-Frankl-Rödl [EFR86]) *For any graph H, we have*

$$\mathrm{Forb}(n, H) = 2^{(1+o(1))\mathrm{ex}(n,H)}.$$

If we want to establish analogous results for graphs containing no *induced* subgraph isomorphic to H, then the first difficulty we have to face is the following: unless H is a complete graph, the maximum number of edges that a graph of n vertices can have without containing an induced copy of H is $\binom{n}{2}$. Thus, Theorem 2.1 does not have a direct analogue. Nevertheless, set

$$\mathrm{ex}^*(n, H) := \left(1 - \frac{1}{\tau(H) - 1}\right)\frac{n^2}{2} + o(n^2),$$

where the relevant quantity, $\tau(H)$, taking the place of the chromatic number is defined as follows.

We say that H is (r, s)*-colorable* for some $0 \le s \le r$ if there is an r-coloring of the vertex set $V(H)$, in which the first s color classes are *cliques* (i.e., induce complete subgraphs) and the remaining $r - s$ color classes are *independent sets* (i.e., induce empty subgraphs). Let $\mathcal{C}(r, s)$ denote the class of all (r, s)-colorable graphs, i.e.,

$$\mathcal{C}(r, s) = \{H \ : \ H \text{ is } (r, s)\text{–colorable}\}.$$

Let $\tau(H)$ be the minimum integer r such that H is (r, s)-colorable for all $0 \le s \le r$. Clearly, we have $\tau(H) \ge \chi(H)$, for every H.

Let $\mathrm{Forb}^*(n, H)$ stand for the number of graphs on n labeled vertices which does contain H as an induced subgraph.