

---

# RECONSTRUCTION OF FUNCTION FIELDS

*by*

Fedor Bogomolov and Yuri Tschinkel

---

ABSTRACT. — We study the structure of abelian subgroups of Galois groups of function fields of surfaces.

## Contents

|   |    |
|---|----|
| Introduction                            | 2  |
| 2. Overview                             | 3  |
| 3. Basic algebra and geometry of fields | 6  |
| 4. Projective structures                | 16 |
| 5. Flag maps                            | 20 |
| 6. Galois groups                        | 29 |
| 7. Valuations                           | 30 |
| 8. A dictionary                         | 33 |
| 9. Flag maps and valuations             | 34 |
| 10. Galois groups of curves             | 36 |
| 11. Valuations on surfaces              | 42 |
| 12. $\ell$ -adic analysis: generalities | 44 |
| 13. $\ell$ -adic analysis: curves       | 52 |
| 14. $\ell$ -adic analysis: surfaces     | 53 |
| 15. Proof                               | 58 |
| References                              | 60 |

---

KEY WORDS AND PHRASES. — Galois groups, function fields.

ect:introduction

## Introduction

We fix two distinct primes  $p$  and  $\ell$ . Let  $k = \bar{\mathbb{F}}_p$  be an algebraic closure of the finite field  $\mathbb{F}_p$ . Let  $X$  be an algebraic variety defined over  $k$  and  $K = k(X)$  its function field. Let  $\mathcal{G}_K^a$  be the abelianization of the pro- $\ell$ -quotient  $\mathcal{G}_K$  of the absolute Galois group of  $K$ . Under our assumptions on  $k$ ,  $\mathcal{G}_K^a$  is a torsion-free  $\mathbb{Z}_\ell$ -module. Let  $\mathcal{G}_K^c$  be its canonical central extension - the second lower central series quotient of  $\mathcal{G}_K$ . It determines the following structure: a fan  $\Sigma_K$  of distinguished (primitive) subgroups of  $\mathcal{G}_K^a$  which are finite rank  $\mathbb{Z}_\ell$ -modules. A topologically noncyclic subgroup  $\sigma \in \Sigma_K$  iff

- $\sigma$  lifts to an abelian subgroup of  $\mathcal{G}_K^c$ ;
- $\sigma$  is maximal: there are no abelian subgroups  $\sigma' \subset \mathcal{G}_K^a$  which lift to an abelian subgroup of  $\mathcal{G}_K^c$  and contain  $\sigma$  as a proper subgroup.

thm:main

**THEOREM 1.** — *Let  $K$  and  $L$  be function fields over algebraic closures of finite fields of characteristic  $\neq \ell$ . Assume that  $K = k(X)$  is a function field of a surface  $X/k$  and that there exists an isomorphism*

$$\Psi = \Psi_{K,L} : \mathcal{G}_K^a \simeq \mathcal{G}_L^a$$

*of abelian pro- $\ell$ -groups inducing a bijection of sets*

$$\Sigma_K = \Sigma_L.$$

*Then, for some  $c \in \mathbb{Z}_\ell^*$ ,  $c\Psi$  is induced by an isomorphism  $\bar{\Psi}$  of the perfect closure of  $K$  with the perfect closure of  $L$ ; the pair  $(c, \bar{\Psi})$  is unique up to*

$$(c, \bar{\Psi}) \mapsto (p^a c, (x \mapsto x^{p^a} \circ \bar{\Psi})).$$

We implement the program outlined in [1] and [2] describing the correspondence between higher-dimensional function fields and their abelianized Galois groups. For results concerning the reconstruction of function fields from their (full) Galois groups (the birational Grothendieck program) we refer to the works of Pop, Mochizuki and Efrat (see [9], [8],[5]).

**Acknowledgments.** Both authors were partially supported by the NSF. The second author was employed by the Clay Mathematics Institute. We are grateful to Laurent Lafforgue and Barry Mazur for their interest and the referee for many useful remarks. Comments by Pierre Deligne were of tremendous help.

sect:ober

## 2. Overview

In this section we outline our strategy of reconstruction, or rather recognition, of the function field  $K$  of an algebraic variety  $X$  over an algebraic closure  $k$  of a finite field from a certain quotient of its Galois group.

Let  $\mathcal{G}_K^a$  be the pro- $\ell$ -quotient of the abelianization

$$G_K/[G_K, G_K],$$

of the absolute Galois group  $G_K = \text{Gal}(\bar{K}/K)$  of  $K$ ,  $\ell \neq \text{char}(k)$ . By Kummer theory,  $\mathcal{G}_K^a$  determines the pro- $\ell$ -completion  $\hat{K}^*$  of the multiplicative group  $K^*$ .

A Galois-theoretic characterization of the field  $K$  involves the recognition of the subgroup  $K^*/k^* \subset \hat{K}^*$ , and of the canonical projective structure, the projectivization of the *additive* group  $K$ , considered as a vector space over  $k$ . The necessary information is encoded in  $\mathcal{G}_K^c$ , the maximal pro- $\ell$ -quotient of

$$G_K/[[G_K, G_K], G_K].$$

Our main Galois-theoretic object is a pair  $(\mathcal{G}_K^a, \Sigma_K)$ , where the *fan*  $\Sigma_K$  is the set of all maximal (by inclusion) topologically noncyclic subgroups  $\sigma \subset \mathcal{G}_K^a$  whose set-theoretic preimage in  $\mathcal{G}_K^c$  is an abelian group. It turns out that the  $\mathbb{Z}_\ell$ -rank of such liftable subgroups  $\sigma \subset \mathcal{G}_K^a$  is bounded by the dimension of  $X$ ,

Theorem 1 states that if for two function fields  $K = k(X)$ ,  $L = l(Y)$ , where  $X/k$  is an algebraic surface,  $Y/l$  an algebraic variety,  $k$  and  $l$  are algebraic closures of finite fields of characteristic  $\neq \ell$  there is an isomorphism

$$\Psi : (\mathcal{G}_K^a, \Sigma_K) \rightarrow (\mathcal{G}_L^a, \Sigma_L)$$

then  $k \simeq l$ ,  $Y$  is a surface and  $L$  is isomorphic to a purely inseparable extension of  $K$ .

Define a subfan  $\Sigma_K^{\text{div}} \subset \Sigma_K$  as the set of those maximal liftable subgroups which have nontrivial intersection with at least one other subgroup in  $\Sigma_K$ . There is a geometric reason to distinguish  $\Sigma_K^{\text{div}}$ . Let  $K$  be the function field

of a surface  $X$  over  $k$ ,  $D$  an irreducible divisor on  $X$  and  $\nu = \nu_D$  the corresponding nonarchimedean divisorial valuation. Its abelian decomposition group  $\mathcal{D}_\nu^a \subset \mathcal{G}_K^a$  is a (noncanonical) direct product of the inertia subgroup  $\mathcal{I}_\nu^a \simeq \mathbb{Z}_\ell$  and the group  $\mathcal{G}_{k(D)}^a$  of the field  $k(D)$ . Now a subgroup  $\sigma \subset \mathcal{D}_\nu^a$  of  $\mathbb{Z}_\ell$ -rank 2 is liftable if and only if it contains  $\mathcal{I}_\nu^a$ . Thus  $\Sigma_K^{\text{div}}$  contains all liftable subgroups of  $\mathbb{Z}_\ell$ -rank 2 which are contained in groups  $\mathcal{D}_\nu^a$ .

The first important result says that  $\Sigma_K^{\text{div}}$  exactly coincides with the set of all liftable subgroups of  $\mathbb{Z}_\ell$ -rank 2 contained in the groups  $\mathcal{D}_\nu^a$ , for different  $\nu = \nu_D$ . This gives an purely group-theoretic description of the groups  $\mathcal{D}_\nu^a$ : the nontrivial intersection of two liftable groups  $\sigma, \sigma'$  is always  $\mathcal{I}_\nu^a$ , for some divisorial valuation  $\nu = \nu_D$ , and  $\mathcal{D}_\nu^a$  “centralizes”  $\mathcal{I}_\nu^a$ , it consists of all those elements in  $\mathcal{G}_K^a$  which commute with  $\mathcal{I}_\nu^a$ , after lifting to  $\mathcal{G}_K^c$ .

The proof is based on Kummer theory and the interpretation of  $\mathcal{G}_K^a$  as a space of special (logarithmic)  $\mathbb{Z}_\ell$ -valued maps on the infinite-dimensional projective space  $\mathbb{P}_k(K) = K^*/k^*$  over  $k$ . The description of liftable subgroups is then reduced to questions in finite-dimensional projective geometry. Complete proofs of these results for  $K = \mathbb{F}_q(X)$  are contained in Section 5. The case of arbitrary algebraically closed ground fields  $k$  is treated in [3],[2].

At this stage we characterized all pairs  $\mathcal{D}_{\nu_D}^a, \mathcal{I}_{\nu_D}^a$  inside  $\mathcal{G}_K^a$ , or, vaguely speaking, we recovered “all curves” on all models  $X$  of  $K$  (and  $Y$  of  $L$ ). Next we recover the “points” on  $D$ , as inertia groups  $\mathcal{I}_w^a \subset \mathcal{G}_{k(D)}^a$ , using various subgroups  $\mathcal{I}_{\nu_{D'}}^a$ , as follows: the image of  $\mathcal{I}_w^a$  under any homomorphism of  $\mathcal{G}_K^a$  to a finite group, which is trivial on  $\mathcal{I}_{\nu_D}^a$ , coincides with the image of some  $\mathcal{I}_{\nu_{D'}}^a$  (see Section 10). Conversely, for any  $\gamma \in \mathcal{D}_{\nu_D}^a \setminus \mathcal{I}_w^a$ , for some divisorial valuation on  $k(D)$  (a point), there exists such a homomorphism with the property that the image of  $\gamma$  is not contained in the image of any inertia subgroup  $\mathcal{I}_{\nu_{D'}}^a$ .

Now we can recover the genus of  $D$  and distinguish the set of divisorial valuations of  $K$  which on some model of  $K$  are represented by curves of genus  $> 0$ . Note that these valuations have 1-dimensional centers on *every* model of  $K$ .

We switch our attention to the dual space  $\hat{K}^*$  of  $\mathcal{G}_K^a$ . We seek to recover the subset  $L^*/l^* \subset \hat{K}^* = \hat{L}^*$  using Galois-theoretic data. This is done in several steps, each time obtaining a smaller subgroup:

- $\mathcal{FS}(K) \subset \hat{K}^*$  - elements in  $\hat{K}^*$  with finite nonrational support,
- $K^*/k^* \otimes \mathbb{Z}_\ell \subset \mathcal{FS}(K)$  and
- $K^*/k^* \otimes \mathbb{Z}_{(\ell)}$ .

Elements of  $\hat{K}^*$  can be thought of as infinite products of elements  $f_i^{\ell^i} \in K^*$ , modulo natural identifications, and they can be represented by, in general, infinite sums of irreducible divisors on a projective model  $X$  of the field with  $\mathbb{Z}_\ell$ -adic coefficients which converge to 0 in the  $\ell$ -adic topology. The subgroup  $\mathcal{FS}(K) \subset \hat{K}^*$  consists of elements whose support contains only finitely many nonrational divisors (characterized above). Of course,  $\mathcal{FS}(K)$  contains  $K^*/k^*$  and  $L^*/l^*$  but it is still rather big - elements in  $\mathcal{FS}(K)$  may have infinite rational “tails” if  $X$  contains infinitely many rational curves.

Next we use an  $\ell$ -adic analog of a symbol  $(f, g) \bmod \ell^n \in K_2(K)/\ell^n$ . Notice that  $(f, g) = 0 \bmod \ell^n$  for any  $n \in \mathbb{N}$  if  $f, g$  belong to the same one-dimensional subfield in  $K$ . In particular, for any  $f \in K \setminus k$  there is an element  $g$  which is not a power of  $f$  and such that  $(f, g) = 0$  (we can take  $g = f + 1$ ). This imposes a strong condition on  $f$  since for a generic element in  $\hat{K}^*$  the “commutator” of  $f$  consists of  $\ell$ -adic powers of  $f$  only. This allows us to characterize

$$K^*/k^* \otimes \mathbb{Z}_\ell = L^*/l^* \otimes \mathbb{Z}_\ell \subset \mathcal{FS}(K).$$

The next step involves a normalization. Inside  $K^*/k^* \otimes \mathbb{Z}_\ell$  we cannot Galois-theoretically distinguish  $L^*/l^* \otimes \mathbb{Z}_{(\ell)}$  from  $a \cdot K^*/k^* \otimes \mathbb{Z}_{(\ell)}$ , for  $a \in \mathbb{Z}_\ell^*$ . However, this conformance invariance is the only freedom there is. If we fix the values of  $f \in L^*/l^* \otimes \mathbb{Z}_\ell$  on one (arbitrary) irreducible divisor on a model  $X$  of  $K$  then  $L^*/l^* \otimes \mathbb{Z}_{(\ell)}$  is naturally identified inside  $K^*/k^* \otimes \mathbb{Z}_\ell$ . Thus, after multiplication by  $a \in \mathbb{Z}_\ell^*$ , we can assume that  $L^*/l^* \subset K^*/k^* \otimes \mathbb{Z}_{(\ell)}$ .

Now we have  $K^*/k^*$  and  $L^*/l^*$  inside  $K^*/k^* \otimes \mathbb{Z}_{(\ell)} = L^*/l^* \otimes \mathbb{Z}_{(\ell)}$ . We also know that subgroups generated by elements  $f, g$  with pairwise trivial symbol  $(f, g) = 0$  correspond to one-dimensional subfields in  $K, L$ , respectively.

Most one-dimensional subfields in  $K$  are isomorphic to  $k(x)$ , for some  $x$ , and Galois data allow us to recognize these subfields. Hence if  $k(x) \subset K$  then  $k(x)^*/k^* \otimes \mathbb{Z}_{(\ell)} = l^*(t)/l^* \otimes \mathbb{Z}_{(\ell)} \subset K^*/k^* \otimes \mathbb{Z}_{(\ell)}$ , for some  $t \in L$ .

Next we show that the corresponding groups  $k(x)^*/k^*$  and  $l^*(x)/l^*$  intersect in  $k(x)^*/(k^*)^r = l^*(x)/(l^*)^s$  for some rational  $r, s$ . This property implies that  $L^*/l^*$  is isomorphic (as a multiplicative group) to  $K_1^*/k^*$  where  $K/K_1$  is a purely inseparable extension.

Now we add the projective structure over  $k, l$ , respectively. We notice that the sets of lines  $\{\mathbb{P}(k \oplus kx)\}$  and  $\{\mathbb{P}(l \oplus lt)\}$  in  $K^*/k^*$  and  $L^*/l^*$ , over all  $x, t$  generating closed subfields  $k(x) \subset K$  and  $l(t) \subset L$ , are the same. It turns out that the set of these lines and their (multiplicative) translations is compatible with a unique projective structure on the (multiplicative) groups  $K^*/k^*$  and  $L^*/l^*$  - namely the one coming from the field structure. This defines a unique additive structure and finishes the proof of our main result.

### 3. Basic algebra and geometry of fields

sect:basicalg

nota:k

NOTATIONS 3.1. — Throughout,  $k$  is an algebraic closure of the finite field  $\mathbb{F} = \mathbb{F}_p$  and  $K = k(X)$  the function field of an algebraic variety  $X/k$  over  $k$  (its *model*).

In this paper we use the fact that two-dimensional function fields  $K$  have “nice” models: smooth projective surfaces  $X$  over  $k$  with  $K = k(X)$ , whose geometric properties play an important role in the recognition procedure. In this section we collect some technical results about function fields of curves and surfaces and their models.

We assume familiarity with

- basic notions of field theory (transcendence degree, purely inseparable extensions);
- basic notions of algebraic geometry:  $k$ -rational points  $X(k)$ , Picard group  $\text{Pic}(X)$ , Néron-Severi group  $\text{NS}(X)$ .

lemm:C

LEMMA 3.2. — *Let  $C/k$  be a smooth curve and  $Q \subset C(k)$  a finite set. Then there exists an  $n = n_Q \in \mathbb{N}$  such that for every degree zero divisor  $D$  with support in  $Q$  the divisor  $nD$  is principal.*

*Proof.* — Finitely generated subgroups of torsion groups are finite. The group of degree zero divisors  $\text{Pic}^0(C)$  (over any finite field) is torsion and every subgroup of divisors with support in a finite set  $Q \subset C(k)$  is finitely generated.  $\square$

lemm:ress

LEMMA 3.3. — *Let  $X/k$  be a surface,  $C_1, \dots, C_s$  a finite set of (pairwise distinct) curves on  $X$  and  $f_i \in k(C_i)^*$ , for  $i = 1, \dots, s$ . Then there exists an  $f \in k(X)^*$  whose restriction to  $C_i$  is defined and equal to  $f_i$ , for all  $i$ .*

*Proof.* — Well known.  $\square$

lemm:new

LEMMA 3.4. — *For some ample smooth curve  $i : C \hookrightarrow X$  the restriction  $i^* : \text{Pic}^0(X) \rightarrow \text{Pic}^0(C)$  is an injection of abstract groups (of  $k$ -points).*

*Proof.* — Let  $H$  be a polarization on  $X$ . There exists an  $n \in \mathbb{N}$  such that for all pairs  $L, L' \in \text{Pic}^0(X)$  one has  $H^1(X, -(L - L' + nH)) = 0$ . Indeed, the property

$$H^1(X, -(L - L' + mH)) = 0$$

is open in  $\text{Pic}^0(X) \times \text{Pic}^0(X) \setminus \Delta$  (where  $\Delta$  the diagonal subgroup), since  $\text{Pic}^0(X)$  is an algebraic group scheme. Denote by

$$U_{mH} \subset (\text{Pic}^0(X) \times \text{Pic}^0(X) \setminus \Delta)$$

the corresponding subset. If we consider a increasing sequence

$$U_{n_i H}, \quad n_i \in \mathbb{Z}, \quad U_{mH} \subset U_{nH}, \quad \text{for } m < n,$$

the union of all  $U_{n_i H}$  is equal to  $\text{Pic}^0(X) \times \text{Pic}^0(X) \setminus \Delta$ . Then there is an  $n \in \mathbb{N}$  such that  $U_{nH} = \text{Pic}^0(X) \times \text{Pic}^0(X) \setminus \Delta$  (due to algebraicity of  $\text{Pic}^0(X) \times \text{Pic}^0(X) \setminus \Delta$  and all  $U_{n_i H}$ ). Exact sequence in cohomology gives:

$$H^0(X, -L + L') = H^0(C_n, -L + L'),$$

where  $C_n$  is a smooth curve in the class  $[nH]$ . Since  $H^0(X, -L + L') = 0$ , for  $L \neq L'$ , the same holds for their restrictions. In particular,

$$i^* : \text{Pic}^0(X) \rightarrow \text{Pic}^0(nH)$$

is a set-theoretic embedding (on the set of  $k$ -points).  $\square$

rem:str

REMARK 3.5. — A more delicate analysis shows that for  $n \gg 0$  the map  $i^* : \text{Pic}^0(X) \rightarrow \text{Pic}^0(nH)$  is an embedding of algebraic groups. Note that over a closure of a finite field the map

$$i : \text{Pic}(X) \rightarrow \text{Pic}(C)$$

is never an embedding if  $\text{rk NS}(X) > 1$  (in contrast with characteristic zero).

lemm:gogo

LEMMA 3.6. — *Let  $X/k$  be a smooth projective surface,  $C \subset X$  an irreducible curve and  $Q$  a finite set of points on  $C$ . Then there exists a diagram*

$$\begin{array}{ccc} \tilde{C} \subset \tilde{X} & \xrightarrow{\pi} & Y \\ \downarrow \tilde{\pi} & & \\ C \subset X & & \end{array}$$

where  $\tilde{X} = \text{Bl}(X) \rightarrow X$  is a blowup of  $X$  with center supported in  $C \setminus Q$  and  $\pi$  is an isomorphism on  $\tilde{X} \setminus \tilde{C}$  (the strict transform of  $C$  under  $\tilde{\pi}$ ) which maps  $\tilde{C}$  to a point on  $Y$ .

*Proof.* — There is a polarization  $H$  on  $X$  such that  $H - C$  restricted to  $X \setminus C$  is very ample (induces an embedding of  $X \setminus C$  into a projective space). Let  $\mathbb{P}_C^r$  be the projective space spanned by  $C$  under the embedding  $X \subset \mathbb{P}^n$  by  $H$ . By our choice of  $H$ ,  $r < n$ . A generic hyperplane  $\mathbb{P}^{r-1} \subset \mathbb{P}_C^r$  intersects  $C$  transversally in finitely many smooth points  $q_1, \dots, q_s$  which are contained in  $C \setminus Q$  (here we use Bertini's theorem for embedded curves, which in this case is evident over any algebraically closed field). The projection from this  $\mathbb{P}^{r-1}$  (inside  $\mathbb{P}^n$ ) induces a proper map from the blowup  $\tilde{X}$  of  $X$  with center in  $\cup q_i$  onto a projective surface  $Y \subset \mathbb{P}^{n-r}$ . Note that the image of  $C$  under the projection is a point  $q \in Y$ .

By construction,  $\mathbb{P}_C^r$  intersects  $X$  exactly in  $C$ . Hence, the proper preimage of  $q$  in  $\tilde{X}$  is  $\tilde{C}$ . Any other  $\mathbb{P}^r \subset \mathbb{P}^n$  intersects  $X \setminus C$  in at most one point and transversally (by assumption on  $H^0(X, H - C)$ ). It follows that the projection induces an isomorphism between  $\tilde{X} \setminus \tilde{C}$  and  $Y \setminus q$ .  $\square$

lemm:go

LEMMA 3.7. — *Let  $X/k$  be a smooth projective surface,  $C \subset X$  a curve and  $Q \subset C(k)$  a finite set. Let  $\mathcal{L}$  be a line bundle on  $X$  whose restriction to  $C$  is trivial ( $\mathcal{L}_C \simeq \mathcal{O}_C$ ). Then there exists a diagram*

$$\begin{array}{ccc} \tilde{C} \subset \tilde{X} & \xrightarrow{\pi} & Y \\ \downarrow \tilde{\pi} & & \\ C \subset X & & \end{array}$$



where  $\tilde{X} = \text{Bl}(X) \rightarrow X$  is a blowup with center supported in finitely many points on  $C \setminus Q$  and  $\pi$  is a proper map as in Lemma 3.6 (contracting  $\tilde{C}$ ) such that the pullback  $\tilde{\mathcal{L}} = \tilde{\pi}^* \mathcal{L}$  is induced from  $Y$ .

*Proof.* — By Lemma 3.6, we may assume that  $C$  is (already) contractible. Since  $\mathcal{L}$  is trivial on  $C$  we have  $\mathcal{L} \simeq \mathcal{O}(R_1 - R_2)$ , where  $R_1, R_2$  are divisors on  $X$  intersecting  $C$  transversally, and

$$R_1 \cap C = R_2 \cap C \subset C \setminus Q.$$

Indeed, we can find a polarization  $H$ , so that  $\mathcal{L} + H$  is also a polarization, giving surjective maps

$$\begin{array}{ccc} H^0(X, \mathcal{L} + H) & \rightarrow & H^0(C, (\mathcal{L} + H)_C) \\ H^0(X, H) & \rightarrow & H^0(C, H_C). \end{array}$$

Let  $i_C : (\mathcal{L} + H)_C \xrightarrow{\sim} H_C$  be an isomorphism. We can find a pair of sections

$$s_1 \in H^0(X, \mathcal{L} + H), \quad s_2 \in H^0(X, H)$$

with  $i_C(s_1)_C = (s_2)_C$ . Let  $R_i$  be the zero divisor of  $s_i$ . Then  $\mathcal{O}(R_1 - R_2) \simeq \mathcal{L}$  and  $R_1, R_2$  intersect  $C$  transversally with

$$R_1 \cap C = R_2 \cap C \subset C \setminus Q,$$

as claimed.

Consider the smooth surface  $\tilde{\pi} : \tilde{X} \rightarrow X$  obtained by blowing up  $R_i \cap C$ . The proper preimages  $\tilde{R}_i$  of  $R_i$  in  $\tilde{X}$  don't intersect the proper preimage  $\tilde{C} \subset \tilde{X}$  of  $C$ . The divisor of  $\pi^* \mathcal{L} = \pi^*(R_1 - R_2)$  doesn't contain components which are exceptional curves lying over points in  $C$ . Hence  $\pi^* \mathcal{L}$  is trivial on the open quasi-projective neighborhood  $\tilde{X} \setminus \text{supp}(\pi^*(R_1 - R_2))$  containing  $\tilde{C}$ . Therefore, the bundle  $\mathcal{L}$  is induced from  $Y$  (as in Lemma 3.6).  $\square$

lemm:cc

LEMMA 3.8. — *Let  $K/k$  be the function field of a surface,  $C/k$  a smooth curve on a model of  $K$  and  $Q = \{q_0, \dots, q_s\} \subset C(k)$  a finite set of points. Then there exist a model  $X$  of  $K$ , irreducible divisors  $D_j, H_j, H'_j$  on  $X$ , with  $j = 0, \dots, s$ , and a positive integer  $n = n_Q$  such that:*

- (1)  $X$  is smooth and contains  $C$ ;
- (2)  $D_j \cap C = q_j$  for all  $j = 1, \dots, s$ ;
- (3)  $n(D_j - D_0)$  restricted to  $C$  is a principal divisor;
- (4)  $n(D_j - D_0) + (H_j - H'_j)$  is a principal divisor on  $X$ ;
- (5) the divisors  $D_j$  are pairwise disjoint;

- (6) *all intersections between  $D_j, H_i$  and  $H'_i$  are transversal, pairwise distinct and outside  $C$ ;*
- (7)  *$H_j, H'_j$  don't intersect  $C$ .*

*Proof.* — Let  $X$  be a smooth projective model of  $K$  containing  $C$  as a smooth curve. Choose divisors  $D_j \subset X$  passing (transversally) through  $q_j$  (for all  $j = 0, \dots, s$ ). Blowing up points in  $C \setminus Q$  we can insure that the (strict transform of)  $C$  becomes contractible and that the image of the surface under a contracting morphism is *projective* (by Lemma 3.6).

Blowing up again (if necessary) and removing components of exceptional divisors, we can insure that the (strict transforms)  $D_j \cap C = q_j$  (for all  $j$ ). By Lemma 3.2, there exists an  $n = n_Q$  such that the restriction of  $n(D_j - D_0)$  to  $C$  is a principal divisor. We continue to blow up (outside  $Q$ ) so that each  $n(D_j - D_0)$  becomes a trivial line bundle on some open neighborhood of  $C$  in some model  $X$  (using Lemma 3.7).

Throughout,  $C$  remains contractible and we write

$$\pi : X \rightarrow Y$$

for the corresponding blow-down. Now  $n(D_j - D_0)$  is induced from a line bundle on  $Y$  (which is projective). In particular, there exist *ample* classes  $[H_j], [H'_j] \in \text{Pic}(Y)$  such that

$$[n(D_j - D_0)] + ([H_j] - [H'_j])$$

is a principal divisor on  $X$  (here we identified  $[H_j], [H'_j]$  with their full transforms in  $X$ ). Finally, we can choose representatives  $H_j, H'_j \subset Y$  of these classes which are disjoint from  $\pi(C)$ , irreducible and satisfy all required transversality assumptions.

More precisely, choose classes  $[H_i]$  so that

$$[n(D_j - D_0)] + ([H_j]), \quad [n(D_j - D_0)] + ([H'_j]), \quad [H_j], [H'_j]$$

provide an embedding of  $Y$  into a projective space. Consider an embedding of  $Y$  into a projective space defined by one of the series  $[H_j], [H'_j]$ . For any finite set of irreducible divisors in  $Y$  we find a hyperplane section intersecting the union of these divisors transversally and not containing the given finite set of points in  $Y$ . Using induction on  $j$  we find representatives of  $[H_j], [H'_j]$  satisfying the lemma.  $\square$

lemm:purely

LEMMA 3.9. — *Let  $K/\mathfrak{K}$  be a purely inseparable extension. Then*

- $\mathfrak{K} \supset k$ ;
- $K/\mathfrak{K}$  is a finite extension;
- $\mathfrak{K} = k(X')$  for some algebraic variety  $X'$ .

**defn:gener**

**DEFINITION 3.10.** — We write  $\overline{E}^K \subset K$  for the normal closure of a subfield  $E \subset K$  (elements in  $K$  which are algebraic over  $E$ ). We say that  $x \in K \setminus k$  is generating if  $\overline{k(x)}^K = \overline{E}^K$ .

**rem:just**

**REMARK 3.11.** — If  $E \subset K$  is 1-dimensional then for all  $x \in E \setminus k$  one has  $\overline{k(x)}^K = \overline{E}^K$  (a finite extension of  $E$ ).

**lemm:subf**

**LEMMA 3.12.** — For any subfield  $E \subset K$  there is a sequence

$$X \xrightarrow{\pi_E} Y' \xrightarrow{\iota_E} Y,$$

where

- $\pi_E$  is rational dominant with irreducible generic fiber;
- $\iota_E$  is quasi-finite and dominant;
- $k(Y') = \overline{E}^K$  and  $k(Y) = E$ .

For generating  $x \in K$  we write

$$\pi_x : X \rightarrow Y$$

for the morphism from Lemma 3.12, with  $k(Y) = k(x)$ . For  $y \in K \setminus k(x)$  define  $\deg_x(y)$  (the degree of  $y$  on the generic fiber of  $\pi_x$ ) as the degree of the corresponding surjective map from the generic fiber of  $\pi_x$  under  $\pi_y$ .

**prop:geome**

**PROPOSITION 3.13.** — Let  $K = k(X)$  be the function field of a smooth surface,  $C \subset X$  a smooth irreducible curve and  $f_1, \dots, f_s \in K^*$  rational functions on  $X$ , restricting nontrivially to  $C$ . Then there exists a model  $\tilde{X}$  of  $K$  (a blowup of  $X$ ) such that for every point  $q$  in (the strict transform) of  $C \subset \tilde{X}$  there exists an irreducible divisor  $D_q \subset \tilde{X}$  (possibly the zero divisor) with the property that for all  $i = 1, \dots, s$  the order of  $D_q$  in the divisor of  $f_i$  is equal to the order of  $f_i$  in  $q$ .

*Proof.* — Consider the divisors  $C$  and  $\text{div}(f_i)$ ,  $i = 1, \dots, s$  and a model  $\tilde{X}$  of  $K$  such that the total preimage  $\tilde{D} \subset \tilde{X}$  of the union of all these divisors in  $\tilde{X}$  has strict normal crossings (resolution of singularities for surfaces). After further blowups we can assume that each irreducible component of  $\tilde{D}$  (distinct from  $C$ ) intersects (the strict transform of)  $C$  in at most one point. For each

$q \in C \cap (\tilde{D} \setminus C)$  let  $D_q$  be this component. For all other  $q$  let  $D_q$  be the zero divisor. These divisors have the required properties.  $\square$

lemm:XX

LEMMA 3.14. — *Let  $K = k(X)$  be the function field of a surface and  $x, y \in K \setminus k$  be such that*

$$\deg_x(y) = \min_{f \in K \setminus \overline{k(x)}^K} (\deg_x(f))$$

*and  $\overline{k(y)}^K = k(y')$  for some  $y' \in K^*$ . Then  $y$  is generating:  $k(y) = \overline{k(y)}^K$ .*

*Proof.* — If  $y$  is not generating then  $y = z(y')$  for some  $y' \in K$  and some function  $z \in k(y')^*$  of degree  $\geq 2$ . This implies that  $\deg_x(y) \geq 2 \deg_x(y')$ , contradicting minimality.  $\square$

lemm:xCxC

LEMMA 3.15. — *Let  $X$  be a model of  $K$  containing a rational curve  $C$  and  $x \in K^*$  a function such that its restriction  $x_C$  to  $C$  is defined and such that  $k(C) = k(x_C)$ . Then  $x$  is generating:  $\overline{k(x)}^K = k(x)$ .*

*Proof.* — The restriction map extends to  $\overline{k(x)}^K$  and hence is an isomorphism between  $k(x_C)$  and  $k(x) = \overline{k(x)}^K$ .  $\square$

The next proposition characterizes multiplicative groups of fields  $\mathfrak{K} \subset K$  such that  $K/\mathfrak{K}$  is a purely inseparable extension. Notice that for a one-dimensional field  $k(C)$  the subfield  $\mathfrak{K}$  is always of the form  $k(C)^{p^n}$ , for some  $n \in \mathbb{N}$ . Thus for any one-dimensional subfield  $E \subset K$  there is an  $r(E) \in \mathbb{N}$  such that the intersection of  $\mathfrak{K}^*$  with  $E^*$  consists exactly of  $r(E)$ -powers of the elements of  $E^*$ . Below we show that this property of intersection with subfields of the special form  $k(x) = \overline{k(x)}^K$  already characterizes multiplicative groups of such  $\mathfrak{K}$  among multiplicative subsets in  $K^*$ .

defi:new

DEFINITION 3.16. — *Let  $\mathfrak{K}^* \subset K^*$  be a (multiplicative) subgroup such that for any subfield  $E = k(x) = \overline{k(x)}^K \subset K$  there exists an  $r = r(E)$  with the property that  $\mathfrak{K}^* \cap E^* = (E^*)^r$  ( $r$ -powers of elements of  $E^*$ ). For every  $t \in E^* \setminus k^*$  we define  $r(t) = r(E)$ .*

REMARK 3.17. — Note that  $r(t)$  is not defined for  $t \in K^* \setminus k^*$  iff  $\overline{k(t)}^K$  is the function field of a curve of genus  $\geq 1$ .

`defi:power`

DEFINITION 3.18. — We will say that  $y \in K^*$  is a power if there exist an  $x \in K^*$  and an integer  $n \geq 2$  such that  $y = x^n$ .

`prop:KKK`

PROPOSITION 3.19. — Let  $K = k(X)$  be the function field of a surface and  $\mathfrak{K}^* \subset K^*$  a subset such that

- (1)  $\mathfrak{K}^*$  is a multiplicative subgroup of  $K^*$ ;
- (2) for every  $E = k(x) = \overline{k(x)}^K \subset K$  there exists an  $r = r(E) \in \mathbb{N}$  with
$$\mathfrak{K}^* \cap E^* = (E^*)^r;$$
- (3) there exists a  $y \in K \setminus k$  with  $r(y) = 1$ .

Then  $\mathfrak{K} := \mathfrak{K}^* \cup 0$  is a field, whose multiplicative group is  $\mathfrak{K}^*$  and  $K/\mathfrak{K}$  is a purely inseparable finite extension.

*Proof.* — Once we know that  $\mathfrak{K}$  is a field we can conclude that every  $x \in K^*$  is either in  $\mathfrak{K}^*$  or some power of it is in  $\mathfrak{K}^*$ . Of course, it can only be a power of  $p$  so that  $K/\mathfrak{K}$  is a purely inseparable extension, of finite degree (by Lemma 3.9).

By (3),  $k \subset \mathfrak{K}$ . To conclude that  $\mathfrak{K}$  is a field, it suffices to show that for every  $x \in \mathfrak{K}$  one has  $x + 1 \in \mathfrak{K}$  (and then use multiplicativity). For every  $x \in \mathfrak{K} \setminus k$  with  $r(x) = 1$  we have  $\mathfrak{K}^* \cap k(x)^* = k(x)^*$  and

$$x + \kappa \in \mathfrak{K}^*, \text{ for all } \kappa \in k.$$

In particular, this holds for  $y$ .

Consider  $x \in \mathfrak{K}^*$  with  $r(x) > 1$  or not defined. We claim that for some  $\kappa \in k$

$$z := \frac{x + y + \kappa}{y + \kappa - 1} \in \mathfrak{K} \text{ and } r(z) = 1.$$

This implies that

$$z - 1 = (x + 1)/(y + \kappa - 1) \in \mathfrak{K}^* \text{ and } x + 1 \in \mathfrak{K}^*,$$

(by multiplicativity). We can assume that  $K/k(C)(y)$ , where  $k(C) = \overline{k(x)}^K$ , is a finite separable extension. (Otherwise, we can let  $K$  be a minimal proper subfield in  $\mathfrak{K}' \subset K$  containing  $k(C)(y)$  and such that  $K/\mathfrak{K}'$  is purely inseparable and use the intersection of  $\mathfrak{K}$  with  $\mathfrak{K}'$  instead of  $\mathfrak{K}$ .)

To prove the claim, choose a model  $X$  of  $K$  such that both maps

$$\begin{aligned} \pi_x : X &\rightarrow C, & k(C) &= \overline{k(x)}^K \\ \pi_y : X &\rightarrow \mathbb{P}^1 = (y : 1) \end{aligned}$$

are proper morphisms (as in Lemma 3.12). Since  $x$  and  $y$  are algebraically independent ( $r(x) > 1$ ), only finitely many components of the fibers of  $\pi_x$  are contained in the fibers of  $\pi_y$  and there exists a  $\kappa \in k$  such that both fibers

$$\pi_y^{-1}(-\kappa) \text{ and } \pi_y^{-1}(1 - \kappa)$$

are transversal to the fibers of  $\pi_x$ , since we assume that  $K/k(C)(y)$  is separable. Note that

$$\operatorname{div}_0(y + \kappa - 1) \not\subset \operatorname{div}(x + y + \kappa),$$

since  $y + \kappa = -1$  on  $\operatorname{div}_0(y + \kappa - 1)$  and  $x$  is nonconstant on these fibers (where  $\operatorname{div}_0$  is the divisor of zeroes). It follows in the first case that *both*

$$t := (y + \kappa)/x \text{ and } z := (x + y + \kappa)/(y + \kappa - 1)$$

are not powers.

Note that  $t, z$  are generating elements. Indeed, if we blow up the smooth point  $q$  of transversal intersection  $\{y + \kappa = 0\} \cap \{x = 0\}$  then  $t$  restricts nontrivially to  $\mathbb{P}_q^1$  and similarly

$$z := (x + y + \kappa)/(y + \kappa - 1) = x + 1/(y + \kappa - 1) + 1$$

restricts nontrivially to  $\mathbb{P}_{q'}^1$ , where  $q' = \{x = -1\} \cap \{y = 1 - \kappa\}$ .

Note that  $t \in \mathfrak{K}^*$  and since it is not a power  $r(t) = 1$  and

$$(1/t) + 1 = (x + y + \kappa)/(y + \kappa) \in \mathfrak{K}.$$

To show that  $z \in \mathfrak{K}$  observe that both  $x, y + \kappa \in \mathfrak{K}$  so that  $t \in \mathfrak{K}$ . Therefore,

$$t + 1 = (x + y + \kappa)/x \in \mathfrak{K}$$

and, by (1),  $x + y + \kappa \in \mathfrak{K}$ . Finally, since  $(y + \kappa - 1) \in \mathfrak{K}$  we get  $z \in \mathfrak{K}$ .  $\square$

REMARK 3.20. — If assumption (3) is not satisfied then we can take

$$(\mathfrak{K}^*)^{1/r(y)} \bigcap K^*,$$

which satisfies all the conditions of the lemma. Thus in general without the assumption (3) we have  $\mathfrak{K} = (\mathfrak{K}')^r$ , where  $K/\mathfrak{K}'$  is purely inseparable and  $r \in \mathbb{N}$ .

In our analysis of Galois groups we need to keep track of rational curves on a surface.

lemm:surface

LEMMA 3.21. — *Let  $X$  be a surface over  $k$ . There three mutually disjoint possibilities:*

- (1)  $\text{Pic}^0(X) = 0$ ;
- (2)  $\text{Pic}^0(X) \neq 0$  and  $X$  contains finitely many rational curves;
- (3)  $\text{Pic}^0(X) \neq 0$  and, after a finite purely inseparable extension,  $X$  admits a fibration over a curve  $C$  of genus  $g(C) \geq 1$  with generic fiber a rational curve.

*Proof.* — Follows from the classification of surfaces. Indeed, if  $X$  is smooth and  $\text{Pic}^0(X) \neq 0$  then there is a nontrivial map into the Albanese variety of  $X$ , and all rational curves lie in fibers. The generic fiber of this map is either rational or there are only finitely many rational curves on  $X$ .  $\square$

Let  $X$  be a surface over  $k$  and  $\text{Alb}(X)$  its Albanese variety. Recall that  $\text{Alb}(X)$  is a principal homogeneous space for an abelian variety  $A^0(X)$ , with  $\dim A^0(X) = \dim \text{Alb}(X)$ .

lemm:surface2

LEMMA 3.22. — *Let  $\mathcal{D} := \{D_j\}_{j \in J}$  be a finite set of irreducible divisors on  $X$ . Assume that there is an  $f \in k(X)^*$  whose divisor is supported in  $\mathcal{D}$ . Let  $B \subset A^0$  be the smallest abelian subvariety such that the image of  $D_j$  under the map  $\alpha : \text{Alb}(X) \rightarrow A := \text{Alb}(X)/B$  is a point, for all  $j \in J$ .*

*Assume that  $B \neq A^0(X)$ . Then the image of  $X$  in  $A$  is a curve  $C$  and  $A$  is isomorphic to the Jacobian  $\text{Jac}(C)$  of degree 1 zero-cycles on  $C$ .*

*Proof.* — First of all,  $\dim \alpha(X) \geq 1$ : the surface  $X$  is connected and  $\alpha(X)$  generates  $A$ . Further,  $\alpha(X)$  is not a surface: otherwise if  $X' \rightarrow \alpha(X)$  is the normalization, then there is a map  $\mu : X \rightarrow X'$  and the image of  $\{D_j\}_{j \in J}$  is a finite set of points on  $X'$ . The intersection matrix of the set of irreducible components in the divisorial support of  $\mu^{-1}(x')$ , for any  $x' \in X'$ , is negative definite, contradicting the assumption that there is a function supported in  $\mathcal{D}$ .

Let  $C := \alpha(X) \subset A$ , we have  $k(C) \subset K$ . Let  $C'$  be a curve with function field  $k(C') = \overline{k(C)}^K \subset K$ . The map  $C' \rightarrow C$  is finite. The map  $\alpha : X \rightarrow A$  factors through the Jacobian  $\text{Jac}(C')$ : we have

$$\begin{array}{ccc} X & \xrightarrow{\alpha_{C'}} & \text{Jac}(C') \\ & & \downarrow \\ & & A \end{array}$$

The image of  $\{D_j\}_{j \in J}$  under  $\alpha_{C'}$  is a finite set of points in  $\text{Jac}(C')$ . We have surjections  $\text{Jac}(C') \rightarrow \text{Jac}(C) \rightarrow A$  and a canonical map  $\text{Alb}(X) \rightarrow \text{Jac}(C')$ . Then  $B = \text{Ker}(\alpha_{C'})$  and  $C' = C$ .  $\square$

#### 4. Projective structures

sect:proj-str

In this section we explain the connection between fields and axiomatic projective geometry. We follow closely the exposition in [7].

defi:proj

DEFINITION 4.1. — A projective structure is a pair  $(S, \mathfrak{L})$  where  $S$  is a (nonempty) set (of points) and  $\mathfrak{L}$  a collection of subsets  $\mathfrak{l} \subset S$  (lines) such that

- P1 there exist an  $s \in S$  and an  $\mathfrak{l} \in \mathfrak{L}$  such that  $s \notin \mathfrak{l}$ ;
- P2 for every  $\mathfrak{l} \in \mathfrak{L}$  there exist at least three distinct  $s, s', s'' \in \mathfrak{l}$ ;
- P3 for every pair of distinct  $s, s' \in S$  there exists exactly one

$$\mathfrak{l} = \mathfrak{l}(s, s') \in \mathfrak{L}$$

such that  $s, s' \in \mathfrak{l}$ ;

- P4 for every quadruple of pairwise distinct  $s, s', t, t' \in S$  one has

$$\mathfrak{l}(s, s') \cap \mathfrak{l}(t, t') \neq \emptyset \Rightarrow \mathfrak{l}(s, t) \cap \mathfrak{l}(s', t') \neq \emptyset.$$

For  $s \in S$  and  $S' \subset S$  define the *join*

$$s \vee S' := \{s'' \in S \mid s'' \in \mathfrak{l}(s, s') \text{ for some } s' \in S'\}.$$

For any finite set of points  $s_1, \dots, s_n$  define

$$\langle s_1, \dots, s_n \rangle := s_1 \vee \langle s_2 \vee \dots \vee s_n \rangle$$

(this does not depend on the order of the points). Write  $\langle S' \rangle$  for the join of a finite set  $S' \subset S$ . A finite set  $S' \subset S$  of pairwise distinct points is called *independent* if for all  $s' \in S'$  one has

$$s' \notin \langle S' \setminus \{s'\} \rangle.$$

A set of points  $S' \subset S$  *spans* a set of points  $T \subset S$  if

- $\langle S'' \rangle \subset T$  for every finite set  $S'' \subset S'$ ;
- for every  $t \in T$  there exists a finite set of points  $S_t \subset S'$  such that  $t \in \langle S_t \rangle$ .



A set  $T \subset S$  spanned by an independent set  $S'$  of points of cardinality  $\geq 1$  is called a projective *subspace* of dimension  $|S'| - 1$ .


The axioms imply that projective subspaces of a given projective space  $S$  form a lattice and that the dimension function is well defined, i.e.,

$$\dim(T \cup T') + \dim(T \cap T') = \dim(T) + \dim(T')$$

for all pairs of projective subspaces  $T, T' \subset S$ . Here we put  $\dim(\emptyset) := -1$ .

defn:pappus

DEFINITION 4.2. — *A projective structure  $(S, \mathfrak{L})$  satisfies Pappus' axiom if PA for all 2-dimensional subspaces and every configuration of six points and lines in these subspaces as below*



*the intersections are collinear.*

The main theorem of abstract projective geometry is:

thm:abstr

THEOREM 4.3. — *Let  $(S, \mathfrak{L})$  be a projective structure of dimension  $n \geq 2$  which satisfies Pappus' axiom. Then there exists a vector space  $V$  over a field  $L$  and an isomorphism*

$$\sigma : \mathbb{P}_L(V) \xrightarrow{\sim} S.$$

*Moreover, for any two such triples  $(V, L, \sigma)$  and  $(V', L', \sigma')$  there is an isomorphism*

$$V/L \xrightarrow{\sim} V'/L'$$

*compatible with  $\sigma, \sigma'$  and unique up to homothety  $v \mapsto \lambda v, \lambda \in L^*$ .*

*Proof.* — See [7], Chapter 6. □

defi:inject

DEFINITION 4.4. — *A morphism of projective structures*

$$\rho : (S, \mathfrak{L}) \rightarrow (S', \mathfrak{L}')$$

is an injection of sets  $\rho : S \hookrightarrow S'$  such that  $\rho(\mathfrak{l}) \in \mathfrak{L}'$  for all  $\mathfrak{l} \in \mathfrak{L}$ .

exa:basic

EXAMPLE 4.5. — Let  $k$  be a field and  $\mathbb{P}_k^n$  the usual projective space over  $k$  of dimension  $n \geq 2$ . Then  $\mathbb{P}_k^n$  carries a projective structure: the set of lines is the set of usual projective lines  $\mathbb{P}_k^1 \subset \mathbb{P}_k^n$ .

Let  $K/k$  be an extension of fields (not necessarily finite). Then the set

$$S := \mathbb{P}_k(K) = (K \setminus 0)/k^*$$

carries a natural (possibly, infinite-dimensional) projective structure. Moreover, multiplication by elements in the group  $K^*/k^*$  preserves this structure.

thm:skk

THEOREM 4.6. — *Let  $K/L$  and  $K'/L'$  be field extensions of degree  $\geq 3$  and*

$$\bar{\phi} : S = \mathbb{P}_L(K) \rightarrow \mathbb{P}_{L'}(K') = S'$$

*a bijection of sets which is an isomorphism of abelian groups and of projective structures. Then*

$$L \simeq L' \text{ and } K \simeq K'.$$

*Proof.* — Consider  $V := K$  as a vector space over  $L$ . By Theorem 4.3, to  $S$  there are canonically attached the  $L$ -algebra  $\text{End}(V)$  and  $\text{GL}(V) \subset \text{End}(V)$ , as the set of elements preserving the collineations of the projective space  $S$  (because the action of homotheties on  $\text{End}(V)$  is trivial). This allows to recover the field  $K$  as the subfield of the  $L$ -algebra  $\text{End}(V)$  given by

$$\{0\} \cup \{x \in \text{GL}(V) \subset \text{End}(V) \mid x \text{ induces a group-translation on } S\}.$$

□

defn:basic2

DEFINITION 4.7. — *Let  $K/k$  be the function field of an algebraic variety  $X$  of dimension  $\geq 2$  and  $S = \mathbb{P}_k(K)$  the associated projective structure from Example 4.5. The lines passing through 1 and a generating element of  $K$  (see Definition 3.10) and their multiplicative translations by elements in  $K^*/k^*$  will be called primary.*

lemm:primary-ell

LEMMA 4.8. — *Let  $K = k(X)$  be the function field of a surface. For every line  $\mathfrak{l} = \mathfrak{l}(1, x)$  there exists a  $\mathbb{P}^2 \subset \mathbb{P}_k(K)$  such that all other lines in this  $\mathbb{P}^2$  are primary.*

*Proof.* — Choose a smooth model  $X$  of  $K$  and two points  $q_1, q_2 \in X$  such that  $x(q_1) = 0, x(q_2) = 1$ . Blow up  $q_1, q_2$  and let  $\mathbb{P}_i^1$  be the corresponding exceptional curves. Let  $y \in K^*$  be an element restricting to a generator of  $k(\mathbb{P}_i^1)$ . The restriction map extends to the normal closure  $\overline{k(y)} \subset K$ . Hence the normal closure  $\overline{k(y)} \subset K$  coincides with  $k(y)$ .

To prove that every line  $\mathfrak{l} \neq \mathfrak{l}(1, x) \subset \mathbb{P}^2 = \mathbb{P}_k(k \oplus kx \oplus ky)$  is primary we need to show that  $(y + a + bx)/(y + c + dx)$  is generating, provided  $(a, b) \neq (c, d)$ . If  $a \neq c$  then the restriction of  $(y + a + bx)/(y + c + dx)$  to  $\mathbb{P}_{q_1}^1$  is equal to  $(y + a)/(y + c)$  and hence is a generator of  $k(\mathbb{P}_{q_1}^1)$ . By the argument of the previous lemma,  $(y + a + bx)/(y + c + dx)$  is generating. If  $a = c, b \neq d$  then  $(y + a + bx)/(y + c + dx)$  on  $\mathbb{P}_{q_2}^1$  coincides with  $(y + a + b)/(y + c + d)$  and is also generating since  $a + b \neq c + d$ , by assumption.  $\square$

lemm:4.10

LEMMA 4.9. — Assume that a set  $S$  has two projective structures  $(S, \mathfrak{L}_1)$  and  $(S, \mathfrak{L}_2)$ , both of dimension  $\geq 2$ , and that for some  $\mathbb{P}_1^2$  (in the first projective structure) every line  $\mathfrak{l}_1 \in (\mathfrak{L}_1 \cap \mathbb{P}_1^2)$ , except possibly one line, is also a line in the second structure. Then the set  $\mathbb{P}_1^2$  is a projective plane in the second structure  $(S, \mathfrak{L}_2)$ , projectively isomorphic to  $\mathbb{P}_1^2 \in (S, \mathfrak{L}_1)$ .

*Proof.* — Let  $\hat{\mathbb{P}}_1^2$  be the set of all lines in  $\mathbb{P}_1^2$  and  $\hat{\mathbb{P}}_1^2 \setminus \mathfrak{l}$  the set of lines which remain projective lines in  $\mathbb{P}_2^2$ . Let  $\mathfrak{l}_1, \mathfrak{l}_2, \mathfrak{l}_3$  be three lines from  $\hat{\mathbb{P}}_1^2 \setminus \mathfrak{l}$  which don't have a common intersection point. Then  $\mathfrak{l}_1, \mathfrak{l}_2, \mathfrak{l}_3$  lie in the same plane  $\mathbb{P}_2^2$ . Since every other line  $\mathfrak{l}' \in \hat{\mathbb{P}}_1^2 \setminus \mathfrak{l}$  intersects  $\mathfrak{l}_1, \mathfrak{l}_2, \mathfrak{l}_3$  then  $\mathfrak{l}' \subset \mathbb{P}_2^2$ . Thus all lines from  $\hat{\mathbb{P}}_1^2 \setminus \mathfrak{l}$  are in  $\mathbb{P}_2^2$  which contains all the points of  $\mathbb{P}_1^2$ .

They are isomorphic since it is an isomorphism between lines and every point, except possibly one point, is an intersection of two lines in  $\hat{\mathbb{P}}_1^2 \setminus \mathfrak{l}$ . Since  $\hat{\mathbb{P}}_2^2$  coincides with  $\hat{\mathbb{P}}_1^2$  outside of one point they coincide.  $\square$

coro:final

COROLLARY 4.10. — Let  $K/k$  and  $K'/k'$  be function fields of algebraic surfaces

$$\bar{\phi} : S = \mathbb{P}_k(K) \rightarrow S' = \mathbb{P}_{k'}(K')$$

an isomorphism of (multiplicative) abelian groups inducing a bijection on the set of primary lines in the corresponding projective structures. Then  $\bar{\phi}$  is an isomorphism of projective structures and

$$k \simeq k' \quad \text{and} \quad K \simeq K'.$$

*Proof.* — By Lemma 4.8 and Lemma 4.9  $\bar{\phi}$  induces an isomorphism of projective structures. It remains to apply Theorem 4.6.  $\square$

## 5. Flag maps

sect:af

nota:pr

NOTATIONS 5.1. — We fix two distinct prime numbers  $\ell$  and  $p$ . Let

- $\mathbb{F} = \mathbb{F}_q$  be a finite field with  $q = p^n$  and  $\mathbb{F}^*$  its multiplicative group;
- $\text{Vect}_{\mathbb{F}}$  - the set of finite-dimensional  $\mathbb{F}$ -vector spaces;
- $A$  a vector space over  $\mathbb{F}$  and  $\mathbb{P}(A) = \mathbb{P}_{\mathbb{F}}(A) = (A \setminus 0)/\mathbb{F}^*$  its projectivization;
- $\mathcal{M}(A)$  the set of maps from  $A \setminus \{0\}$  to  $\mathbb{Z}_{\ell}$ ;
- for  $\mu \in \mathcal{M}(A)$  and  $B \subset A$  an  $\mathbb{F}$ -linear subspace,  $\mu_B$  the restriction of  $\mu$  to  $B \setminus \{0\}$ .

defn:homo

DEFINITION 5.2. — A map  $\mu \in \mathcal{M}(A)$  will be called  $\mathbb{F}^*$ -invariant if for all  $a \in A \setminus \{0\}$  and all  $\kappa \in \mathbb{F}^*$  one has

$$\mu(\kappa \cdot a) = \mu(a).$$

defn:inv

DEFINITION 5.3. — A map  $\mu$  on  $A \setminus \{0\}$ , for a (possibly infinite-dimensional) vector space  $A$ , will be called an  $\mathbb{F}$ -flag map, if

- $\mu$  is  $\mathbb{F}^*$ -invariant;
- every finite-dimensional  $\mathbb{F}$ -vector space  $B \subset A$  has a flag of  $\mathbb{F}$ -subspaces

$$B = B_0 \supset B_1 \supset \dots \supset B_d = 0$$

such that  $\mu_B$  is constant on  $B_n \setminus B_{n+1}$ , for all  $n = 0, \dots, d-1$ .

The value of  $\mu$  on  $B = B_0 \setminus B_1$  is called the generic value of  $\mu$  on  $B$ ; we denote it by  $\mu^{\text{gen}}(B)$ . The set of  $\mathbb{F}$ -flag maps will be denoted by  $\Phi_{\mathbb{F}}(A)$ .

exam:main-flag

EXAMPLE 5.4. — Let  $K = k(X)$  be a function field. We can consider it as a vector space over  $k$  or over any of the finite subfields  $\mathbb{F} \subset k$ . Let  $\nu$  be a nonarchimedian valuation on  $K$  and  $\chi : \Gamma_{\nu} \rightarrow \mathbb{Z}_{\ell}$  a homomorphism from the value group of  $\nu$  (see Section 7). Then  $\chi \circ \nu \in \Phi_k(K)$ .

defn:lf

DEFINITION 5.5. — Let  $A$  be an  $\mathbb{F}$ -algebra (without zero-divisors). A map  $\mu \in \mathcal{M}(A)$  will be called logarithmic if

$$\mu(a \cdot a') = \mu(a) + \mu(a'), \text{ for all } a, a' \in A \setminus 0.$$

The set of such maps will be denoted by  $\mathcal{L}_{\mathbb{F}}(A)$ .

Since  $\mathbb{F}$  is torsion, a logarithmic map to  $\mathbb{Z}_\ell$  is  $\mathbb{F}^*$ -invariant.

c:pair

**DEFINITION 5.6.** — *Let  $A$  be an  $\mathbb{F}$ -vector space. Two maps  $\mu, \mu' \in \mathcal{M}(A)$  will be called a  $c$ -pair (commuting pair) if for all two-dimensional  $\mathbb{F}$ -subspaces  $B \subset A$  there exist constants  $\lambda, \lambda', \lambda'' \in \mathbb{Z}_\ell$  (depending on  $B$ ) with  $(\lambda, \lambda') \neq (0, 0)$  such that for all  $b \in B \setminus 0$  one has*

$$\lambda\mu_B(b) + \lambda'\mu'_B(b) = \lambda''.$$

thm:cp

**THEOREM 5.7.** — *Let  $\mathbb{F} \subset k$  be a finite field with  $\#\mathbb{F} \geq 11$ , and  $\mu, \mu' \in \mathcal{L}_\mathbb{F}(K)$  nonproportional maps forming a  $c$ -pair. Then there exists a pair  $(\lambda, \lambda') \in \mathbb{Z}_\ell^2 \setminus (0, 0)$  such that  $\lambda\mu + \lambda'\mu' \in \Phi_\mathbb{F}(K)$ .*

*Proof.* — This is a special case of the main theorem of [3], where it is proved over general ground fields  $k$ . However, the case when  $k = \bar{\mathbb{F}}_q$  is easier. Following the request of the referee, we now give a complete proof in this special case. The main steps in the proof are:

- characterization of flag maps by their restriction to 2-dimensional  $\mathbb{F}$ -linear subspaces, for  $\#\mathbb{F} \geq 11$  (see Lemma 5.16);
- reduction to linear spaces over prime fields, resp.  $\mathbb{F}_4$ , see Lemma 5.18: if  $\mu \notin \Phi_{\mathbb{F}'}(A)$ , for a finite field  $\mathbb{F}'$ , and  $\mu$  is  $\mathbb{F}^*$ -invariant with respect to a large finite extension  $\mathbb{F}/\mathbb{F}'$  then there is a subgroup  $C \simeq \mathbb{F}_p^2 \subset A$ , (resp.  $\mathbb{F}_4^2$ ), so that  $\mu_C \notin \Phi_{\mathbb{F}_p}(C)$ .
- reduction to dimension 3: if the rank two  $\mathbb{Z}_\ell$ -module  $\sigma := \langle \mu, \mu' \rangle$  does not contain a flag map then there is a subgroup  $B \simeq \mathbb{F}_p^3 \subset A$  (resp.  $\mathbb{F}_4^3$ ), such that for any nontrivial  $\mu'' \in \sigma$  there is a proper subspace  $C = C_{\mu''} \subsetneq B$  where  $\mu''_C \notin \Phi_{\mathbb{F}_p}(C)$  (this step uses the logarithmic property);
- geometry of collineations on  $\mathbb{P}^2 = \mathbb{P}_\mathbb{F}(B)$  over prime fields  $\mathbb{F} = \mathbb{F}_p$  (resp.  $\mathbb{F}_4$ ): such subgroups  $B$  cannot exist. This shows the existence of the desired flag map on  $A$ .

□

lemm:linearr

**LEMMA 5.8.** — *If  $A \in \text{Vect}_\mathbb{F}$  and  $\mu \in \Phi_\mathbb{F}(A)$  then there exists a canonical  $\mathbb{F}$ -flag  $(A_n)_{n=0, \dots, d}$  such that*

$$\mu^{\text{gen}}(A_n) \neq \mu^{\text{gen}}(A_{n+1}),$$

*for all  $n = 0, \dots, d-1$ .*

*Proof.* — Put  $A_0 = A$  and let  $A_{n+1}$  be the additive subgroup of  $A_n$  spanned by  $a$  with  $\mu(a) \neq \mu^{\text{gen}}(A_n)$ . Since  $\mu$  is  $\mathbb{F}^*$ -invariant,  $A_{n+1}$  is an  $\mathbb{F}$ -vector space. Indeed, for  $a, a' \in A_{n+1}$  and  $\kappa, \kappa' \in \mathbb{F}^*$  write

$$a = \sum_{i \in I} b_i, \quad a' = \sum_{j \in J} b'_j$$

with finite  $I, J$ . Since

$$\mu(b_i) \neq \mu^{\text{gen}}(A_n), \quad \mu(b'_j) \neq \mu^{\text{gen}}(A_n),$$

for all  $i \in I, j \in J$ , we have

$$\mu(\kappa b_i) = \mu(b_i) \neq \mu^{\text{gen}}(A_n) \quad \text{and} \quad \mu(\kappa' b'_j) = \mu(b'_j) \neq \mu^{\text{gen}}(A_n)$$

so that  $\kappa a + \kappa' a' \in A_{n+1}$ . □

REMARK 5.9. — Since a flag map  $\mu$  is  $\mathbb{F}^*$ -invariant, it defines a unique map on  $(A \setminus \{0\})/\mathbb{F}^* = \mathbb{P}_{\mathbb{F}}(A)$ . Conversely, a map  $\mu$  on  $\mathbb{P}_{\mathbb{F}}(A)$  gives rise to an  $\mathbb{F}^*$ -invariant maps on  $A \setminus \{0\}$ . An  $\mathbb{F}$ -flag map on  $A \in \text{Vect}_{\mathbb{F}}$  defines a flag by projective subspaces on  $\mathbb{P}_{\mathbb{F}}(A)$ . We denote by *generic* elements of  $\mathbb{P}_{\mathbb{F}}(A)$  the image of generic elements from  $A$ .

NOTATIONS 5.10. — We denote by  $\hat{\mathbb{P}}(A) = \hat{\mathbb{P}}_{\mathbb{F}}(A)$  the set of codimension one projective  $\mathbb{F}$ -subspaces of  $\mathbb{P}(A)$ .

defn:duall

DEFINITION 5.11. — Assume that  $A \in \text{Vect}_{\mathbb{F}}$ , and for all codimension one  $\mathbb{F}$ -subspaces  $B \subset A$  one has  $\mu_B \in \Phi_{\mathbb{F}}(B)$ . Define  $\hat{\mu}$  by

$$\begin{aligned} \hat{\mathbb{P}}(A) &\rightarrow \mathbb{Z}_{\ell} \\ B &\mapsto \hat{\mu}(\mathbb{P}(B)) := \mu^{\text{gen}}(B). \end{aligned}$$

lemm:1pt

LEMMA 5.12. — If  $A \in \text{Vect}_{\mathbb{F}}$  and  $\mu \in \Phi_{\mathbb{F}}(A)$  then either  $\hat{\mu}$  is constant on  $\hat{\mathbb{P}}(A)$  or it is constant on the complement to one point.

*Proof.* — Consider the canonical flag  $(A_n)_{n=0,\dots,d}$ . If  $\text{codim}(A_1) \geq 2$  then for every  $\mathbb{P}(B) \in \hat{\mathbb{P}}(A)$  one has  $\mu^{\text{gen}}(B) = \mu^{\text{gen}}(A)$  and  $\hat{\mu}$  is constant. Otherwise,  $\mu^{\text{gen}}(B) = \mu^{\text{gen}}(A)$ , on any  $B \neq A_1$  (and differs at  $\mathbb{P}(A_1) \in \hat{\mathbb{P}}(A)$ ). □

We need the following elementary

lemm:grass

LEMMA 5.13. — *Let  $\mathbb{F} = \mathbb{F}_q$  be a finite field with  $q \geq 11$  and  $\mathbb{P}^m = \mathbb{P}_{\mathbb{F}}^m$ ,  $m \geq 2$  a projective space over  $\mathbb{F}$ . For any four projective hyperplanes and any ten projective subspaces of codimension at least two (all defined over  $\mathbb{F}$ ) there exists a line (over  $\mathbb{F}$ ) not contained in any of the above hyperplanes and not intersecting any of the ten codimension two subspaces.*

*Proof.* — One has

$$\#\mathrm{Gr}(2, m)(\mathbb{F}) \leq \#\mathrm{Gr}(2, m+1)(\mathbb{F})/q^2.$$

The number of  $\mathbb{F}$ -lines intersecting a subspace of codimension two in  $\mathbb{P}_{\mathbb{F}}^m$  is bounded by  $\#\mathrm{Gr}(2, m+1)(\mathbb{F})/q^2$ . Our claim holds for  $q \geq 11$ .  $\square$

lemm:c11

LEMMA 5.14. — *Let  $\mathbb{F} = \mathbb{F}_q$  be a finite field with  $q \geq 11$ ,  $A \in \mathrm{Vect}_{\mathbb{F}}$  and  $\mu \in \mathcal{M}(A)$  an  $\mathbb{F}^*$ -invariant map. Assume that there exist  $\mathbb{F}$ -subspaces  $B_i \subset A$ ,  $\mathrm{codim}(B_i) = 1$ , for  $i = 1, \dots, 4$  such that*

- (1) *either  $\#\{\mu^{\mathrm{gen}}(B_i)\} \geq 3$  or*
- (2)  *$\mu^{\mathrm{gen}}(B_1) = \mu^{\mathrm{gen}}(B_2) \neq \mu^{\mathrm{gen}}(B_3) = \mu^{\mathrm{gen}}(B_4)$ .*

*Then there exists an  $\mathbb{F}$ -subspace  $C \subset A$ ,  $\dim_{\mathbb{F}}(C) = 2$  such that  $\mu_C \notin \Phi_{\mathbb{F}}(C)$ .*

*Proof.* — By Lemma 5.13, there exists a  $\mathbb{P}^1 = \mathbb{P}(C) \in \mathbb{P}(A)$  such that its intersection points with  $\mathbb{P}(B_i)$  are pairwise distinct and generic in the corresponding  $\mathbb{P}(B_i)$  (the nongeneric points of  $\mathbb{P}(B_i)$  are contained in 4 subspaces in  $\mathrm{codim}_{\mathbb{F}} \geq 2$ , the intersections of  $B_i$  give rise to 6 more subspaces). Then either  $\mu$  takes at least three distinct values on  $\mathbb{P}(C)$  or has distinct values in at least two pairs of points. In both cases  $\mu \notin \Phi_{\mathbb{F}}(C)$ .  $\square$

coro:one

COROLLARY 5.15. — *Assume that  $\mu_B \in \Phi_{\mathbb{F}}(B)$  for all  $\mathbb{P}(B) \in \hat{\mathbb{P}}(A)$  (and  $\#\mathbb{F} \geq 11$ ). Then  $\hat{\mu}$  is constant outside of one point.*

*Proof.* — The map  $\hat{\mu}$  takes two different values on  $\hat{\mathbb{P}}(B)$ . By Lemma 5.14, among any three hyperplanes two have the same generic value, so that there can be at most three such values. If there are hyperplanes  $h_1, h_2, h_3 \in \hat{\mathbb{P}}(A)$ , where  $\hat{\mu}(h_1) = \hat{\mu}(h_2) \neq \hat{\mu}(h_3)$  then for any other  $h \in \hat{\mathbb{P}}(A)$  we have  $\hat{\mu}(h) = \hat{\mu}(h_1)$  and  $\hat{\mu}$  is constant outside of  $h_3$ .  $\square$

lemm:1

LEMMA 5.16. — *Let  $A \in \mathrm{Vect}_{\mathbb{F}}$ , with  $\#\mathbb{F} \geq 11$ , and  $\mu \in \mathcal{M}(A)$  be an  $\mathbb{F}^*$ -invariant map such that for every two-dimensional  $\mathbb{F}$ -subspace  $B \subset A$ ,  $\mu_B \in \Phi_{\mathbb{F}}(B)$ . Then  $\mu \in \Phi_{\mathbb{F}}(A)$ .*

*Proof.* — Assume the statement holds if  $\dim(A) \leq n - 1$ . Then  $\hat{\mu}$  is defined and, by Corollary 5.15, either  $\hat{\mu}$  is constant on  $\hat{\mathbb{P}}(A)$  or constant on the complement to one point.

If  $\hat{\mu}$  is constant, then the  $\mathbb{F}$ -linear envelope of points  $b \in A$  such that  $\mu(b) \neq \hat{\mu}$  has codimension at least two. Indeed, if there is a codimension one subspace  $B \subset A$  generated by such  $b$  then by assumption  $\mu \in \Phi_{\mathbb{F}}(B)$  and  $\mu^{\text{gen}}(B) \neq \hat{\mu}$ , contradicting the assumption that  $\hat{\mu}$  is constant. Otherwise, put  $A_1 := B$ . By induction,  $\mu \in \Phi_{\mathbb{F}}(B)$  and is constant on  $A \setminus B$ . Hence  $\mu \in \Phi_{\mathbb{F}}(A)$ .

Assume that  $\hat{\mu}$  is nonconstant and let  $B \subset A$  be the unique codimension one subspace with differing  $\mu^{\text{gen}}(B)$ . Choose an  $\mathbb{F}$ -basis  $b_1, \dots, b_{n-1}$  in  $B$  such that  $\mu(b_i) = \mu^{\text{gen}}(B)$ . Assume that there is a point  $a \in A \setminus B$  such that  $\mu(a) \neq$  the generic value of  $\hat{\mu}$  and let  $B'$  be the codimension one  $\mathbb{F}$ -subspace spanned  $b_1, \dots, b_{n-2}, a$ . Then  $\mu^{\text{gen}}(B') \neq$  the generic value of  $\hat{\mu}$ , contradicting the uniqueness of  $B$ . It follows that  $\mu$  is constant on  $A \setminus B$ .  $\square$

REMARK 5.17. — Let  $\mathbb{F}/\mathbb{F}'$  be a finite extension,  $A \in \text{Vect}_{\mathbb{F}}$ , considered as an  $\mathbb{F}'$ -vector space, and  $\mu \in \Phi_{\mathbb{F}'}(A)$ . If  $\mu$  is  $\mathbb{F}^*$ -invariant, then  $\mu \in \Phi_{\mathbb{F}}(A)$ . Indeed, by Lemma 5.8, the canonical  $\mathbb{F}'$ -flag is a flag of  $\mathbb{F}$ -vector spaces. We use this observation to reduce our problem to prime fields (resp.  $\mathbb{F}_4$ ).

lemm:444

LEMMA 5.18. — Let  $\mathbb{F}/\mathbb{F}'$  be a quadratic extension, with  $\#\mathbb{F}' > 2$ . Let  $A$  be an  $\mathbb{F}$ -vector space of dimension 2, considered as an  $\mathbb{F}'$ -vector space of dimension 4. Let  $\mu \in \mathcal{M}(A)$  be an  $\mathbb{F}^*$ -invariant map such that for every  $\mathbb{F}'$ -subspace  $C \subset A$ ,  $\dim_{\mathbb{F}'}(C) = 2$ , one has  $\mu_C \in \Phi_{\mathbb{F}'}(C)$ . Then  $\mu \in \Phi_{\mathbb{F}}(A)$ .

*Proof.* — First assume that  $\mu$  takes only two values on  $A \setminus \{0\}$ , say 0, 1, and that  $\mu \notin \Phi_{\mathbb{F}}(A)$ . Since  $\mathbb{P}_{\mathbb{F}}(A) = \mathbb{P}_{\mathbb{F}}^1$  there exist elements  $a_1, a_2, a_3, a_4 \in A \setminus \{0\}$  such that the orbits  $\mathbb{F}^* \cdot a_i$  do not intersect and

$$0 = \mu(a_1) = \mu(a_2) \neq \mu(a_3) = \mu(a_4) = 1.$$

Then  $\mathbb{F}^* \cdot a_i = \Lambda_i \setminus \{0\}$ , where  $\Lambda_i$  is a linear subspace over  $\mathbb{F}'$ . The  $\mathbb{F}'$ -span  $\Lambda_{12}$  of two nonzero vectors  $x_1 \in \Lambda_1, x_2 \in \Lambda_2$  has  $\mu^{\text{gen}}(\Lambda_{12}) = 0$ . Hence  $\Lambda_{12}$  contains at most one  $\mathbb{F}'$ -subspace  $\langle b \rangle$  of  $\mathbb{F}'$ -dimension 1 with generic value 1. The union of the spaces  $\Lambda_{12}$ , for different choices of  $x_1, x_2$ , covers  $A$  and

$$\#\{b \in A \mid \mu(b) = 1\} \leq (q + 1)^2,$$



where  $\#\mathbb{F}' = q$ . Similarly, there are at most  $(q+1)^2$  such nongeneric  $c \in A$  with  $\mu(c) = 0$ . Since  $\#\mathbb{P}^3(\mathbb{F}') = q^3 + q^2 + q + 1 > 2(q^2 + 2q + 1)$ , for  $q > 2$ , we get a contradiction.

Assume now that  $\mu$  takes at least 3 distinct values on  $A \setminus \{0\}$ , say 0, 1, 2, and that there are two vectors  $a_1, a_2 \in A$  such that the orbits  $\mathbb{F}^* \cdot a_1, \mathbb{F}^* \cdot a_2$  don't intersect and  $0 = \mu(a_1) = \mu(a_2)$ . Such a configuration must exist (take two  $\mathbb{F}'$ -spaces of  $\mathbb{F}'$ -dimension two spanned by  $\mathbb{F}^*$ -orbits; the  $\mathbb{F}'$  span of two generic vectors in these spaces contains elements whose  $\mu$ -value coincides with the value of  $\mu$  on one of the two orbits). The modified map, given by

$$\tilde{\mu}(a) := \begin{cases} 0 & \text{if } \mu(a) = 0 \\ 1 & \text{otherwise} \end{cases},$$

satisfies the conditions of the Lemma, and by the above argument  $\tilde{\mu} \in \Phi_{\mathbb{F}}(A)$ . In particular,  $\tilde{\mu} = 0$  outside one  $\mathbb{F}^*$ -orbit on  $A \setminus \{0\}$ . Since  $\mu$  is  $\mathbb{F}^*$ -invariant it follows that  $\mu$  takes two values, and not three as we assumed. Contradiction.  $\square$

lemm:p22

LEMMA 5.19. — *Let  $\mathbb{F}' = \mathbb{F}_p$  (resp.  $\mathbb{F}_4$ ), and  $\mathbb{F}/\mathbb{F}'$  be an extension of degree divisible by 4. Consider  $K = k(X)$  as an  $\mathbb{F}$ -vector space and let  $\mu, \mu' \in \mathcal{L}_{\mathbb{F}}(K)$  be a  $c$ -pair such that the linear span  $\sigma = \langle \mu, \mu', 1 \rangle_{\mathbb{Z}_\ell}$  does not contain an  $\Phi_{\mathbb{F}}$ -map. Then there exist an  $\mathbb{F}'$ -subspace  $B \subset K$  with  $\dim_{\mathbb{F}'}(B) = 3$ , two distinct  $\mathbb{F}'$ -subspaces  $C, C' \subset B$  of dimension 2 and maps  $\tilde{\mu}, \tilde{\mu}' \in \sigma$  such that*

- $\tilde{\mu}_C \notin \Phi_{\mathbb{F}'}(C)$  and  $\tilde{\mu}_{C'}$  is constant;
- $\tilde{\mu}'_{C'} \notin \Phi_{\mathbb{F}'}(C')$  and  $\tilde{\mu}'_C$  is constant;

*In particular, for every (nonzero) map  $\mu'' \in \sigma$  there exists an  $\mathbb{F}'$ -subspace  $C'' \subset B$ ,  $\dim_{\mathbb{F}'} C'' = 2$  with the property that  $\mu''_{C''} \notin \Phi_{\mathbb{F}'}(C'')$ .*

*Proof.* — We consider  $K$  as an  $\mathbb{F}$ -vector space as well as an  $\mathbb{F}'$ -vector space. Let  $\mu$  be an  $\mathbb{F}^*$ -invariant map on  $K$ . If  $\mu$  were an  $\mathbb{F}'$ -flag map on every two-dimensional  $\mathbb{F}'$ -subspace of  $K$  then, by Lemma 5.18,  $\mu$  would be an  $\mathbb{F}$ -flag map on every  $\mathbb{F}$ -subspace  $B \subset K$  of  $\dim_{\mathbb{F}} B = 2$ . Since  $\#\mathbb{F} \geq 11$  we could apply Lemma 5.16 and conclude that  $\mu \in \Phi_{\mathbb{F}}(K)$ .

Thus, since  $\mu \notin \Phi_{\mathbb{F}}(K)$ -map, there is an  $\mathbb{F}'$ -subspace  $C \subset K$ ,  $\dim_{\mathbb{F}'}(C) = 2$  such that  $\mu_C \notin \Phi_{\mathbb{F}'}(C)$ . If  $\mu'_C$  is constant, put  $\tilde{\mu}' := \mu$ . Otherwise, using the  $c$ -pair property on  $C$  we find constants  $d_C, d'_C, d''_C$ , with  $d'_C \neq 0$ , such that

$$d_C \mu + d'_C \mu'_C = d''_C \quad \text{and put } \tilde{\mu}' = \mu' - \frac{d''_C - d_C \mu}{d'_C}.$$

Then  $\tilde{\mu}'_C = 0$ . Since the linear combination  $\tilde{\mu}'$  is not a flag map, there exists a  $C'$ ,  $\dim_{\mathbb{F}'}(C') = 2$ , where  $\tilde{\mu}' \notin \Phi_{\mathbb{F}'}(C')$ . If  $\mu_{C'}$  is constant, put  $\tilde{\mu} := \mu$ . Otherwise, using the  $c$ -pair property on  $C'$  we find constants  $d_{C'}, d'_{C'}, d''_{C'}$ , with  $d'_{C'} \neq 0$ , such that

$$d_{C'}\mu + d'_{C'}\mu'_{C'} = d''_{C'} \quad \text{and put } \tilde{\mu} = \mu - \frac{d''_{C'} - d'_{C'}\tilde{\mu}'}{d_{C'}}.$$

Then  $\tilde{\mu}_{C'} = 0$  and  $\tilde{\mu}_C \notin \Phi_{\mathbb{F}'}(C)$  (since  $\tilde{\mu}'_C$  is constant). Now put

$$B := C + \frac{c}{c'} \cdot C',$$

for some nonzero  $c \in C$  and  $c' \in C'$ . Then  $\dim_{\mathbb{F}'}(B) = 3$ , the maps  $\tilde{\mu}_B, \tilde{\mu}'_B$  are linearly independent, and they satisfy the required conditions, by the logarithmic property. For  $s \neq 0$ , we have  $s\tilde{\mu} + s'\tilde{\mu}' \notin \Phi_{\mathbb{F}'}(C)$ . Otherwise,  $s\tilde{\mu} + s'\tilde{\mu}' \notin \Phi_{\mathbb{F}'}(\frac{c}{c'} \cdot C')$ .  $\square$

A detailed analysis of  $c$ -pairs on projective planes as above shows that such planes cannot exist. This will complete the proof of the main theorem.

emm:line-point0

LEMMA 5.20 (Lemma 4.3.2 in [3]). — *Let  $V \subset \mathbb{Z}_\ell^2$  be such that for any two pairs of distinct points the affine line through one pair and the affine line through the other have a common point and that this point of intersection is contained in  $V$ . Then  $V$  is contained in a line union one point.*

*Proof.* — Otherwise,  $V$  contains four points in general position. Embed  $\mathbb{Z}_\ell^2$  into  $\mathbb{P}^2(\mathbb{Q}_\ell)$ , choose coordinates for these four points

$$(1 : 0 : 0), (0 : 1 : 0), (0 : 0 : 1) \text{ and } (1 : 1 : 1)$$

and close  $V$  for the operation

$$x, y, z, t \mapsto \mathbb{I}(x, y) \cap \mathbb{I}(z, t), \quad \text{when } x \neq y, z \neq t, \mathbb{I}(x, y) \neq \mathbb{I}(z, t).$$

The closure  $\bar{V}$  of  $V$  satisfies the axioms of a projective plane (see Definition 4.1). For example, to verify that any “line” in  $\bar{V}$  contains at least three points it suffices to pick one of the four initial points not on this line and to draw lines through this point and the remaining three points in the initial set.

By the fundamental theorem of projective geometry,  $\bar{V} = \mathbb{P}^2(\mathbb{Q})$ . On the other hand,  $\mathbb{P}^2(\mathbb{Q})$  is dense in  $\mathbb{P}^2(\mathbb{Q}_\ell)$ . In particular, it cannot be contained in  $\mathbb{Z}_\ell^2$ . Contradiction.  $\square$

coro:line-point

COROLLARY 5.21. — *Let  $B = \mathbb{F}^3$  and  $\mu, \mu' \in \mathcal{M}(B)$  be a  $c$ -pair of  $\mathbb{F}^*$ -invariant maps. Then the image of  $\mathbb{P}(B)$  under map*

$$\begin{aligned} \varphi : \mathbb{P}(B) &\rightarrow \mathbb{A}^2(\mathbb{Z}_\ell) \\ b &\mapsto (\mu(b), \mu'(b)) \end{aligned}$$

*is contained in a union of an affine line and (possibly) one more point.*

*Proof.* — The  $c$ -pair condition for  $\mu, \mu'$  implies that the image of every  $\mathbb{P}^1 \subset \mathbb{P}(B)$  is contained in an affine line in  $\mathbb{Z}_\ell^2$ . Next, for any two pairs of distinct points  $(a, b), (a', b')$  in  $\varphi(\mathbb{P}(B))$  the affine lines  $\mathfrak{l} = \mathfrak{l}(a, b), \mathfrak{l}' = \mathfrak{l}'(a', b')$  in  $\mathbb{A}^2 = \mathbb{Z}_\ell^2$  through these pairs of points must intersect. (Choose  $\tilde{a}, \tilde{b}, \tilde{a}', \tilde{b}'$  in the preimages of  $a, b, a', b'$ ; the projective lines  $\tilde{\mathfrak{l}}, \tilde{\mathfrak{l}}' \subset \mathbb{P}(B) = \mathbb{P}^2$  through these points intersect in some  $x$  and, by the first observation,  $\varphi(x)$  must lie on both  $\mathfrak{l}$  and  $\mathfrak{l}'$ ). Now it suffices to apply Lemma 5.20.  $\square$

assu:proof

ASSUMPTION 5.22. — *We may now assume that*

- $\mathbb{F} = \mathbb{F}_p$  or  $\mathbb{F}_4$ ;
- $\mu, \mu' \in \mathcal{L}_{\mathbb{F}}(A)$  is a  $c$ -pair of linearly independent maps as in Lemma 5.19,
- $B$  is as in Lemma 5.19: for every two-dimensional  $C'' \subset B$  there exists a  $\mu'' \in \langle \mu, \mu' \rangle$  such that  $\mu''_{C''} \notin \Phi_{\mathbb{F}}(C'')$ .

*We can exclude the following degenerate cases, which contradict our assumption that no linear combination of  $\mu, \mu'$  is a flag map on  $B$ :*

- (1)  $\varphi(\mathbb{P}(B))$  is contained in a line; this means that  $\mu, \mu'$  are linearly dependent (modulo constants);
- (2)  $\varphi(\mathfrak{l})$  is a point, for some  $\mathfrak{l} \subset \mathbb{P}(B)$ ; this implies that  $\varphi(\mathfrak{l}) \in \varphi(\mathfrak{l}')$ , for all  $\mathfrak{l}' \subset \mathbb{P}(B)$  and  $\varphi(\mathbb{P}(B))$  is contained in a line, contradiction to (1);
- (3)  $\varphi$  is constant outside one line; here the affine map  $\mathbb{Z}_\ell^2 \rightarrow \mathbb{Z}_\ell$  projecting  $\varphi(\mathfrak{l})$  to one point gives a nontrivial flag map in the span of  $\mu, \mu'$ .

lemm:pi-x

LEMMA 5.23. — *Let  $\mathfrak{l}, \mathfrak{l}' \subset \mathbb{P}^2$  be distinct lines. Let  $x \in \mathbb{P}^2$  be a point such that  $\varphi(x) \notin (\varphi(\mathfrak{l}) \cup \varphi(\mathfrak{l}'))$ . Then there is a natural projective isomorphism  $\pi_{x, \mathfrak{l}'} : \mathfrak{l} \rightarrow \mathfrak{l}'$  respecting the level sets of  $\varphi$ . Namely, for every pair  $y_1, y_2 \in \mathfrak{l}$  with  $\varphi(y_1) = \varphi(y_2)$  one has*

$$\varphi(\pi_{x, \mathfrak{l}'}(y_1)) = \varphi(\pi_{x, \mathfrak{l}'}(y_2))$$

(and vice versa). In particular, if  $\varphi(\mathfrak{l}) \subset \varphi(\mathfrak{l}')$  then  $\varphi(\mathfrak{l}) = \varphi(\mathfrak{l}')$ .

*Proof.* — The images  $\varphi(\mathfrak{l}(x, y_1))$  and  $\varphi(\mathfrak{l}(x, y_2))$  span the same affine line  $L_x$ . We have  $\varphi(\mathfrak{l}') \not\subset L_x$ . Define  $\pi_{x, \mathfrak{l}'}(y_i) := \mathfrak{l}(x, y_i) \cap \mathfrak{l}'$ . By Corollary 5.21,  $\varphi(\pi_{x, \mathfrak{l}'}(y_i))$  are contained in the intersection of  $\varphi(\mathfrak{l}')$  and  $L_x$ , so that  $\varphi(\pi_{x, \mathfrak{l}'}(y_1)) = \varphi(\pi_{x, \mathfrak{l}'}(y_2))$ .  $\square$

coro:con4

COROLLARY 5.24. — *If there exist a line  $\mathfrak{l} \subset \mathbb{P}^2$  and a point  $x \in \mathfrak{l}$  such that  $\varphi$  is constant on  $\mathfrak{l} \setminus x$  then there is a nontrivial flag map in the span of  $\mu, \mu'$ .*

*Proof.* — By Assumption 5.22,  $\varphi$  is nonconstant on every line. Assume that there exists a point  $a \in \varphi(\mathbb{P}^2)$  such that  $\varphi^{-1}(a)$  consists exactly of  $x$ . Then for all  $\mathfrak{l}', \mathfrak{m}, \mathfrak{l}''$  not containing  $x$  one has  $\varphi(\mathfrak{l}') = \varphi(\mathfrak{l}'')$  and  $\varphi$  is constant on the complement to  $x$  on every line through  $x$ . Then a linear combination of  $\mu, \mu'$  is constant on  $\mathbb{P}^2 \setminus x$ , thus a flag map, contradicting the assumption.

Let  $x'$  be a point in  $\mathbb{P}^2 \setminus \mathfrak{l}$  with  $\varphi(x') = \varphi(x)$ . The lines  $\mathfrak{l}$  and  $\mathfrak{l}(x, x')$  are not equivalent,  $\varphi(\mathfrak{l}) \neq \varphi(\mathfrak{l}')$ . For any line  $\mathfrak{l}'' \neq \mathfrak{l}(x, x')$  through  $x'$  we have  $\varphi(\mathfrak{l} \cap \mathfrak{l}'') \neq \varphi(x)$ . Using a point on  $y \in \mathfrak{l}'$  with  $\varphi(y) \neq \varphi(x)$  and applying Lemma 5.23 we find that  $\varphi(\mathfrak{l}'') = \varphi(\mathfrak{l})$ . For any  $y \notin (\mathfrak{l} \cup \mathfrak{l}')$  consider the line  $\mathfrak{l}(x', y)$ . It follows that  $\varphi(y)$  equals the value of  $\varphi$  on  $\mathfrak{l} \setminus x$ , thus  $\varphi$  is constant on the complement to  $\mathfrak{l}'$ , contradicting Assumption 5.22(3).  $\square$

coro:same

COROLLARY 5.25. — *Let  $x, y \in \mathbb{P}_{\mathbb{F}}^2$  be distinct points so that  $\varphi(x), \varphi(y) \notin (\varphi(\mathfrak{l}) \cup \varphi(\mathfrak{l}'))$  and the line  $\mathfrak{l}(x, y)$  through  $x, y$  passes through the intersection  $q_0 := \mathfrak{l} \cap \mathfrak{l}'$ . Then the composition*

$$\pi_{x, \mathfrak{l}} \circ \pi_{y, \mathfrak{l}}^{-1} : \mathfrak{l} \rightarrow \mathfrak{l}$$

*induces a nontrivial translation on  $\mathfrak{l}$ , with fixed point  $q_0$ , preserving the level sets of  $\varphi$ . (By symmetry we have a similar translation on  $\mathfrak{l}'$ .)*

$$\pi_{x, \mathfrak{l}'} \circ \pi_{y, \mathfrak{l}'}^{-1} : \mathfrak{l} \rightarrow \mathfrak{l}$$

*In particular, if  $\mathbb{F} = \mathbb{F}_p$  (the prime field) then this translation is transitive on  $\mathfrak{l} \setminus (\mathfrak{l} \cap \mathfrak{l}')$  and  $\varphi$  is constant on this complement. If  $\mathbb{F} = \mathbb{F}_4$  then the complement to  $\mathfrak{l} \setminus (\mathfrak{l} \cap \mathfrak{l}')$  is a union of two (two point) orbits of this translation and  $\varphi$  is constant on each orbit.*

*Proof of Theorem 5.7.* — We keep the Assumptions 5.22.

For every point  $x \in \mathbb{P}^2$  and every line  $\mathfrak{l}$  through  $x$  there exist lines  $\mathfrak{l}', \mathfrak{l}''$  through  $x$  such that  $\varphi(\mathfrak{l}) = \varphi(\mathfrak{l}')$  and  $\varphi(\mathfrak{l}') \neq \varphi(\mathfrak{l}'')$ . Indeed, consider a line  $\tilde{\mathfrak{l}}$

with  $\varphi(x) \notin \varphi(\tilde{l})$ . If on all such lines  $\varphi$  takes more than two values, then all these lines are equivalent and  $\varphi$  is constant on the complement to  $x$  on every line through  $x$ , contradiction to Corollary 5.24. Otherwise, each value on  $\tilde{l}$  will be taken at least twice, hence the claim.

Corollary 5.25 gives a translation on  $l \setminus x$  preserving the level sets of  $\varphi$ . Over the prime field  $\mathbb{F}_p$ ,  $p > 2$ ,  $\varphi$  restricted to  $l$  is constant on the complement to  $x$  and we can apply Corollary 5.24.

Over  $\mathbb{F}_4$ ,  $\varphi$  is either constant on  $l \setminus x$ , contradicting Corollary 5.24, or the level sets of  $\varphi$  on  $l \setminus x$  fall into two orbits of cardinality two. Since we can pick  $x$  on  $l$  arbitrarily,  $\varphi$  must be constant on  $l$ , contradicting Assumption 5.22(2).  $\square$

## 6. Galois groups

sect:gal

Let  $k$  be an algebraic closure of a finite field of characteristic  $\neq \ell$ ,  $K$  the function field of an algebraic variety  $X$  over  $k$ ,  $\mathcal{G}_K^a$  the abelianization of the pro- $\ell$ -quotient  $\mathcal{G}_K$  of the Galois group  $G_K$  of a separable closure of  $K$ ,

$$\mathcal{G}_K^c = \mathcal{G}_K / [[\mathcal{G}_K, \mathcal{G}_K], \mathcal{G}_K] \xrightarrow{\text{pr}} \mathcal{G}_K^a$$

its canonical central extension and  $\text{pr}$  the natural projection.

defn:lift

DEFINITION 6.1. — We say that  $\gamma, \gamma' \in \mathcal{G}_K^a$  form a commuting pair if for some (and therefore any) of their preimages  $\tilde{\gamma}, \tilde{\gamma}'$  in  $\mathcal{G}_K^c$  one has  $[\tilde{\gamma}, \tilde{\gamma}'] = 0$ . A subgroup  $\mathcal{H}$  of  $\mathcal{G}^a$  is called liftable if any two elements in  $\mathcal{H}$  form a commuting pair.

defn:fan

DEFINITION 6.2. — The fan  $\Sigma_K = \{\sigma\}$  on  $\mathcal{G}_K^a$  is the set of all topologically noncyclic liftable subgroups  $\sigma \subset \mathcal{G}_K^a$  which are not properly contained in any other liftable subgroup of  $\mathcal{G}_K^a$ .

REMARK 6.3. — For function fields  $K/k$  of surfaces all groups  $\sigma \in \Sigma_K$  are isomorphic to torsion-free primitive  $\mathbb{Z}_\ell$ -submodules  $\sigma$  of rank 2, see Section 9.

nota:kkk

NOTATIONS 6.4. — Let

$$\mu_{\ell^n} := \{ \sqrt[n]{1} \}$$

and

$$\mathbb{Z}_\ell(1) = \lim_{n \rightarrow \infty} \mu_{\ell^n}.$$

We often identify  $\mathbb{Z}_\ell$  and  $\mathbb{Z}_\ell(1)$  (since  $k$  is algebraically closed). Write

$$\hat{K}^* := \lim_{n \rightarrow \infty} K^*/(K^*)^{\ell^n}$$

for the multiplicative group of (formal) rational functions on  $X$ .

thm:ga

**THEOREM 6.5 (Kummer theory).** — *The group  $K^*/k^*$  is a free  $\mathbb{Z}$ -module. One has*

- $K^*/(K^*)^{\ell^n} = (K^*/k^*)/\ell^n$ , for all  $n \in \mathbb{N}$ ;
- the discrete group  $K^*/(K^*)^{\ell^n}$  and the compact profinite group  $\mathcal{G}_K^a/\ell^n$  are Pontryagin dual to each other, for a  $\mu_{\ell^n}$ -duality;
- for  $K^*/k^* \xrightarrow{\sim} \mathbb{Z}^I$ , one has  $K^*/(K^*)^{\ell^n} \xrightarrow{\sim} (\mathbb{Z}/\ell^n)^I$  and

$$\mathcal{G}_K^a/\ell^n \xrightarrow{\sim} (\mathbb{Z}/\ell^n(1))^I,$$

hence the duality between  $\hat{K}^* = \widehat{K^*/k^*}$  and  $\mathcal{G}_K^a$  is modelled on that between

$$\{\text{functions } I \rightarrow \mathbb{Z}_\ell \text{ tending to } 0 \text{ at } \infty\} \text{ and } \mathbb{Z}_\ell^I.$$

rem:2

**LEMMA 6.6.** — *Let  $E/k$  be the function field of a curve. Then  $\Sigma_E = \emptyset$ .*

*Proof.* — By a result of Grothendieck, the pro- $\ell$  fundamental group  $(\pi_1)_\ell^\wedge$  of a curve punctured in finitely many points is free. We have

$$\mathcal{G}_E^a = \varprojlim_{J \subset I} \mathbb{Z}_\ell^J, \quad \mathcal{G}_E^c = \varprojlim_{J \subset I} \wedge^2(\mathbb{Z}_\ell^J),$$

with the commutation map equal to  $\wedge$ . This implies that a liftable subgroup of  $\mathcal{G}_E^a$  is topologically cyclic.  $\square$

## 7. Valuations

sect:val

In this section we recall basic results concerning valuations and valued fields (we follow [4]). Most of this material is an adaptation of well-known facts to our context.

nota:vg

**NOTATIONS 7.1.** — A *value group*, denoted by  $\Gamma$ , is a totally ordered (torsion-free) abelian group. We use the additive notation “+” for the group law and  $\geq$  for the order. We have

$$\Gamma = \Gamma^+ \cup \Gamma^-, \quad \Gamma^+ \cap \Gamma^- = \{0\} \quad \text{and} \quad \gamma \geq \gamma' \text{ iff } \gamma - \gamma' \in \Gamma^+.$$

Then  $\Gamma_\infty = \Gamma \cup \{\infty\}$  is a totally ordered monoid, by the conventions

$$\gamma < \infty, \gamma + \infty = \infty + \infty = \infty, \forall \gamma \in \Gamma.$$

**defn:valu**

**DEFINITION 7.2.** — A (nonarchimedean) valuation on a field  $K$  is a pair  $\nu = (\nu, \Gamma_\nu)$  consisting of a value group  $\Gamma_\nu$  and a map

$$\nu : K \rightarrow \Gamma_{\nu, \infty}$$

such that

- $\nu : K^* \rightarrow \Gamma_\nu$  is a surjective homomorphism;
- $\nu(\kappa + \kappa') \geq \min(\nu(\kappa), \nu(\kappa'))$  for all  $\kappa, \kappa' \in K$ ;
- $\nu(0) = \infty$ .

**rem:zeta**

**REMARK 7.3.** — In particular, since  $\Gamma_\nu$  is nontorsion,  $\nu(\zeta) = 0$  for every element  $\zeta$  of finite order in  $K^*$ .

A valuation is called *trivial* if  $\Gamma = \{0\}$ . If  $K = k(X)$  is a function field over an algebraic closure  $k$  of a finite field then every valuation of  $K$  restricts to a trivial valuation on  $k$  (every element in  $k^*$  is torsion).

**lemm:zll**

**LEMMA 7.4.** — Let  $K = k(X)$  and  $\nu$  be a nonarchimedean valuation on  $k(X)$ . Then  $\text{Hom}(\Gamma_\nu, \mathbb{Z}_\ell)$  is a finitely generated  $\mathbb{Z}_\ell$ -module.

*Proof.* — Note that the  $\mathbb{Q}$ -rank of  $\nu$  is bounded by  $\dim(X)$  (see [10]).  $\square$

**nota:ko**

**NOTATIONS 7.5.** — We denote by  $K_\nu$ ,  $\mathfrak{o}_\nu$ ,  $\mathfrak{m}_\nu$  and  $\mathbf{K}_\nu$  the completion of  $K$  with respect to  $\nu$ , the ring of  $\nu$ -integers in  $K$ , the maximal ideal of  $\mathfrak{o}_\nu$  and the residue field

$$\mathbf{K}_\nu := \mathfrak{o}_\nu / \mathfrak{m}_\nu.$$

If  $X$  (over  $k$ ) is a model for  $K$  then the *center*  $\mathfrak{c}(\nu)$  of a valuation is the irreducible subvariety defined by the prime ideal  $\mathfrak{m}_\nu \cap k[X]$  (provided  $\nu$  is nonnegative on  $k[X]$ ).

It is useful to keep in mind the following exact sequences:

**eqn:1**

$$(7.1) \quad 1 \rightarrow \mathfrak{o}_\nu^* \rightarrow K^* \rightarrow \Gamma_\nu \rightarrow 1$$

and

**eqn:2**

$$(7.2) \quad 1 \rightarrow (1 + \mathfrak{m}_\nu) \rightarrow \mathfrak{o}_\nu^* \rightarrow \mathbf{K}_\nu^* \rightarrow 1.$$

**nota:ine**

**NOTATIONS 7.6.** — Write  $\mathcal{I}_\nu^a \subset \mathcal{D}_\nu^a \subset \mathcal{G}_K^a$  for the images of the inertia and the decomposition group of the valuation  $\nu$  in  $\mathcal{G}_K^a$ .

`nota:iner`

NOTATIONS 7.7. — If  $\chi : \Gamma_\nu \rightarrow \mathbb{Z}_\ell(1)$  is a homomorphism then

$$\chi \circ \nu : K^* \rightarrow \mathbb{Z}_\ell(1)$$

defines an element of  $\mathcal{G}_K^a$ , called an inertia element of the valuation  $\nu$ . The group of such elements is  $\mathcal{I}_\nu^a \subset \mathcal{G}_K^a$ .

NOTATIONS 7.8. — The decomposition group  $\mathcal{D}_\nu^a$  is by definition equal to the image of  $\mathcal{G}_{K_\nu}^a$  in  $\mathcal{G}_K^a$ .

LEMMA 7.9. — *There is a natural embedding  $\mathcal{G}_{K_\nu}^a \hookrightarrow \mathcal{G}_K^a$  and a (canonical) isomorphism*

$$\mathcal{D}_\nu^a / \mathcal{I}_\nu^a \simeq \mathcal{G}_{K_\nu}^a.$$

*Proof.* — See Theorem 19.6 in [6], for example. □

`defn:div`

DEFINITION 7.10. — *Let  $K = k(X)$  be a function field. Its valuation  $\nu$  is*

- positive-dimensional if  $\text{tr deg}_k \mathbf{K}_\nu \geq 1$ ;
- divisorial if  $\text{tr deg}_k \mathbf{K}_\nu = \dim(X) - 1$ .

`nota:cv`

NOTATIONS 7.11. — We let  $\mathcal{V}_K$  be the set of all nontrivial (nonarchimedean) valuations of  $K$  and  $\mathcal{DV}_K$  the subset of divisorial valuations. If  $\nu \in \mathcal{DV}_K$  is realized by a divisor  $D$  on a model  $X$  of  $K$  (see Example 7.13) we sometimes write  $\mathcal{I}_D^a$ , resp.  $\mathcal{D}_D^a$ , for the corresponding inertia, resp. decomposition group.

`exam:curve`

EXAMPLE 7.12. — Let  $E = k(C)$  be the function field of a smooth curve. Every point  $q \in C(k)$  defines a nontrivial valuation  $\nu_q$  on  $E$  (the order of a function  $f \in E^*$  at  $q$ ). Conversely, every nontrivial valuation  $\nu$  on  $E$  defines a point  $q := \mathfrak{c}(\nu)$  on  $C$ .

`exam:suff`

EXAMPLE 7.13. — Let  $K = k(X)$  be the function field of a surface.

- Every positive-dimensional valuation is divisorial.
- Every (irreducible) curve  $D \subset X$  defines a valuation  $\nu_D$  on  $K$  with value group  $\mathbb{Z}$ . Conversely, every valuation on  $K$  with value group  $\mathbb{Z}$  and non-algebraically closed residue field defines a curve  $D$  on some model  $X$  of  $K$ .
- Every flag  $(D, q)$ , (curve, point on its normalization), defines a valuation  $\nu_{D,q}$  on  $K$  with value group  $\mathbb{Z}^2$ .
- There exist valuations on  $K$  with value group  $\mathbb{Q}$  and center supported in a point (on every model).



lemm:surf-valuation

LEMMA 7.14. — *Let  $K = k(X)$  be the function field of a surface. If  $\mathcal{D}_\nu^a/\mathcal{I}_\nu^a$  is nontrivial then  $\nu$  is divisorial.*

*Proof.* — The only 1-dimensional valuations on function fields of surfaces are divisorial valuations. For other valuations, the residue field  $K_\nu = k$  is algebraically closed and  $\mathcal{G}_{K_\nu}^a$  trivial.  $\square$

## 8. A dictionary

sect:dict

Write

$$\begin{aligned}\mathcal{L}_K &:= \mathcal{L}_k(K) = \{ \text{homomorphisms } K^* \rightarrow \mathbb{Z}_\ell(1) \} \\ \Phi_K &:= \Phi_k(K) = \{ \text{flag maps } K \rightarrow \mathbb{Z}_\ell(1) \}\end{aligned}$$

prop:decomp

PROPOSITION 8.1. — *One has the following identifications:*

$$\begin{aligned}\mathcal{G}_K^a &= \mathcal{L}_K, \\ \mathcal{D}_\nu^a &= \{ \mu \in \mathcal{L}_K \mid \mu \text{ trivial on } (1 + \mathfrak{m}_\nu) \}, \\ \mathcal{I}_\nu^a &= \{ \mu \in \mathcal{L}_K \mid \mu \text{ trivial on } \mathfrak{o}_\nu^* \}.\end{aligned}$$

*If two nonproportional  $\mu, \mu' \in \mathcal{G}_K^a$  form a commuting pair then the corresponding maps  $\mu, \mu' \in \mathcal{L}_K$  form a  $c$ -pair (in the sense of Definition 5.6).*

*Proof.* — The first identification is a consequence of Kummer theory 6.5. The second identification can be checked on one-dimensional subfields of  $K$ , where it is evident. For this and the third identification we use (7.1) and (7.2). For the last statement, assume that  $\mu, \mu' \in \mathcal{L}_K$  don't form a  $c$ -pair. Then there is an  $x \in K$  such that the restrictions of  $\mu, \mu' \in \mathcal{L}_K$  to the subgroup  $\langle 1, x \rangle$  are linearly independent. Therefore,  $\mu, \mu' \in \mathcal{G}_K^a$  define a rank 2 liftable subgroup in  $\mathcal{G}_{k(x)}^a$ . Such subgroups don't exist since  $\mathcal{G}_{k(x)}$  is a free pro- $\ell$ -group.  $\square$

exam:cp

EXAMPLE 8.2. — If  $\mu \in \mathcal{D}_\nu^a$  and  $\alpha \in \mathcal{I}_\nu^a$  then  $\mu, \alpha$  form a commuting pair.

prop:may

PROPOSITION 8.3. — *Let  $K$  be a field and  $\alpha \in \Phi_K \cap \mathcal{L}_K$ . Then there exists a unique valuation  $\nu = (\nu_\alpha, \Gamma_{\nu_\alpha})$  (up to equivalence) and a homomorphism  $\text{pr} : \Gamma_{\nu_\alpha} \rightarrow \mathbb{Z}_\ell(1)$  such that*

$$\alpha(f) = \text{pr}(\nu_\alpha(f))$$

*for all  $f \in K^*$ . In particular,  $\alpha \in \mathcal{I}_\nu^a$  (under the identification of Proposition 8.1).*

*Proof.* — Let  $\mathbb{F}$  be a finite subfield of  $k$  and assume that  $\alpha(f) \neq \alpha(f')$  for some  $f, f' \in K$  and consider the line  $\mathbb{P}^1 = \mathbb{P}(\mathbb{F}f + \mathbb{F}f')$ . Since  $\alpha$  is a flag map, it is constant outside one point on this  $\mathbb{P}^1$  so that either  $\alpha(f + f') = \alpha(f)$  or  $\alpha(f + f') = \alpha(f')$ . This defines a relation:  $f' >_\alpha f$  (in the first case) and  $f >_\alpha f'$  (otherwise). If  $\alpha(f) = \alpha(f')$  and there exists an  $f''$  such that  $\alpha(f) \neq \alpha(f'')$  and  $f >_\alpha f'' >_\alpha f'$  then we put  $f >_\alpha f'$ . Otherwise, we put  $f =_\alpha f'$ .

It was proved in [3], Section 2.4, that the above definitions are correct and that  $>_\alpha$  is indeed an order which defines a filtration on the additive group  $K$  by subgroups  $(K_\gamma)_{\gamma \in \Gamma}$  such that

- $K = \bigcup_{\gamma \in \Gamma} K_\gamma$  and
- $\bigcap_{\gamma \in \Gamma} K_\gamma = \emptyset$ ,

where  $\Gamma$  is the set of equivalence classes with respect to  $=_\alpha$ . Since  $\alpha \in \mathcal{L}_K$  this order is compatible with multiplication in  $K^*$ , so that the map  $K \rightarrow \Gamma$  is a valuation and  $\alpha$  factors as  $K^* \rightarrow \Gamma \rightarrow \mathbb{Z}_\ell \simeq \mathbb{Z}_\ell(1)$ . By (7.1),  $\alpha \in \mathcal{I}_\nu^a$ .  $\square$

coro:lift

**COROLLARY 8.4.** — *Every (topologically) noncyclic liftable subgroup of  $\mathcal{G}_K^a$  contains an inertia element of some valuation.*

*Proof.* — By Theorem 5.7, every such liftable subgroup contains an  $\Phi$ -map, which by Proposition 8.3 belongs to some inertia group.  $\square$

## 9. Flag maps and valuations

sect:af-val

In this section we give a Galois-theoretic description of inertia and decomposition subgroups of divisorial valuations.

lemm:am

**LEMMA 9.1.** — *Let  $\alpha \in \Phi_K \cap \mathcal{L}_K$ ,  $\nu = \nu_\alpha$  be the associated valuation and  $\mu \in \mathcal{L}_K$ . Assume that  $\alpha, \mu$  form a c-pair. Then*

$$\mu(1 + \mathfrak{m}_\nu) = \mu(1).$$

*In particular, the restriction of  $\mu$  to  $\mathfrak{o}_\nu$  is induced from  $\mathbf{K}_\nu$ .*

*Proof.* — We have

- (1)  $\alpha(\kappa) = 0$  for all  $\kappa \in \mathfrak{o}_\nu \setminus \mathfrak{m}_\nu$ ;
- (2)  $\alpha(\kappa + m) = \alpha(\kappa)$  for all  $\kappa$  and  $m$  as above;
- (3)  $\mathfrak{m}_\nu$  is generated by  $m \in \mathfrak{o}_\nu$  such that  $\alpha(m) \neq 0$ .

If  $m \in \mathfrak{m}_\nu$  is such that  $\alpha(m) \neq 0$  and  $\kappa \in \mathfrak{o}_\nu \setminus \mathfrak{m}_\nu$  then  $\alpha$  is nonconstant on the subgroup  $A := \langle \kappa, m \rangle$ . Then

$$\mu(\kappa + m) = \mu(\kappa).$$

Indeed, if  $\mu$  is nonconstant on  $A$  the restriction  $\mu_A$  is proportional to  $\alpha_A$  (by the  $c$ -pair property) and  $\alpha$  satisfies (2). In particular, for such  $m$  we have  $\mu(1 + m) = \mu(1)$ .

If  $\alpha(m) = 0$  then there exists  $m', m'' \in \mathfrak{m}_\nu$  such that  $m = m' + m''$  and  $\alpha(m') = \alpha(m'') \neq 0$ . Indeed, there exists an  $m' \in \mathfrak{m}_\nu$  such that  $m > m' > 1$  and  $\alpha(m') \neq \alpha(1) = 0$ . Since  $\alpha$  takes only two values on the subgroup  $\langle m', m \rangle \subset \mathfrak{m}_\nu$  we have

$$\alpha(m'') = \alpha(-m' + m) = \alpha(m').$$

Therefore,

$$0 = \mu(1 + m') + \mu(1 + m'') = \mu(1 + m + m'm'').$$

Put  $\kappa = 1 + m + m'm''$  and observe that  $\alpha(-m'm'') = 2\alpha(m') \neq 0$ . By the argument above

$$\mu(\kappa - m'm'') = \mu(\kappa) = \mu(1 + m' + m'') = \mu(1 + m),$$

as claimed.  $\square$

coro:divv

**COROLLARY 9.2.** — *Inertia elements  $\alpha \in \mathcal{I}_\nu^a$  commute only with elements  $\mu \in \mathcal{D}_\nu^a$ .*

prop:delta

**PROPOSITION 9.3.** — *Let  $K = k(X)$  be the function field of a surface. Every  $\sigma \in \Sigma_K$  has  $\text{rk}_{\mathbb{Z}_\ell} \sigma = 2$ . Moreover, it defines a unique valuation  $\nu = \nu_\sigma$  of  $K$  so that either every element of  $\sigma$  is inertial for  $\nu$ , or  $\nu$  is divisorial and there is an element  $\mu \in \sigma$  which is not inertial for  $\nu$ , but  $\mu \in \mathcal{D}_\nu^a$ .*

*If distinct  $\sigma, \sigma' \in \Sigma_K$  have a nonzero intersection then there exists a divisorial valuation  $\nu''$  such that*

- $\sigma, \sigma' \in \mathcal{D}_{\nu''}^a$ ;
- $\sigma \cap \sigma' = \mathcal{I}_{\nu''}^a$ .

*Conversely, if  $\sigma \in \Sigma_K$  is not contained in a  $\mathcal{D}_{\nu''}^a$  for any divisorial valuation  $\nu''$  then for all  $\sigma' \in \Sigma_K$ ,  $\sigma' \neq \sigma$ , one has  $\sigma \cap \sigma' = 0$ .*

*Proof.* — We saw that  $\sigma \in \Sigma_K$  contains an inertia element  $\alpha$  for some valuation  $\nu$ . Since  $\sigma$  is topologically noncyclic there is a  $\mu \in \sigma$ ,  $\mathbb{Z}_\ell$ -independent on  $\alpha$ , and commuting with  $\alpha$ . If  $\mu$  is not inertial, that is,  $\mu \notin \Phi_K$ , then  $\nu$  gives

a nontrivial element in the (abelianized) Galois group of the residue field  $K_\nu$  of  $\nu$ . Thus  $\nu$  is divisorial,  $K_\nu$  is 1-dimensional and every liftable subgroup in  $\mathcal{G}_{K_\nu}^a$  has  $\mathbb{Z}_\ell$ -rank equal to one. Hence  $\text{rk}_{\mathbb{Z}_\ell} \sigma = 2$  in this case and, by Corollary 9.2,  $\mu \in \mathcal{D}_\nu^a$ . Such a valuation  $\nu$  is unique, since  $\mathcal{I}_\nu^a \cap \mathcal{I}_{\nu'}^a = 0$  for distinct divisorial  $\nu, \nu'$ .

If  $\sigma$  contains *only* inertia elements, then there exists a unique valuation  $\nu$  such that  $\sigma \in \mathcal{I}_\nu^a$ . Indeed, either  $\mathfrak{m}_\nu + \mathfrak{m}_{\nu'} = K$  or we may assume that  $\mathfrak{m}_\nu \subset \mathfrak{m}_{\nu'}$  (and  $\mathfrak{o}_\nu \supset \mathfrak{o}_{\nu'}$ ). The first case is impossible since the corresponding inertia groups don't intersect. In the second case,  $\mathcal{I}_\nu^a \subset \mathcal{I}_{\nu'}^a$ , as claimed. Moreover, it follows that  $\text{rk}_{\mathbb{Z}_\ell} \sigma = 2$ , since the  $\mathbb{Q}$ -rank of any valuation on a surface (over  $\bar{\mathbb{F}}_q$ ) is at most two. This gives  $\nu = \nu_\sigma$  in this case.

If distinct  $\sigma, \sigma'$  have a nontrivial intersection, then the subgroup  $\mathcal{D} \subset \mathcal{G}_K^a$  generated by  $\sigma, \sigma'$  is not the inertia group of any valuation (the rank of those is  $\leq 2$ , as we have seen above). If the  $\sigma \cap \sigma'$  contains a nontrivial inertia element  $\alpha$  then  $\mathcal{D}$  is contained in the decomposition group of this element (all elements of  $\mathcal{D}$  commute with  $\alpha$ ) and the corresponding valuation is divisorial. If  $\mu \in \sigma \cap \sigma'$  is not an inertia element then there exist inertia elements  $\alpha \in \sigma$  and  $\alpha' \in \sigma'$  corresponding to distinct *divisorial* valuations  $\nu, \nu'$ . The decomposition groups of distinct divisorial valuations don't intersect.  $\square$

Proposition 9.3 allows us to identify intrinsically (in terms of the Galois group) inertia subgroups of divisorial valuations as well as their decomposition groups as follows. Every pair of distinct groups  $\sigma, \sigma' \in \Sigma_K$  with a nontrivial intersection defines a divisorial valuation  $\nu$ , whose inertia group

$$\mathcal{I}_\nu^a = \sigma \cap \sigma'.$$

The corresponding decomposition subgroup is

$$\mathcal{D}_\nu^a = \cup_{\sigma \supset \mathcal{I}_\nu^a} \sigma.$$

## 10. Galois groups of curves

sect:valc

Here we give a Galois-theoretic characterization of subgroups  $\sigma \in \Sigma_K$  which are inertia subgroups of rank two valuations of  $K$  arising from a flag  $(C, q)$ , where  $D$  is a smooth irreducible curve (on some model of  $K$ ) and  $q \in C(k)$  is a point (see Example 7.13). We show that Galois-theoretic data determine the genus of  $C$  and all “points” on  $C$ , as special liftable subgroups of rank two inside  $\mathcal{G}_{k(C)}^a$ .

Throughout,  $E = k(D)$  is the function field of a smooth curve of genus  $g$ . We have an exact sequence

$$0 \rightarrow E^*/k^* \rightarrow \text{Div}(C) \rightarrow \text{Pic}(C) \rightarrow 0$$

(where  $\text{Div}(C)$  can be identified with the free abelian group generated by points in  $C(k)$ ). This gives a dual sequence

$$(10.1) \quad 0 \rightarrow \mathbb{Z}_\ell \xrightarrow{\Delta} \mathcal{M}(C(k), \mathbb{Z}_\ell) \rightarrow \mathcal{G}_E^a \rightarrow \mathbb{Z}_\ell^{2g} \rightarrow 0,$$

with the identifications

- $\text{Hom}(\text{Pic}(C), \mathbb{Z}_\ell) = \Delta(\mathbb{Z}_\ell)$  (since  $\text{Pic}^0(C)$  is torsion);
- $\mathcal{M}(C(k), \mathbb{Z}_\ell) = \text{Hom}(\text{Div}(C), \mathbb{Z}_\ell)$  is the  $\mathbb{Z}_\ell$ -linear space of maps from  $C(k) \rightarrow \mathbb{Z}_\ell$ ;
- $\mathbb{Z}_\ell^{2g} = \text{Ext}^1(\text{Pic}^0(C), \mathbb{Z}_\ell)$ .

Using this model and the results in Section 6, we can interpret

$$(10.2) \quad \mathcal{G}_E^a \subset \mathcal{M}(C(k), \mathbb{Q}_\ell) / \text{constant maps}$$

as the  $\mathbb{Z}_\ell$ -linear subspace of all maps  $\mu : C(k) \rightarrow \mathbb{Q}_\ell$  (modulo constant maps) such that

$$[\mu, f] \in \mathbb{Z}_\ell \text{ for all } f \in E^*/k^*.$$

Here  $[\cdot, \cdot]$  is the pairing:

$$(10.3) \quad \begin{aligned} \mathcal{M}(C(k), \mathbb{Q}_\ell) \times E^*/k^* &\rightarrow \mathbb{Q}_\ell \\ (\mu, f) &\mapsto [\mu, f] := \sum_q \mu(q) f_q, \end{aligned}$$

where  $\text{div}(f) = \sum_q f_q q$ . In detail, let  $\gamma \in \mathcal{G}_E^a$  be an element of the Galois group. By Kummer theory,  $\gamma$  is a homomorphism  $K^*/k^* \rightarrow \mathbb{Z}_\ell(1) \simeq \mathbb{Z}_\ell$ . Choose a point  $c_0 \in C(k)$ . For every point  $c \in C(k)$ , there is an  $m_c \in \mathbb{N}$  such that the divisor  $m_c(c - c_0)$  is principal. Define a map

$$\begin{aligned} \mu_\gamma : C(k) &\rightarrow \mathbb{Q}_\ell, \\ c &\mapsto \gamma(m_c(c - c_0))/m_c. \end{aligned}$$

Changing  $c_0$  we get maps differing by a constant map.

In this interpretation, an element of an inertia subgroup  $\mathcal{I}_w^a \subset \mathcal{G}_E^a$  corresponds to a “delta”-map (constant outside the point  $q_w$ ). Each  $\mathcal{I}_w^a$  has a canonical (topological) generator  $\delta_w$ , given by  $\delta_w(f) = \nu_w(f)$ , for all  $f \in E^*/k^*$ .

The (diagonal) map  $\Delta \in \mathcal{M}(C(k), \mathbb{Q}_\ell)$  from (10.1) is then given by

$$\Delta = \sum_{w \in \mathcal{V}_E} \delta_w = \sum_{q_w \in C(k)} \delta_{q_w}.$$

defn:sup

DEFINITION 10.1. — We say that the support of a subgroup  $\mathcal{I} \subset \mathcal{G}_E^a$  is  $\leq s$  and write

$$|\text{supp}(\mathcal{I})| \leq s$$

if there exist valuations  $w_1, \dots, w_s \in \mathcal{V}_E$  such that

$$\mathcal{I} \subset \langle \mathcal{I}_{w_1}^a, \dots, \mathcal{I}_{w_s}^a \rangle_{\mathbb{Z}_\ell} \subset \mathcal{G}_E^a.$$

Otherwise, we write  $|\text{supp}(\mathcal{I})| > s$ .

lemm:cu

LEMMA 10.2. — Let  $\mathcal{I} \subset \mathcal{G}_E^a$  be a topologically cyclic subgroup such that  $|\text{supp}(\mathcal{I})| > s \geq 2$ . Then there exist a finite set  $\{f_j\}_{j \in J} \subset E^*$  and an  $m \in \mathbb{N}$  such that the map

$$\begin{aligned} \psi : \mathcal{G}_E^a &\rightarrow V := \bigoplus_{j \in J} \mathbb{Z}/\ell^m \\ \mu &\mapsto ([\mu, f_j]_m)_{j \in J} \end{aligned}$$

has the following property: for every set  $\{w_1, \dots, w_s\} \subset \mathcal{V}_E$

$$\psi(\mathcal{I}) \not\subset \langle \psi(\mathcal{I}_{w_1}^a), \dots, \psi(\mathcal{I}_{w_s}^a) \rangle_{\mathbb{Z}_\ell}.$$

*Proof.* — Let  $\iota \in \mathcal{G}_E^a \subset \mathcal{M}(C(k), \mathbb{Q}_\ell)$  be a representative, as in (10.2), of a topological generator of  $\mathcal{I}$ , where  $\text{supp}(\mathcal{I}) > s$ . There are three possibilities:

- (1)  $\iota(C(k)) \subset \mathbb{Q}_\ell$  is infinite;
- (2) there is a  $b \in \iota(C(k)) \subset \mathbb{Q}_\ell$  such that  $\iota^{-1}(b)$  is infinite and there exist at least  $s+1$  distinct points  $q_{s+2}, \dots, q_{2s+2} \in C(k)$  such that  $\iota(q_j) \neq b$  for all  $j = s+2, \dots, 2s+2$ ;
- (3) otherwise:  $\iota(C(k))$  is finite, there is a  $b$  with  $\iota^{-1}(b)$  infinite and there are at most  $s$  distinct points with values differing from  $b$ .

In Case (3),  $|\text{supp}(\mathcal{I})| \leq s$ .

In Case (1), choose any set  $Q = \{q_1, \dots, q_{2s+2}\} \subset C(k)$  of points with pairwise distinct values. In Case (2) choose distinct  $q_1, \dots, q_{s+1} \in \iota^{-1}(b)$  and put  $Q := \{q_1, \dots, q_{2s+2}\}$ . In both cases, if  $Q' \subset Q$  is any subset of cardinality  $|Q'| = s$  then  $\iota$  is nonconstant on  $Q \setminus Q'$ . In particular, there exist points  $q_{s_1}, q_{s_2} \in Q \setminus Q'$  such that

eqn:iota

$$(10.4) \quad \iota(q_{s_1}) \neq \iota(q_{s_2}).$$

We may assume that  $\iota(Q) \subset \mathbb{Z}_\ell$  (replacing  $\iota$  by a sufficiently high multiple, if necessary). Now we choose an  $m'' \in \mathbb{N}$  such that all values of  $\iota$  on  $Q$  remain pairwise distinct modulo  $\mathbb{Z}/\ell^{m''}$ . Let  $\text{Div}_Q^0(C)$  be the abelian group of degree zero divisors on  $C$  supported in  $Q$ . By Lemma 3.2, there is an  $n = n_Q \in \mathbb{N}$  such that  $nD$  is principal for every  $D \in \text{Div}_Q^0(C)$ . In particular, for every  $q_{s_1}, q_{s_2} \in Q$  there is a function  $f \in E^*$  such that  $\text{div}(f) = n(q_{s_1} - q_{s_2})$ . Write  $n = \ell^{m'} \bar{n}$ , with  $\gcd(\bar{n}, \ell) = 1$ , and put  $m = m' + m''$ .

We have a pairing (Kummer theory)

$$\begin{aligned} \mathcal{G}_E^a \times n\text{Div}_Q^0(C) &\rightarrow \mathbb{Z}/\ell^m \\ (\mu, f) &\mapsto [\mu, f]_m. \end{aligned}$$

Notice that  $[\mathcal{I}_w^a, f] = 0$  for all  $w$  with  $q_w \notin Q$  and all  $f \in E^*$  supported in  $Q$ . Further, for every  $Q' \subset Q$  with  $|Q'| = s$  and points  $q_{s_1}, q_{s_2} \in Q \setminus Q'$  as in (10.4) there is an  $f \in E^*$  with divisor  $\text{div}(f) = n(q_{s_1} - q_{s_2})$  such that

$$[\iota, f] = n \cdot (\iota(q_{s_1}) - \iota(q_{s_2})) \neq 0 \pmod{\ell^m}$$

and

$$[\mathcal{I}_{w'}^a, f] = 0$$

for all  $\mathcal{I}_{w'}^a$  of  $q' \in Q'$ . Let  $\{f_j\}_{j \in J}$  be a basis for  $\ell^m \cdot \text{Div}_Q^0(C)$ , with  $f_j \in E^*$ . The map

$$\begin{aligned} \psi : \mathcal{G}_E^a &\rightarrow \bigoplus_{j \in J} \mathbb{Z}/\ell^m \\ \mu &\mapsto ([\mu, f_j]_m)_{j \in J} \end{aligned}$$

satisfies the required properties.  $\square$

The next step is an *intrinsic* definition of inertia subgroups

$$\mathcal{I}_w^a \subset \mathcal{D}_\nu^a / \mathcal{I}_\nu^a = \mathcal{G}_{k(C)}^a.$$

We have a projection

$$\pi_\nu : \mathcal{G}_K^a \rightarrow \mathcal{G}_K^a / \mathcal{I}_\nu^a$$

and an inclusion

$$\mathcal{G}_{K_\nu}^a = \mathcal{D}_\nu^a / \mathcal{I}_\nu^a \hookrightarrow \mathcal{G}_K^a / \mathcal{I}_\nu^a$$

prop:intri

PROPOSITION 10.3. — *Let  $\nu$  be a divisorial valuation of  $K$ . A subgroup*

$$\mathcal{I} \subset \mathcal{D}_\nu^a / \mathcal{I}_\nu^a$$

*is the inertia subgroup of a divisorial valuation of  $k(C) = K_\nu$  iff for every homomorphism*

$$\psi : \mathcal{G}_K^a / \mathcal{I}_\nu^a \rightarrow V$$

onto a finite abelian group  $V$  there exists a divisorial valuation  $\nu_\psi$  such that

$$\psi(\mathcal{I}) = \psi \circ \pi_\nu(\mathcal{I}_{\nu_\psi}^a).$$

*Proof.* — Let  $C$  be the smooth model for  $\mathbf{K}_\nu = k(C)$ ,

$$\mathcal{I} = \mathcal{I}_w^a \subset \mathcal{D}_\nu^a / \mathcal{I}_\nu^a$$

the inertia subgroup of a divisorial valuation of  $k(C)$  corresponding to a point  $q = q_w \in C(k)$  and

$$\psi : \mathcal{G}_K^a / \mathcal{I}_\nu^a \rightarrow V$$

a homomorphism onto a finite abelian group. Since  $\mathcal{G}_K^a$  is a pro- $\ell$ -group, we may assume that

$$V = \bigoplus_{j \in J} \mathbb{Z} / \ell^{n_j},$$

for some  $n_j \in \mathbb{N}$ . Let  $n = \max_j(n_j)$ . By Kummer theory,

$$\mathrm{Hom}(\mathcal{G}_K^a, \mathbb{Z} / \ell^n) = K^* / (K^*)^{\ell^n}$$

so that  $\psi$  determines elements

$$\bar{f}_j \in K^* / (K^*)^{\ell^n}$$

(for all  $j \in J$ ). Choose functions  $f_j$  projecting to  $\bar{f}_j$ . They define a finite set of divisors  $D_{ij}$  on  $X$ . Moreover,  $f_j$  are not simultaneously constant on  $C$  (otherwise,  $\psi(\mathcal{G}_{k(C)}^a) = \psi(\mathcal{I}_{k(C)}^a)$ ). Changing the model  $\tilde{X} \rightarrow X$ , if necessary, we may assume that

- $C$  is smooth (and irreducible);
- there exists exactly one irreducible component  $D$  in the full preimage of  $\bigcup D_{ij}$  which intersects  $C$  in  $q$ . Moreover, this intersection is transversal (see Section 3). Then the image of  $\mathcal{I}_D^a$  under  $\psi$  is equal to the image of  $\mathcal{I}_w^a$ .

Conversely, we need to show that if  $\mathcal{I} \neq \mathcal{I}_w^a$  (for some  $w \in \mathcal{DV}_{\mathbf{K}_\nu}$ ), then there exists a homomorphism

$$\psi : \mathcal{G}_K^a / \mathcal{I}_\nu^a \rightarrow V$$

onto a finite abelian group  $V$  such that for all  $\nu' \in \mathcal{DV}_K$  one has

$$\psi(\mathcal{I}) \neq \psi \circ \pi_{\nu'}(\mathcal{I}_{\nu'}^a).$$

We consider two cases

- (1) there exist two points  $q, q' \in C(k)$  such that  $\mathcal{I} \subset \langle \mathcal{I}_w^a, \mathcal{I}_{w'}^a \rangle$ ;
- (2) otherwise.



*Case 1.* There exists a rational map  $\pi : X \rightarrow \mathbb{P}^1$  such that its restriction

$$\pi : C \rightarrow \mathbb{P}^1$$

is surjective, unramified at  $q, q'$  and  $\pi(q) \neq \pi(q')$ . Under the induced map of Galois groups

$$\pi_*(\mathcal{I}) \subset \langle \mathcal{I}_{\pi(w)}^a, \mathcal{I}_{\pi(w')}^a \rangle$$

but is not contained in either  $\mathcal{I}_{\pi(q)}^a$  or  $\mathcal{I}_{\pi(q')}^a$ . Thus there exist a finite abelian group  $V$  and a map  $\psi : \mathcal{G}_{k(\mathbb{P}^1)}^a \rightarrow V$  such that  $\psi(\mathcal{I}) \notin \psi(\mathcal{I}_{w''}^a)$  for any  $q'' \in \mathbb{P}^1$ . It follows that

$$\psi \circ \pi_*(\mathcal{I}) \notin \psi \circ \pi_*(\mathcal{I}_\nu^a)$$

for any  $\nu \in \mathcal{DV}_K$ .

*Case 2.* By Lemma 10.2, there exist a finite set of functions  $\bar{f}_j \in k(C)$ , with support in a finite set  $Q = \{q_0, \dots, q_s\} \subset C(k)$ , and an  $m \in \mathbb{N}$  such that the homomorphism

$$\begin{aligned} \bar{\psi} : \mathcal{G}_{k(C)}^a &\rightarrow V = \bigoplus_{j \in J} \mathbb{Z}/\ell^m \\ \mu &\mapsto ([\mu, \bar{f}_j]_m)_{j \in J} \end{aligned}$$

has the property that for all  $w, w' \in \mathcal{DV}_{k(C)}$

$$\psi(\mathcal{I}) \notin \langle \psi(\mathcal{I}_w^a), \psi(\mathcal{I}_{w'}^a) \rangle_{\mathbb{Z}_\ell}.$$

Next we choose a model for  $X$  and  $C$  as in Lemma 3.8. In particular, there exist functions  $g_j$  with divisor

$$\text{div}(g_j) = n \cdot (D_j - D_0) + (H_j - H'_j)$$

such that all the divisors are irreducible, with transversal intersections and  $\text{div}(g_j)|_C = n(q_j - q_0)$ . These functions  $g_j$  define a homomorphism

$$\psi : \mathcal{G}_K^a / \mathcal{I}_\nu^a \rightarrow V.$$

If  $D$  is a divisor on  $X$  then  $\psi \circ \pi_\nu(\mathcal{I}_D^a) = 0$  unless  $D = D_j$  for some  $j$ . In this case  $\psi \circ \pi_\nu(\mathcal{I}_{D_j}^a) = \psi(\mathcal{I}_{w_j}^a)$ .

Let  $\nu' \in \mathcal{DV}_K$  and  $\mathfrak{c}(\nu') \subset X$  be its center on  $X$ . There are three cases:

- $\mathfrak{c}(\nu') \not\subset D_j$  for any  $j$ : then  $\psi \circ \pi_{\nu'}(\mathcal{I}_{\nu'}^a) = 0$ ;
- $\mathfrak{c}(\nu') \in D_j^0$ , where  $D_j^0 = D_j \setminus (\cup_{j' \neq j} D_j \cap D_{j'})$ : then

$$\psi \circ \pi_{\nu'}(\mathcal{I}_{\nu'}^a) \subset \psi(\mathcal{I}_{w_j}^a);$$

–  $\mathfrak{c}(\nu') \in D_j \cap D_{j'}$  for some  $j, j'$ : then

$$\psi \circ \pi_\nu(\mathcal{I}_{\nu'}^a) \subset \langle \psi(\mathcal{I}_{w_j}^a), \psi(\mathcal{I}_{w_{j'}}^a) \rangle_{\mathbb{Z}_\ell}.$$

All three possibilities contradict our assumptions.  $\square$

lemm:gen-type

LEMMA 10.4. — *Let  $E = k(C)$  be the function field of a curve. Then  $g(C) \geq 1$  iff there exists a non-zero homomorphism from  $\mathcal{G}_E^a$  to a finite (abelian) group which maps all inertia elements to 0.*

*Proof.* — Indeed, every curve of genus  $\geq 1$  over a finite field of characteristic  $p$  has unramified coverings of degree  $\ell$ . These coverings define maps of Galois groups, which are trivial on all inertia elements. If  $C$  is rational then  $\mathcal{G}_E^a$ , and hence its image under every homomorphism (onto any finite group), is generated by inertia elements (see the exact sequence (10.1)).  $\square$

rem:gen-tt

REMARK 10.5. — Combining this with Proposition 10.3 we can decide in purely Galois-theoretic terms which divisorial valuations of  $K$  correspond to nonrational (irreducible) curves  $C$  on some model  $X$  of  $K$ . We call such valuations *nonrational*.

sect:recc

## 11. Valuations on surfaces

Next we are lead to the following problem: How to characterize subgroups  $\widehat{k(C)^*} \subset \hat{K}^*$ ? We recall a geometric argument (from algebraic K-theory) characterizing pairs  $f, g \in K^*$  which are contained in  $k(C)^* \subset K^*$ , for some curve  $C$  (such curves correspond to projections  $X \rightarrow C$ ).

Let  $\nu$  be a divisorial valuation of  $K$  and

$$\nu : K^* \rightarrow \mathbb{Z}$$

the valuation map. We have the residue map

$$\text{res}_\nu : \text{Ker}(\nu) \rightarrow \mathbf{K}_\nu^*$$

and a bilinear (with respect to multiplication) symbol

eqn:ress

$$(11.1) \quad \begin{array}{ccc} K^* \times K^* & \xrightarrow{\varrho_\nu} & \mathbf{K}_\nu^* \\ f, g & \mapsto & f^{\nu(g)} / g^{\nu(f)}. \end{array}$$

On a smooth model  $X$  of  $K$ , where  $\nu = \nu_D$  for a divisor  $D \subset X$ , we define

eqn:res-1

$$(11.2) \quad \varrho_\nu = \varrho_D : K^* \times K^* \rightarrow \mathbf{K}_\nu^*$$

as follows:

- $\varrho_\nu(f, g) = 1$  if both  $f, g$  are invertible on  $D$ ;
- $\varrho_\nu(f, g) = f_D^m$  if  $f$  is invertible ( $f_D$  is the restriction to  $D$ ) and  $g$  has multiplicity  $m$  along  $D$ ;
- $\varrho_\nu(f, g) = (f^{m_g}/g^{m_f})_D$  in the general case, when  $f, g$  have multiplicities  $m_f, m_g$ , respectively.

The definition does not depend on the choice of the model.

The following is a standard result in K-theory. We include a proof since we will need its  $\ell$ -adic version.

lemm:fg

LEMMA 11.1. — For  $f, g \in K^*$

$$\varrho_\nu(f, g) = 1 \quad \forall \nu \in \mathcal{DV}_K \iff f, g \in E = k(C) \subset K \text{ for some curve } C.$$

*Proof.* — ( $\Leftarrow$ ) On an appropriate model  $X$  we have  $\nu = \nu_D$  for a divisor  $D \subset X$  and  $\pi : X \rightarrow C$  is regular and flat with irreducible generic fiber (and  $f, g \in k(C)^*$ ). By definition,  $\varrho_\nu(f, g) = 1$  if  $D$  is not in a fiber of  $\pi$ . If  $D$  is in a fiber then there is a  $t \in k(C)^*$ ,  $\nu_D(t) \neq 0$  such that both  $ft^{m_f}, gt^{m_g}$  are regular and constant on  $D$  (for some  $m_f, m_g \in \mathbb{N}$ ) so that  $\varrho_\nu(f, g) = 1$ .

( $\Rightarrow$ ) Assume that  $\varrho_\nu(f, g) = 1$  for every  $\nu \in \mathcal{DV}_K$ . Every nonconstant function  $f$  defines a unique map (with irreducible generic fiber)

$$\pi_f : X \rightarrow C_f$$

which corresponds to the algebraic closure of  $k(f)$  in  $K$  (we will say that  $f$  is induced from  $C_f$ ). We claim that  $\pi_f = \pi_g$ .

Since  $f$  is induced from  $C_f$ , we have

$$\operatorname{div}(f) = \sum_{q \in Q} a_q D_q,$$

where  $Q \subset C_f(k)$  is finite and  $D_q = \pi^{-1}(q)$ . Then  $D_q^2 = 0$  and  $D_q$  is either a multiple of a fiber of  $\pi_g$  or it has an irreducible component  $D \subset D_q$  which dominates  $C_g$  (under  $\pi_g$ ). In the second case, the restriction of  $g$  to  $D_q$  is a nonconstant element in  $k(D_q)$ . Then  $\nu_D(f) \neq 0$ , while  $\nu_D(g) = 0$ . Hence  $\varrho_D(f, g) \neq 1$  since it coincides with  $g_D^{-\nu_D(f)} \neq 1$ , a contradiction. Therefore,

all  $D_q$  are contained in the finitely many fibers  $S$  of  $\pi_g$ . That means  $\text{div}(f)$  does not intersect the fibers  $R_t, t \in C_g, t \notin S$  which implies that  $f$  is constant on such  $R_t$ . Hence  $f$  belongs to the normal closure of  $k(C_g)$  in  $K$ , and in fact  $f \in k(C_g)$  since  $k(C_g)$  is algebraically closed in  $K$ , by construction. Thus  $f$  is induced from  $C_g$  and hence  $C_f = C_g$  and  $\pi_f = \pi_g$ .  $\square$

## 12. $\ell$ -adic analysis: generalities

sect:ella

Let  $X$  be a smooth model of  $K$ . An element  $f \in K^*/k^*$  gives rise to a divisor  $D = D_f$  on  $X$  and conversely, such a  $D$  uniquely determines  $f$ . Recall that the Galois group  $\mathcal{G}_K^a$  determines  $\hat{K}^*$ , a group substantially bigger than  $K^*/k^*$ . Our goal is to characterize the  $\ell$ -adic space  $K^*/k^* \otimes \mathbb{Z}_\ell \subset \hat{K}^*$ . The Galois datum  $(\mathcal{G}_K^a, \Sigma_K)$  allows us to distinguish between rational and nonrational irreducible divisors on  $X$  (via the corresponding valuations) and to describe intrinsically a subspace  $\mathcal{FS}(K) \subset \hat{K}^*$  (of divisors with finite nonrational support, see 12.6 and 12.7). In this section we further shrink  $\mathcal{FS}(K)$ , giving an intrinsic characterization of those elements which have finite divisorial support on every smooth model  $X$ .

For any smooth algebraic variety  $X$  over  $k$  with function field  $K = k(X)$  we have an exact sequence

eqn:seqq (12.1) 
$$0 \rightarrow K^*/k^* \xrightarrow{\rho_X} \text{Div}(X) \xrightarrow{\varphi} \text{Pic}(X) \rightarrow 0,$$

where  $\text{Div}(X)$  is the group of (Weil or Cartier) divisors of  $X$ . Write

$$\text{Div}(X)_\ell := \text{Div}(X) \otimes \mathbb{Z}_\ell \quad \text{and} \quad \text{Pic}(X)_\ell := \text{Pic}(X) \otimes \mathbb{Z}_\ell$$

for the group of *finite*  $\ell$ -adic divisors, resp.  $\ell$ -adic Picard group. We have an exact sequence:

eq:exx (12.2) 
$$0 \rightarrow K^* \otimes \mathbb{Z}_\ell \xrightarrow{\rho_{X,\ell}} \text{Div}(X)_\ell \xrightarrow{\varphi_\ell} \text{Pic}(X)_\ell \rightarrow 0$$

Let

$$\widehat{\text{Div}}(X) := \{D = \sum_{m \in M} \hat{a}_m D_m\}, \quad \text{resp.} \quad \widehat{\text{Div}}_{\text{nr}}(X) \subset \widehat{\text{Div}}(X),$$

be the group of divisors (resp. nonrational divisors) with *decreasing coefficients*:

- $M$  is a countable set;
- for all  $r \in \mathbb{Z}$  the set

$$\{m \mid |\hat{a}_m|_\ell \leq r\}$$

is finite;

- for  $D \in \widehat{\text{Div}}_{\text{nr}}(X)$ , all  $D_m$  are nonrational.

Clearly,  $\text{Div}(X)_\ell \subset \widehat{\text{Div}}(X)$ . Every element

$$\hat{f} \in \hat{K}^* = \lim_{n \rightarrow \infty} K^*/(K^*)^{\ell^n}$$

has a representation

$$\hat{f} = (f_n)_{n \in \mathbb{N}} \text{ or } f = f_0 f_1^\ell f_2^{\ell^2} \cdots,$$

with  $f_n \in K^*$ . We have homomorphisms

$$\begin{aligned} \hat{\rho}_X : \hat{K}^* &\rightarrow \widehat{\text{Div}}(X), \\ \hat{f} &\mapsto \text{div}(\hat{f}) := \sum_{n \in \mathbb{N}} \ell^n \cdot \text{div}(f_n) = \sum_m \hat{a}_m D_m, \end{aligned}$$

$$\hat{\rho}_{X, \text{nr}} : \hat{K}^* \rightarrow \widehat{\text{Div}}(X) \xrightarrow{\text{pr}} \widehat{\text{Div}}_{\text{nr}}(X),$$

where  $D_m \subset X$  are irreducible divisors,

$$\hat{a}_m = \sum_{n \in \mathbb{N}} a_{nm} \ell^n \in \mathbb{Z}_\ell,$$

with  $a_{nm} \in \mathbb{Z}$ , and

$$\text{div}(f_n) = \sum_m a_{nm} D_m.$$

Here  $\text{div}(f_n)$  is the Cartier divisor of  $f_n$  and  $\sum_m a_{nm} D_m$  is its image in the group of Weil divisors. The map  $\hat{\rho}_X$  has a kernel

$$T_\ell(X) := \varprojlim \text{Tor}_1(\mathbb{Z}/\ell^n, \text{Pic}(X)[\ell]),$$

where  $\text{Pic}(X)[\ell] \subset \text{Pic}(X)$  is the  $\ell$ -power torsion subgroup. In particular  $T_\ell(X) = \mathbb{Z}_\ell^{2g}$ , where  $g$  is the dimension of  $\text{Pic}^0(X)$ .

lemm:t-ell

LEMMA 12.1. — *For varieties over  $k$  we have*

- (1) *a morphism  $X \rightarrow Y$  induces a homomorphism  $T_\ell(X) \rightarrow T_\ell(Y)$ ;*
- (2) *the canonical morphism  $X \hookrightarrow \text{Alb}(X)$  into the Albanese variety induces a canonical isomorphism  $T_\ell(\text{Alb}(X)) \rightarrow T_\ell(X)$ ;*
- (3) *if  $X' \rightarrow X$  is a birational isomorphism between smooth varieties then  $T_\ell(X) \rightarrow T_\ell(X')$  is an isomorphism;*

(4) *an exact sequence of abelian varieties*

$$1 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 1$$

*induces an exact sequence*

$$1 \rightarrow T_\ell(A'') \rightarrow T_\ell(A) \rightarrow T_\ell(A').$$

*Proof.* — The follows from the corresponding properties of the functor  $\text{Pic}^0$  for smooth algebraic varieties over  $k$ .  $\square$

We have a diagram

eqn:dia-need

(12.3)

$$\begin{array}{ccccccccccc} 0 & \rightarrow & K^* \otimes \mathbb{Z}_\ell & \xrightarrow{\rho_{X,\ell}} & \text{Div}(X)_\ell & \xrightarrow{\varphi_\ell} & \text{Pic}(X)_\ell & \rightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \rightarrow & T_\ell(X) & \rightarrow & \hat{K}^* & \xrightarrow{\hat{\rho}_X} & \widehat{\text{Div}}(X) & \xrightarrow{\hat{\varphi}_\ell} & \widehat{\text{Pic}}(X) & \rightarrow & 0, \end{array}$$

where

$$\widehat{\text{Pic}}(X) := \varprojlim \text{Pic}(X) \otimes \mathbb{Z}/\ell^n = \text{NS}(X) \otimes \mathbb{Z}_\ell.$$

Every  $\nu \in \mathcal{DV}_K$  gives rise to a homomorphism

$$\hat{\nu} : \hat{K}^* \rightarrow \mathbb{Z}_\ell$$

and a homomorphism

$$\text{r}\hat{\text{e}}\text{s}_\nu : \text{Ker}(\hat{\nu}) \rightarrow \hat{K}_\nu^*$$

and a symbol

$$\hat{\varrho}_\nu : \hat{K}^* \times \hat{K}^* \rightarrow \hat{K}_\nu^*.$$

On a smooth model  $X$ , where  $\nu = \nu_D$  for a divisor  $D \subset X$ ,  $\hat{\nu}(\hat{f})$  is the  $\ell$ -adic coefficient at  $D$  of  $\text{div}(\hat{f})$ , while  $\hat{\varrho}_\nu$  is the natural  $\mathbb{Z}_\ell$ -bilinear generalization of (11.1).

lemm:t-e112

LEMMA 12.2. — *Let  $X$  be a smooth surface or a smooth curve over  $k$  and  $K = k(X)$ . Then*

$$T_\ell(X) = \cap_{\nu \in \mathcal{DV}_K} \text{Ker}(\hat{\nu}).$$

*Proof.* — Follows from Lemma 12.1.  $\square$

In particular, we have the map  $\text{r}\hat{\text{e}}\text{s}_\nu : T_\ell(K) := T_\ell(X) \rightarrow \hat{K}_\nu^*$ .

lemm:tell

LEMMA 12.3. — For all  $\nu \in \mathcal{DV}_K$  we have

$$\text{r\acute{e}s}_\nu(T_\ell(K)) \subset T_\ell(\mathbf{K}_\nu).$$

*Proof.* — Let  $X$  be a model of  $K$  such that  $\nu = \nu_D$ , where  $D$  is a smooth curve. We may assume (after blowing up) that  $X$  contains a divisor  $D'$  intersecting  $D$  in exactly one point. Consider the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Ker}(\nu) & \longrightarrow & \text{Div}(X \setminus D) & \longrightarrow & \text{Pic}(X) \longrightarrow 0 \\ & & \downarrow \text{res}_\nu & & \downarrow & & \downarrow \delta \\ 0 & \longrightarrow & \mathbf{K}_\nu^* & \longrightarrow & \text{Div}(D) & \longrightarrow & \text{Pic}(D) \longrightarrow 0 \end{array}$$

where  $\text{Div}(X \setminus D)$  is the  $\mathbb{Z}$ -module spanned by divisors different from  $D$ . By the choice of  $X$ , the restriction  $\delta$  induces a surjection  $\text{NS}(X) \rightarrow \text{NS}(D)$ .

Tensoring all  $\mathbb{Z}$ -modules with  $\mathbb{Z}/\ell^n$  and passing to the projective limit we obtain a map

$$T_\ell(K) \rightarrow T_\ell(\mathbf{K}_\nu),$$

and the claim.  $\square$

By Lemma 3.21, if  $T_\ell(K) \neq 0$  then either  $X$  contains only finitely many rational curves, or  $X$ , modulo purely inseparable covers, is a rational pencil over a curve  $C$  of genus  $g(C) \geq 1$ .

lemm:DD

LEMMA 12.4. — Let  $X/k$  be a smooth projective surface,  $M$  a finite set and

$$D = \sum_{m \in M} a_m D_m \in \text{Div}(X)_\ell, \quad a_m \in \mathbb{Z}_\ell$$

a divisor such that  $\varphi_\ell(D) = 0$ . Then there exist a finite set  $I$ , functions  $f_i \in K^*$  and numbers  $a_i \in \mathbb{Z}_\ell$ , linearly independent over  $\mathbb{Z}$ , such that for all  $i \in I$

$$\text{supp}_X(f_i) \subset \text{supp}_X(D)$$

and

$$D = \sum a_i \text{div}(f_i).$$

*Proof.* — It suffices to consider equation (12.2) and to observe that any  $\mathbb{Z}_\ell$ -lattice of principal divisors with support in a finite set of divisors contains a generating  $\mathbb{Z}$ -lattice of principal divisors.  $\square$

**defi:commu**

DEFINITION 12.5. — We say that  $\hat{f}, \hat{g} \in \hat{K}^*$  commute if  $\hat{\rho}_\nu(\hat{f}, \hat{g}) = 1$ , for all divisorial  $\nu$ . We say that they have disjoint support if for all divisorial valuations  $\nu \in \mathcal{DV}_K$

$$\hat{\nu}(\hat{f}) \cdot \hat{\nu}(\hat{g}) = 0.$$

We say that  $\hat{f} \in \hat{K}^*$  has nontrivial commutators if there exist  $\hat{g} \in \hat{K}^*$  with disjoint support (from  $\hat{f}$ ) which commute with  $\hat{f}$ .

**nota:supp**

NOTATIONS 12.6. — We put

$$\begin{aligned} \text{supp}_K(\hat{f}) &:= \{ \nu \in \mathcal{DV}_K \mid \hat{f} \text{ nontrivial on } \mathcal{I}_\nu^a \}; \\ \text{supp}_X(\hat{f}) &:= \{ D_m \subset X \mid \hat{a}_m \neq 0 \}. \end{aligned}$$

**rem:alt**

DEFINITION 12.7. — We say that  $\hat{f}$  has finite nonrational support if the set of nonrational  $\nu \in \text{supp}_K(\hat{f})$  is finite (see Lemma 10.4 for the definition and Galois-theoretic characterization of nonrational valuations). Let

$$\mathcal{FS}(K) \subset \hat{K}^*$$

be the subgroup of such elements.

Note that for  $\hat{f} \in \mathcal{FS}(K)$ , its nonrational component  $\hat{\rho}_{X,\text{nr}}(\hat{f})$  is independent of the model  $X$ . More precisely, for any birational morphism  $X' \rightarrow X$  we can identify  $\widehat{\text{Div}}_{\text{nr}}(X') = \widehat{\text{Div}}_{\text{nr}}(X)$ . Under this identification

$$\rho_{X',\text{nr}}(\hat{f}) = \rho_{X,\text{nr}}(\hat{f}).$$

**defn:fs**

DEFINITION 12.8. — We say that  $\hat{f}$  has finite support on the model  $X$  if  $\text{supp}_X(\hat{f})$  is finite. Put

$$\mathcal{FS}_X(K) := \{ \hat{f} \in \hat{K}^* \mid \rho_X(\hat{f}) \in \text{Div}(X)_\ell \}.$$

**lemm:indep**

LEMMA 12.9. — The definition of  $\mathcal{FS}_X(K)$  does not depend on the choice of a smooth model  $X$ .

*Proof.* — For any two smooth models  $X', X''$  we can find a smooth model  $X$  dominating both. The difference between the sets of irreducible divisors  $\text{Div}(X')$ , resp.  $\text{Div}(X'')$ , and  $\text{Div}(X)$  is finite and consists of rational curves.  $\square$



Equation (12.3) implies the exact sequence

$$\text{eqn:diag} \quad (12.4) \quad 0 \rightarrow T_\ell(X) \rightarrow \mathcal{FS}_X(K) \rightarrow K^*/k^* \otimes \mathbb{Z}_\ell \rightarrow 0.$$

Indeed, if  $\hat{f} \in \mathcal{FS}_X(K)$  then  $\text{div}(\hat{f}) \in \text{Div}(X)_\ell$  and its image in  $\text{Pic}(X)_\ell$  is zero. Thus there is an element  $f \in K^*/k^* \otimes \mathbb{Z}_\ell$  with the same  $\ell$ -adic divisor. By definition  $\hat{f}/f \in T_\ell(X)$ .

We proceed to give a Galois-theoretic characterization of  $\mathcal{FS}_X(K)$ .

Case I. Let  $K$  be the function field of a surface  $X$  containing only finitely many rational curves. Then

$$\mathcal{FS}(K) = \mathcal{FS}_X(K).$$

Case II. Assume that, after a purely inseparable extension,  $X$  admits a fibration over a curve of genus  $\geq 1$ , with generic fiber a rational curve.

Let  $\mathcal{FS}'(K) \subset \hat{K}^*$  be the group generated by all  $\hat{f}$  such that

- $\hat{f}$  has nontrivial nonrational support;
- $\hat{f}$  has nontrivial commutators.

Then, for every model  $X$  of  $K$ , we have

$$\mathcal{FS}'(K) = \mathcal{FS}_X(K).$$

Indeed, an infinite rational tail in  $\hat{f}$  in this case consists of an infinite number of fibers. Same holds for  $\hat{g}$ . Thus the divisor of  $\hat{f}$  (resp.  $\hat{g}$ ) intersects all but finitely many fibers in the infinite rational tail of  $\hat{g}$  (resp.  $\hat{f}$ ) with intersection multiplicity some power of  $p = \text{char}(k)$ . Consider  $\nu$  corresponding to rational curves in the divisor of  $\hat{g}$  intersecting the divisor of  $\hat{f}$  as above. Then

$$\hat{\rho}_\nu(\hat{f}, \hat{g}) \neq 1,$$

contradiction.

Case III. By Lemma 3.21, we can now assume that  $\text{Pic}^0(X) = 0$ .

Let  $\mathcal{F}_X(K)$  be the set of all  $f \in K(X)^*/k^*$  such that

- (1)  $\rho_{X, \text{nr}}(f) \neq 0$  and

- (2) for every rational curve  $D \subset X$  with  $\nu = \nu_D$  either  $D \in \text{supp}_X(f)$  or  $\text{res}_\nu(f) \not\equiv 0 \pmod{\ell}$  in  $K_\nu^*/k^*$ .

Geometrically, condition (2) means that if  $D$  is not a component of the divisor of  $f$  then there is a point in  $D \cap \text{div}(f)$  whose multiplicity is prime to  $\ell$ .

lemm:imp

LEMMA 12.10. — *The set  $\mathcal{F}_X(K)$  generates  $K^*/k^*$ .*

*Proof.* — Let  $y \in K^*$  be a function such that the generic fiber of the corresponding rational map  $\pi_y : X \rightarrow \mathbb{P}_y^1$ , is an irreducible nonrational curve. Such  $y$  generate  $K^*$ .

Let  $\tilde{X}$  be a blowup of  $X$  such that the induced map  $\tilde{\pi}_y : \tilde{X} \rightarrow \mathbb{P}_y^1$  is regular. Let  $D \subset \tilde{X}$  be a rational curve surjecting onto  $\mathbb{P}_y^1$ . For generic coprime polynomials  $P, Q \in k[y]$  of degree  $\geq 2$ , choose 4 points  $q_1, \dots, q_4$  in the preimage of  $\phi$ , where

$$\begin{aligned} \phi : \mathbb{P}_y^1 &\rightarrow \mathbb{P}^1 \\ y &\mapsto f(y) := P(y)/Q(y). \end{aligned}$$

If  $f$  were nonconstant on  $D \subset \tilde{X}$  and  $\text{res}_\nu(f)$  were  $0 \pmod{\ell}$  then the local ramification indices of  $f$  and hence of  $y$  were divisible by  $\ell$ . Thus we would have a map  $\pi_y : D \rightarrow \mathbb{P}_y^1$  with all local ramification indices over 4 points divisible by  $\ell$ , and by Hurwitz' theorem,  $g(D) > 0$ , contradicting the rationality of  $D$ . It follows that  $f \in \mathcal{F}_X(K)$ . Clearly, such elements generate  $k(y)^*$ , and thus  $K^*$ .  $\square$

lemm:imp2

LEMMA 12.11. — *For every pair of nonzero commuting elements  $\hat{f}, \hat{g} \in \mathcal{FS}(K)$  with disjoint support such that there exists an  $f \in \mathcal{F}_X(K)$  with*

$$f = \hat{f} \pmod{\ell}, \text{ in } \hat{K}^*$$

*one has  $\hat{f} \in \mathcal{FS}_X(K)$  and  $\hat{g} \in \mathcal{FS}_X(K)$ .*

*Proof.* — Write

$$\begin{aligned} \rho_X(\hat{f}) &= \sum_{i \in I} n_i D_i + \ell \sum_{j=1}^{\infty} n_j C_j, \\ \rho_X(\hat{g}) &= \sum_{i \in I'} n'_i D'_i + \ell \sum_{j=1}^{\infty} n'_j C'_j, \end{aligned}$$

where  $I, I'$  are finite sets and the second sum is an infinite series over distinct rational curves  $C_j, C'_j \subset X$ . By assumption, the sets  $\{D_i\}_{i \in I}$ ,  $\{C_j\}_{j \in \mathbb{N}}$ ,  $\{D'_i\}_{i \in I'}$ ,  $\{C'_j\}_{j \in \mathbb{N}}$  are disjoint.

By assumption,  $\hat{\rho}_\nu(\hat{f}, \hat{g}) = 1$ , for all  $\nu$ . For  $\nu = \nu_D$ , where  $D \in \text{supp}_X(\hat{g})$ , this symbol equals the residue of  $\hat{f}$  on  $D$ , which equals the corresponding residue of  $f \bmod \ell$ . For rational curves in the support of  $\hat{g}$  it is nonzero by (2). It follows that every divisor in  $\text{supp}_X(\hat{g})$  is nonrational and that  $\hat{g} \in \mathcal{FS}_X(K)$ .

Since  $T_\ell(X) = 0$ , we can write  $\text{div}(\hat{g}) = \sum_{m \in M} a_m D_m$ , where  $M$  is a finite set, all  $D_m \subset X$  are nonrational divisors and  $a_m \in \mathbb{Z}_\ell$ , or  $\hat{g} = \prod_{i \in I''} g_i^{b_i}$ , with  $I''$  a finite set,  $g_i \in K^*/k^*$  and  $b_i \in \mathbb{Z}_\ell$ . We may assume that all divisors in  $\text{supp}_X(g_1)$  are nonrational.

The restriction of  $g_1$  to every irreducible component of the divisor of  $\hat{f}$  is identically zero. This means that under the map

$$\pi_{g_1} : X \rightarrow C$$

all components of  $\text{supp}_X(\hat{f})$  map to points (note that  $C = \mathbb{P}^1$ , since  $\text{Pic}^0(X) = 0$ ). Since the components of the divisor of  $g_1$  are nonrational, the generic fiber of  $\pi_{g_1}$  is also nonrational. Thus  $\text{supp}_X(\hat{f})$  contains only a finite number of rational divisors, so that  $\hat{f} \in \mathcal{FS}_X(K)$ .  $\square$

lemm:new-arg

LEMMA 12.12. — *Let  $x, y \in \mathcal{FS}_X(K)$  be noncommuting elements. Assume that the three elements  $x, y, xy$  have nontrivial commutators in  $\mathcal{FS}_X(K)$ . Then there exists a unique  $a \in \mathbb{Z}_\ell/\mathbb{Z}_\ell^*$  such that*

$$x, y, xy \in a \cdot K^*/k^* \subset \mathcal{FS}_X(K)$$

*Proof.* — Let  $P_x, P_y$  and  $P_{xy}$  be minimal  $\mathbb{Z}$ -sublattices of  $\mathcal{FS}_X(K)$  such that  $x \in P_x \otimes \mathbb{Z}_\ell, y \in P_y \otimes \mathbb{Z}_\ell$  and  $xy \in P_{xy} \otimes \mathbb{Z}_\ell$ . Note that  $P_x \cap P_y = 0$ , since there are no  $\mathbb{Z}$ -relations - the elements of  $P_x$ , resp.  $P_y$ , belong to the same pencil and nontrivial elements of different pencils are distinct. The lattice  $P_{xy}$  surjects onto both  $P_x$  and  $P_y$ . If one of the projections had a kernel, there would be an element in  $P_x$  belonging to the pencil  $P_{xy}$ , and similarly for  $P_y$ , but there are no common elements. We can now assume that one of the projections is an isomorphism. The corresponding section in  $P_{xy}$  gives the claimed imbedding.  $\square$

coro:now

COROLLARY 12.13. — *Let  $\mathcal{K} \subset \mathcal{FS}(K)$  be a lattice such that*

- *every element in  $\mathcal{K}$  has a nontrivial commutator and*

- $\mathcal{K}$  surjects onto  $K^*/\ell$ .

Then there is  $a \in \mathbb{Z}_\ell^*$  such that  $a \cdot \mathcal{K} \subset K^* \otimes \mathbb{Z}_{(\ell)}$ .

*Proof.* — Since  $\mathcal{K}$  is generated by the preimages of  $\mathcal{F}_X \subset K^*/\ell$  the lattice  $\mathcal{K}$  is contained in  $\mathcal{FS}_X(K)$ . Then we apply Lemma 12.12.  $\square$

In particular, any such lattice is contained in  $a^{-1} \cdot K^* \otimes \mathbb{Z}_{(\ell)}$  with the same property. Thus there is a unique maximal lattice for any triple of elements  $x, y, xy$  with nontrivial commutators so that  $\rho_\nu(x, y) \not\equiv 0 \pmod{\ell}$  for some  $\nu \in \mathcal{DV}_K$ . It equals  $a \cdot K^* \otimes \mathbb{Z}_{(\ell)}$ , for some  $a \in \mathbb{Z}_\ell^*$ .

### 13. $\ell$ -adic analysis: curves

sect:curves-ell

In this section we begin the process recognition of the lattice  $K^*/k^* \subset \hat{K}^*$ . We solve an analogous problem for the function field of a rational curve. This result will play an essential role in the analysis of surfaces.

prop:cc

PROPOSITION 13.1. — *Let  $\tilde{k}$  be the closure of a finite field,  $\text{char}(\tilde{k}) \neq \ell$ ,  $C$  a curve over  $\tilde{k}$  of genus  $g$  with function field  $E = \tilde{k}(C)$  and*

$$\Psi : \mathcal{G}_{k(\mathbb{P}^1)}^a \rightarrow \mathcal{G}_E^a$$

*an isomorphism of Galois groups inducing an isomorphism on inertia groups of divisorial valuations, that is, a bijection on the set of such groups and isomorphisms of corresponding groups. Let*

$$\Psi^* : \widehat{k(\mathbb{P}^1)^*} \rightarrow \hat{E}^*$$

*be the dual isomorphism. Then  $E = \tilde{k}(\mathbb{P}^1)$  and there is a constant  $a \in \mathbb{Z}_\ell^*$  such that  $\Psi^*(k(\mathbb{P}^1)^*/k^*) = a \cdot E^*/\tilde{k}^*$ .*

*Proof.* — Recalling the exact sequence (10.1), we have a commuting diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}_\ell(\Delta_{C(\tilde{k})}) & \longrightarrow & \mathcal{M}(C(\tilde{k})) & \longrightarrow & \mathcal{G}_E^a \longrightarrow \mathbb{Z}_\ell^{2g} \longrightarrow 0 \\ & & & & & & \downarrow \\ 0 & \longrightarrow & \mathbb{Z}_\ell(\Delta_{\mathbb{P}^1(k)}) & \longrightarrow & \mathcal{M}(\mathbb{P}^1(k)) & \longrightarrow & \mathcal{G}_{k(\mathbb{P}^1)}^a \longrightarrow 0 \end{array}$$

Since  $\Psi$  is an isomorphism on inertia groups  $\mathcal{I}_w^a$ , for each  $w$ , the sets  $C(\tilde{k})$  and  $\mathbb{P}^1(k)$  coincide and we get a *unique* isomorphism of  $\mathbb{Z}_\ell$ -modules (of maps to  $\mathbb{Z}_\ell$ )

$$\mathcal{M}(C(\tilde{k})) = \mathcal{M}(\mathbb{P}^1(\tilde{k})).$$

In particular, we find that  $g = 0$  and  $E = \tilde{k}(\mathbb{P}^1)$ . Further, we have an induced isomorphism

$$\mathbb{Z}_\ell\left(\sum_{w \in \mathcal{V}_E} \delta_w\right) = \mathbb{Z}_\ell\left(\sum_{w' \in \mathcal{V}_{k(\mathbb{P}^1)}} \delta_{w'}\right)$$

so that

$$\left(\sum_{w \in \mathcal{V}_E} \delta_w\right) = a \left(\sum_{w' \in \mathcal{V}_{k(\mathbb{P}^1)}} \delta_{w'}\right)$$

for some  $a \in \mathbb{Z}_\ell^*$ . This implies that  $\delta_w = a\delta_{w'}$ , for all  $w \in \mathcal{V}_E$  and the corresponding  $w' \in \mathcal{V}_{\mathbb{P}^1}$ . In particular, for the dual groups we have

$$E^*/\tilde{k}^* = (K^*/k^*)^a,$$

where  $a \in \mathbb{Z}_\ell^*$ . □

#### 14. $\ell$ -adic analysis: surfaces

sect:surr  
lemm:t-new

LEMMA 14.1. — *Let  $X$  be a smooth surface and  $D$  a smooth irreducible divisor on  $X$ . Let  $\nu = \nu_D$  be the corresponding divisorial valuation on the function field  $K = k(X)$  and  $t \in T_\ell(X)$  an element such that  $\text{r}\hat{\text{e}}\text{s}_\nu(t) = 1 \in \hat{K}_\nu^*$ . Then  $t$  is in the image of  $T_\ell(A)$  in  $T_\ell(X)$ .*

*Proof.* — Let  $B = B_D \subset A^0(X)$  be the abelian variety defined in Lemma 3.22 - the image of  $D$  in the quotient  $A = \text{Alb}(X)/B$  is a point; we have a map  $k(A) \rightarrow k(X)$ .

Let  $\alpha : \text{Alb}(X) \rightarrow A := \text{Alb}(X)/B$  projection. For any  $a \in A$  the preimage  $\alpha^{-1}(a)$  is isomorphic to the image of  $\text{Jac}(D)$  in  $\text{Alb}(X)$ . Since  $T_\ell(D)$

3.22 □

We will need an  $\ell$ -adic version of Lemma 11.1 .

prop:gff

PROPOSITION 14.2. — *Let  $\hat{f}, \hat{g} \in \mathcal{FS}_X(K)$  be elements with nontrivial support such that*

- $\varrho_\nu(\hat{f}, \hat{g}) = 1$  for every  $\nu \in \mathcal{DV}_K$ ;

$$- \operatorname{supp}_K(\hat{f}) \cap \operatorname{supp}_K(\hat{g}) = \emptyset,$$

that is,  $\hat{f}$  has nontrivial commutators. Then there is a 1-dimensional field  $E = k(C) \subset K$  such that  $\hat{f}, \hat{g} \in \hat{E}^*$ .

*Proof.* — By Lemma 12.4,

$$\hat{f} = t_f \cdot f, \quad \text{where } f := \prod_{i \in I} f_i^{a_i}, \quad \text{resp. } \hat{g} = t_g \cdot g, \quad \text{where } g := \prod_{j \in J} g_j^{b_j},$$

where

- $t_f, t_g \in T_\ell(X)$ ;
- $I, J$  are finite sets;
- $f_i, g_j \in K^*$  for all  $i, j$ ;
- $a_i \in \mathbb{Z}_\ell$  (resp.  $b_j \in \mathbb{Z}_\ell$ ) are linearly independent over  $\mathbb{Z}$ .

Fix a valuation  $\nu = \nu_D$ , where  $D$  is in the support of  $\hat{g}$  on a (smooth) model  $X$ . By assumption

$$\operatorname{res}_\nu(t_f \cdot \prod_{i \in I} f_i^{a_i}) = 1 \in \hat{K}_\nu^*.$$

By Lemma 12.3,  $\operatorname{res}_\nu(t_f) \in T_\ell(\hat{K}_\nu)$  so that  $t_f$  has trivial support on  $D$ . We claim that for all  $i \in I$ ,  $\operatorname{res}_\nu(f_i) = 1 \in \hat{K}_\nu^*/k^*$ . The divisor of the restriction of  $f_i$  to  $D$  is  $\sum_{i'} r_{ii'} q_{ii'}$ , where  $q_{ii'}$  are points on  $D$  and  $r_{ii'} \in \mathbb{Z}$ . This gives a relation

$$\sum_{i \in I} a_i \left( \sum_{i'} r_{ii'} q_{ii'} \right) = 0.$$

However,  $a_i$  were linearly independent over  $\mathbb{Z}$  which implies that  $r_{ii'} = 0$ , for all  $i, i'$ . In particular,  $\operatorname{res}_\nu(f_i) \in k^*$ . The same argument for  $g$  shows that  $g$  and  $f$  commute and that all pairs  $f_i, g_j$  commute as well. By Lemma 11.1, all  $f_i, g_j \in E = k(C) \subset K$  for some curve  $C$ .

We now have a diagram:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & T_\ell(E) & \longrightarrow & \hat{E}^* & \longrightarrow & \widehat{\operatorname{Div}}(C) & \longrightarrow & \widehat{\operatorname{Pic}}(C) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & T_\ell(K) & \longrightarrow & \hat{K}^* & \longrightarrow & \widehat{\operatorname{Div}}(X) & \longrightarrow & \widehat{\operatorname{Pic}}(X) & \longrightarrow & 0 \end{array}$$

We need to show that  $t_f$  (resp.  $t_g$ ) is in the image of  $T_\ell(E)$ . Let  $D$  be an irreducible component in the divisor of  $g$  (resp.  $f$ ). Changing the model, we may assume that  $D$  is smooth. We have a diagram

$$\begin{array}{ccccc} D & \longrightarrow & X & & \\ \downarrow & & \downarrow & & \\ \text{Jac}(D) & \xrightarrow{\iota} & \text{Alb}(X) & \xrightarrow{\alpha} & A = \text{Alb}(X)/B \end{array}$$

where  $\alpha$  is a surjection with connected fibers and  $B = B_D$  is as in Lemma 3.22: it is the minimal abelian subvariety of  $A^0(X)$  so that the image of  $D$  in  $\text{Alb}(X)/B$  is a point,  $a_D$ . (note that  $D$  is irreducible). We have

$$\iota(\text{Jac}(D)) = \alpha^{-1}(a_D) \simeq B.$$

Applying Lemma 12.1 (4) we conclude that the induced sequence

$$T_\ell(A) \xrightarrow{\alpha^*} T_\ell(\text{Alb}(X)) \xrightarrow{\iota^*} T_\ell(B)$$

of free finite rank  $\mathbb{Z}_\ell$ -modules is exact in the middle term. We have shown that  $\text{res}_\nu(f_i) = 1$ , for all  $i \in I$ . It follows that  $\text{r\acute{e}s}_\nu(t_f) = 1 \in \mathbf{K}_\nu^*$ , where  $\nu = \nu_D$  is the corresponding valuation. In particular,

$$t_f = 1 \in T_\ell(B) \hookrightarrow T_\ell(\mathbf{K}_\nu) = T_\ell(D).$$

It follows that there is an  $a \in T_\ell(A)$  such that  $\alpha^*(a) = t_f$ .

We apply this argument to every component  $D_j$  of the divisor of  $g$  and find that  $t_f$  is induced from quotients  $\text{Alb}(X)/B_j$ , where  $B_j := B_{D_j}$ , for  $j \in J$ . Let  $B$  be the abelian subvariety of  $A^0(X)$  generated by  $B_j$ . By Lemma 3.22,  $\text{Alb}(A)/B \simeq \text{Jac}(C)$ , and  $X$  maps to  $C$  with connected fibers. We have the diagrams

$$\begin{array}{ccc} X & \longrightarrow & \text{Alb}(X) \\ \downarrow & & \downarrow \\ C & \longrightarrow & \text{Jac}(C) \end{array} \quad \begin{array}{ccc} T_\ell(X) & \longleftarrow & T_\ell(\text{Alb}(X)) \\ \uparrow & & \uparrow \\ T_\ell(C) & \xleftarrow{\sim} & T_\ell(\text{Jac}(C)) \end{array}$$

It follows that  $t_f$ , and similarly  $t_g$ , is in  $T_\ell(C) = T_\ell(E)$ . □

rem:sati

REMARK 14.3. — For every  $f \in K^*$  the element  $g = (f + a)(f + b)$  where  $a \neq b$  and  $ab \neq 0$ , satisfies the conditions of Proposition 14.2.

Proposition 14.2 characterizes Galois-theoretically subgroups  $\hat{E}^* \subset \hat{K}^*$  corresponding to 1-dimensional subfields of  $K$ . We now have:

prop:surfa0

PROPOSITION 14.4. — *The group  $K^*/k^* \otimes \mathbb{Z}_\ell \subset \mathcal{FS}_X(K)$  is generated by subgroups  $\hat{E}^* \cap \mathcal{FS}_X(K)$  with  $E = k(C)$  so that  $T_\ell(E)$  is trivial. Moreover,*

$$(K^*/k^* \otimes \mathbb{Z}_\ell) \cap T_\ell(X) = 1.$$

*Proof.* — First of all  $T_\ell(C) \hookrightarrow T_\ell(X)$ , for every 1-dimensional subfield  $k(C) \subset K$ . It suffices to note that normally closed fields  $E = k(x)$ , with  $T_\ell(E)$  trivial, generate  $K^*/k^*$ .  $\square$

prop:surfa

PROPOSITION 14.5. — *Let  $\mathfrak{K}^* \subset K^*/k^* \otimes \mathbb{Z}_\ell \subset \hat{K}^*$  be a subset with the following properties:*

- $\mathfrak{K}^*$  is closed under multiplication;
- $\mathfrak{K}^* \cap \hat{E}^* = a_E \cdot E^*/k^*$  for every 1-dimensional normally closed subfield  $E = k(x) \subset K$ , with  $a_E \in \mathbb{Z}_\ell^*$ ;
- there exists a  $\nu_0 \in \mathcal{DV}_K$  such that

$$\{[\delta_0, \hat{f}] \mid \hat{f} \in \mathfrak{K}^*\} \simeq \mathbb{Z}$$

*for a topological generator  $\delta_0$  of  $\mathcal{I}_{\nu_0}^a$ .*

*Then  $\mathfrak{K}^* \subset K^*/k^* \otimes \mathbb{Z}_{(\ell)}$ .*

*Proof.* — For  $x \in K \setminus k$  let  $E = k(x)$  be the corresponding 1-dimensional field, assumed to be normally closed in  $K$ . By assumption, there exists an  $a_E \in \mathbb{Z}_\ell^*$  such that

$$\mathfrak{K}^* \cap \hat{E}^* = a_E \cdot E^*/k^*.$$

If some (any) topological generator  $\delta_0$  of  $\mathcal{I}_{\nu_0}^a$  is not identically zero on  $\hat{E}^*$  then there exists a (smooth) model  $X$ , where  $\nu_0$  is realized by a divisor  $D_0$ , together with a morphism

$$X \rightarrow \mathbb{P}^1 = \mathbb{P}_E^1$$

such that  $D_0$  dominates  $\mathbb{P}^1$ . It follows that

$$a_E \in \mathbb{Q} \cap \mathbb{Z}_\ell^* = \mathbb{Z}_{(\ell)}.$$

It remains to observe that every  $x \in K^*$  can be written as a product

$$x = x' \cdot x''$$



such that  $\delta_0$  is nontrivial on both normally closed  $E' = k(x')$  and  $E'' = k(x'')$ .  $\square$

coro:pm

**COROLLARY 14.6.** — *After a choice of  $\delta_0$ , for every 1-dimensional  $E \subset K$  and every  $f \in E^*/k^*$  we can Galois-theoretically distinguish its poles from its zeroes.*

The last essential step is a Galois-theoretic characterization of the partial projective structure on  $\mathfrak{K}^*/k^*$ , more precisely, the characterization of generating elements and primary lines in  $\mathfrak{K}^*/k^*$  (see Definition 3.10 and Definition 4.7).

lemm:unnn

**LEMMA 14.7.** — *Let  $x \in K^*$  be a generating element,  $E := k(x)$  and  $r = r(x) \in \mathbb{N}$  the smallest positive integer such that  $x^r \in \mathfrak{K}^*$ . Then*

- $r = p^m$  for some  $m \in \mathbb{N}$  (with  $p = \text{char}(k)$ );
- $(E^*/k^*) \cap (\mathfrak{K}^*/k^*) = (E^{p^m})^*/k^*$ ;
- (pointwise)  $p^m$ -th powers of primary lines in  $E^*/k^*$  coincide with primary lines in  $(E^{p^m})^*/k^*$ .

*Proof.* — The first property follows since  $K/\mathfrak{K}$  is a finite purely inseparable extension, by Propositions 3.19 and 14.5. Next, we claim that a generator  $y \in \mathfrak{K}$  is a  $p^m$ -th power of a generator of  $K$  (for some  $m$  depending on  $y$ ). Indeed,  $E := \overline{k(y)}^K \subset K$  is a finite and purely inseparable extension of  $k(y)$ ,  $E := k(x)$  (for some  $x \in K$ ). Thus

$$y = (ax^{p^m} + b)/(cx^{p^m} + d) = ((a'x + b')/(c'x + d'))^{p^m}$$

for some  $m \in \mathbb{Z}$ ,  $a, b, c, d \in k$  and their  $p^m$ -th roots  $a', b', c', d' \in k$  (since  $k$  is algebraically closed).

In particular, a generator  $y \in \mathfrak{K}^*$  is in  $E^* \cap \mathfrak{K}^*$  (and is the minimal positive power of a generator in  $E$  contained in  $E^* \cap \mathfrak{K}^*$ ). This implies the third property: the generators of  $E^{p^m}$  are  $p^m$ -th powers of the generators of  $E$ .  $\square$

coro:pro-uni

**COROLLARY 14.8 (Definition).** — *Assume that  $y, y'$  are primitive elements in  $(E^{p^m})^* \subset \mathfrak{K}^*$  such that*

- $y, y'$  have support in 2 points;
- the pole of  $y$  coincides with the pole of  $y'$ .

*Then (the images of)  $y, y'$  in  $\mathfrak{K}^*/k^*$  are contained in a primary line passing through (the images of)  $1, y, y'$ .*

*Proof.* — Definition 10.1 and Lemma 10.2 give a Galois-theoretic characterization of the notion “support in 2 points”. By Corollary 14.6 we can Galois-theoretically distinguish zeroes and poles of  $y \in \mathfrak{K}^*/k^*$ . It remains to apply Lemma 14.7.  $\square$

sect:proof

## 15. Proof

In this section we prove our main theorem: if

$$(\mathcal{G}_K^a, \Sigma_K) = (\mathcal{G}_L^a, \Sigma_L),$$

where  $L$  is a function field over an algebraic closure of a finite field of characteristic  $\neq \ell$ , then  $K$  is a purely inseparable extension of  $L$ .

*Step 1.* We have a nondegenerate pairing

$$\mathcal{G}_K^a \times \hat{K}^* \rightarrow \mathbb{Z}_\ell(1).$$

This implies that  $\hat{K}^* = \hat{L}^*$ .

*Step 2.* We have  $\Sigma_K^{\text{div}} = \Sigma_L^{\text{div}}$  and we identify intrinsically the inertia and decomposition groups of divisorial valuations:

$$\mathcal{I}_\nu^a \subset \mathcal{D}_\nu^a \subset \mathcal{G}_K^a :$$

every liftable subgroup  $\sigma \in \Sigma_K^{\text{div}} \subset \Sigma_K$  contains an inertia element of a divisorial valuation (which is also contained in at least one other  $\sigma' \in \Sigma_K$ ). The corresponding decomposition group is the “centralizer” of the (topologically) cyclic inertia group (the set of all elements which “commute” with inertia). This identifies  $\mathcal{DV}_K = \mathcal{DV}_L$ .

*Step 3.* For every  $\nu \in \mathcal{DV}_K$  we characterize intrinsically

$$\mathcal{I}_w^a \subset \mathcal{D}_\nu^a / \mathcal{I}_\nu^a$$

(see Proposition 10.3).

*Step 4.* We distinguish divisorial valuations with nonrational centers (see Lemma 10.4 and Remark 10.5).

*Step 5.* For  $\hat{f} \in \hat{K}^*$  we have two notions of support:  $\text{supp}_K(\hat{f})$  (intrinsic) and  $\text{supp}_X(\hat{f})$  (depending on a model  $X$ ) and two notions of finiteness:  $\hat{f}$  is nontrivial on at most finitely many nonrational divisorial valuations  $\nu$ , resp.  $\hat{f}$  has finite divisorial support on a model. We defined  $\mathcal{FS}(K) \subset \hat{K}^*$  as the set of elements satisfying the first notion of finiteness. If some (any) model  $X$  of  $K$  contains only finitely many rational curves, both notions of finiteness of support coincide and one obtains an intrinsic Galois-theoretic characterization of  $K^*/k^* \otimes \mathbb{Z}_\ell \subset \hat{K}^*$ , as elements in  $\mathcal{FS}(K)$ . In general, it may happen that some  $g \in L^*/l^*$  has an “infinite rational tail” on some (every) model  $X$  of  $K$ :

$$\rho_X(g) = \rho_{X,\text{nr}}(g) + \sum_{j \geq 1} n_j C_j,$$

where  $C_j$  are irreducible rational curves on  $X$ . In Lemma 12.10 we show that many elements of  $L^*/l^* \subset \mathcal{FS}(L) = \mathcal{FS}(K)$  have finite support on every model  $X$  of  $K$ , and vice versa. In particular, we prove that

$$\mathcal{FS}_0(K) = K^*/k^* \otimes \mathbb{Z}_\ell = L^*/l^* \otimes \mathbb{Z}_\ell$$

(up to an  $\ell$ -torsion group related to  $\text{Pic}^0(X)$ , for some model  $X$  of  $K$ ), where  $\mathcal{FS}_0(K) \subset \mathcal{FS}(K) \subset \hat{K}^*$  has an intrinsic Galois-theoretic description.

*Step 6.* For every pair of elements  $\hat{f}, \hat{g} \in \mathcal{FS}_0(K)$  satisfying

- $\text{supp}_K(\hat{f}) \cap \text{supp}_K(\hat{g}) = \emptyset$ ;
- $\varrho_\nu(\hat{f}, \hat{g}) = 1$  for all  $\nu \in \mathcal{DV}_K$

there exists a subfield  $E = k(C) \subset K$  such that  $\hat{f}, \hat{g} \in \hat{E}^*$  (Proposition 14.2).

*Step 7.* Proposition 13.1 identifies  $E^*/k^*$  inside  $\hat{E}^*$ , up to conformal equivalence, for all one-dimensional  $E = k(x)$ , which are integrally closed in  $K$ .

*Step 8.* Proposition 14.5 identifies  $\mathfrak{K}^* := K^*/k^* \cap L^*/l^*$  (as a multiplicative group) with a multiplicative subgroup of  $K^*/k^* \otimes \mathbb{Z}_\ell$ .

*Step 9.* By Proposition 3.19,  $\mathfrak{K}^*$  is isomorphic to  $K_1^*/k^*$ , and  $L_1^*/l^*$ , where  $K_1/K$  and  $L_1/L$  are finite purely inseparable extensions. Therefore,  $\mathfrak{K}^*$  carries two structures of an abstract projective space compatible with the multiplicative structure (see Example 4.5).

*Step 10.* By Theorem 4.6 the field is uniquely determined by the partial projective structure on  $\mathfrak{K}^*$  consisting of primary lines (see Lemma 4.8 and Lemma 4.9).

*Step 11.* Lemma 14.7 and Corollary 14.8 give a Galois-theoretic characterization of generating elements and primary lines in  $\mathfrak{K}^*$ . These define a (unique) partial projective structure on  $\mathfrak{K}^*$  (in particular, the projective structures induced by  $\mathbb{P}(K_1)$  and  $\mathbb{P}(L_1)$  coincide). In particular, the fields  $K_1$  and  $L_1$  both contain  $k$  and are isomorphic.

*Step 12.* It follows that  $K$  and  $L$  are finite purely inseparable extensions of the same field. This concludes the proof of Theorem 1.

## References

- [B-1] [1] F. A. BOGOMOLOV – “Abelian subgroups of Galois groups”, *Izv. Akad. Nauk SSSR Ser. Mat.* **55** (1991), no. 1, p. 32–67.
- [B-3] [2] F. A. BOGOMOLOV – “On two conjectures in birational algebraic geometry”, in *Algebraic geometry and analytic geometry (Tokyo, 1990)*, ICM-90 Satell. Conf. Proc., Springer, Tokyo, 1991, p. 26–52.
- [BT] [3] F. A. BOGOMOLOV and Y. TSCHINKEL – “Commuting elements in Galois groups of function fields”, in *Motives, Polylogarithms and Hodge theory*, International Press, 2002, p. 75–120.
- [bourb] [4] N. BOURBAKI – *Commutative algebra. Chapters 1–7*, Elements of Mathematics, Springer-Verlag, Berlin, 1998, Translated from the French, Reprint of the 1989 English translation.
- [efrat] [5] I. EFRAT – “Construction of valuations from  $K$ -theory”, *Math. Res. Lett.* **6** (1999), no. 3-4, p. 335–343.
- [endler] [6] O. ENDLER – *Valuation theory*, Springer-Verlag, New York, 1972, To the memory of Wolfgang Krull (26 August 1899–12 April 1971), Universitext.
- [mihalek72] [7] R. J. MIHALEK – *Projective geometry and algebraic structures*, Academic Press, New York, 1972.
- [mochizuki] [8] S. MOCHIZUKI – “The local pro- $p$  anabelian geometry of curves”, *Invent. Math.* **138** (1999), no. 2, p. 319–423.

- pop

 [9] F. POP – “On Grothendieck’s conjecture of birational anabelian geometry”, *Ann. of Math. (2)* **139** (1994), no. 1, p. 145–182.
- zs

 [10] O. ZARISKI and P. SAMUEL – *Commutative algebra. Vol. II*, Springer-Verlag, New York, 1975, Reprint of the 1960 edition, Graduate Texts in Mathematics, Vol. 29.